

Research Article

Radioprotective and *In-Vitro* Cytotoxic Sapogenin from *Euphorbia neriifolia* (Euphorbiaceae) Leaf

Papiya Bigoniya*¹ and Avtar Chand Rana²

¹Radharaman College of Pharmacy, Radharaman Group of Institutes, Ratibad, Bhopal, M.P. 462002,

²B.N College of Pharmacy, B.N Group of Colleges, Udaipur, Rajasthan, India.

Abstract

Purpose: *Euphorbia neriifolia* Linn. (Euphorbiaceae) plant is traditionally used in the treatment of abdominal troubles, bronchitis, tumours, leucoderma, piles, inflammation, and enlargement of spleen. The objective of this study was to evaluate the antioxidant and anticancer activities of a sapogenin isolate of this plant.

Methods: Euphol was isolated as a major constituent from the triterpenoidal sapogenin fraction of *E. neriifolia* leaf. Its *in-vitro* antioxidant activity was evaluated by reducing power assay, 1,1 – diphenyl – 2-picryl hydrazyl (DPPH) assay, as well as hydroxyl radical and superoxide radical scavenging activities. Radioprotective activity was assessed against radiation-induced chromosomal aberrations and cytotoxicity on murine F₁ B16 melanoma.

Results: The sapogenin exerted moderate antioxidant activity with highly significant ($p < 0.001$) reduction of gamma radiation-induced chromosomal aberrations (33.5 % compared to 71.5 % for radiation treatment alone at 4 Gy). It also exhibited cytotoxic activity on melanoma cell lines ($IC_{50} = 173.78 \mu\text{g/ml}$).

Conclusion: The sapogenin fraction showed antioxidant, radioprotective and cytotoxic activities. This study provides a scientific basis for the claimed traditional anticarcinogenic potentials of *E. neriifolia*.

Keywords: *Euphorbia neriifolia*; Euphol; Sapogenin; Antioxidant; Radioprotective; Melanoma; Chromosomal aberration

Received: 25 June 2009

Revised accepted: 11 September 2009

*Corresponding author: E-mail: p_bigoniya2@hotmail.com; Tel: +91-0755-2477941 (Res), 0755-2896237 (Inst); Fax: +91-0755-2896663

INTRODUCTION

Euphorbia neriifolia Linn, belonging to the family, Euphorbiaceae, is found throughout the Deccan Peninsula of India and commonly occurs in the dry hilly rocky grounds of north, central and south India. It is a herb full of spine, and is popularly known as sehund or thohar in Hindi. Ayurveda describes the plant as bitter, pungent, laxative, carminative, improves appetite, as well as useful in abdominal problems, bronchitis, tumours, loss of consciousness, delirium, leucoderma, piles, inflammation, enlargement of spleen, anaemia, ulcers and fever. Its leaves, in the Indian traditional system, are used as aphrodisiac, diuretic, and also in cough and cold, bleeding piles and ano-rectal fistula [1].

Plants of euphorbia species show anticarcinogenic activity due to the presence of several terpenes, anthocyanins, alcohols and steroids; diterpenoid ingenol 3,20-dibenzoate and phorbol 12-tiglate 13-decanoate isolated from Euphorbiaceae plants show antileukaemic activity against the P-388 lymphocytic leukaemia in mice [2]. Euphol, a triterpene alcohol from the roots of *Euphorbia kansui*, has inhibitory activity against mice skin tumour [3].

E. neriifolia, being widely available in large quantities, is potentially a low-cost source of active therapeutic substances. We have previously reported on the mild CNS depressant, wound healing and immunomodulatory activities of the hydro-alcohol leaf extract [4-6]. Little phytopharmacological work, however, has been done on the medicinal application of the leaf. Saponin isolated from the leaf possesses good haemolytic and *in-vitro* antioxidant activity but it is devoid of antibacterial activity up to 10 mg/ml concentration [7]. Since ethnopharmacological exploration has shown the traditional use of *E. neriifolia*, especially its leaf, as antitumour agent, the objective of this study was to isolate sapogenin from the

plant's leaf and study its antioxidant and anticancer activities.

EXPERIMENTAL

Plant material

E. neriifolia leaves were collected from cultivated field hedge plants in the suburban areas of Bhopal (latitude 23.21°, longitude 77.84°, BHOP), Madhya Pradesh, India, in September 2005. The plant was identified with the aid of available literature and authenticated by Dr AP Shrivastava, a taxonomist and Principal, P.K.S Govt. Ayurveda College and Institute, Bhopal, India. A voucher specimen (no. 1085) was deposited in the herbarium of the department.

Reagents

RPMI 1640 media, foetal calf serum and 1,1 – diphenyl –2-picryl hydrazyl (DPPH) were purchased from Sigma Chemicals Co., St. Louis, USA while phytohaemagglutinin was obtained from Difco, USA. Collagen-coated culture flasks were purchased from Nunc, Denmark. and Neubauer hemocytometer from Feinoptik, Germany. Modified Eagle Minimum Essential Media and deoxyribose were purchased from Himedia, Mumbai; vincristin from Cipla, India, nitroblue tetrazolium from E. Merck, Darmstadt, Germany. Photomicrographs were captured using an Olympus D×60 microscope connected to an Olympus DP-50 digital camera.

Extraction, isolation and characterization of compound

One kilogram of the dried powder of the leaf was extracted with 3 L of cold ethanol (70 %) by maceration for seven days and the solvent removed under vacuum. Phytochemical investigation of the extract was performed to detect the presence of reducing sugar, tannin flavonoid, alkaloid, saponin, steroid, glycoside and fixed oil [8,9]. The extract was

re-suspended in 250 ml of water and 500 ml of chloroform in HCl (50 %v/v) was added to effect acid hydrolysis of its saponins content in order to isolate sapogenins. The chloroform phase was separated and concentrated at < 40 °C to a third of its volume. This phase was exhaustively extracted with water-saturated n-butanol (three times) and the solvent removed under vacuum. The brown dried powder obtained represents the total crude sapogenin (yield: 2.41 %) and it tested positive according to Salkowski and Noller's test [10].

Characterisation of sapogenin isolate

The whole sapogenin isolate was subjected to column chromatography on silica gel using chloroform, a solvent gradient of chloroform/ethyl acetate (80:20, 60:40, 40:60 and 20:80), ethyl acetate and methanol. Five fractions were collected and chromatographed on silica gel G plates using CHCl₃ : MeOH (50:50). Fraction 3 was subjected to other assessments including Salkowski, Noller's and Libermann Burchard tests [10]. Its melting point was determined from its DSC thermogram using a Mettler Toledo DSC 821 system in which the sample press-sealed in an aluminium pan with a perforated lid and heated at a rate of 5 °C/min in a nitrogen environment. The UV spectra of 5 µg/ml sapogenin fraction in chloroform were obtained with a Shimadzu (UV-1700 Pharmaspec) spectrophotometer while the IR spectra were recorded on a Shimadzu (Jasco FTIR-5300) spectrometer using KBr pellet at a scanning speed of 2 mm/sec and with resolution set at 4 cm⁻¹. NMR spectra were determined with a Bruker Deltonics (Avance 300) spectrometer in CDCl₃ at 300 MHz and the entire chemical shifts were relative to tetramethyl silane (TMS, δ 0.00). The electron-impact mass spectra of the powdered sample was recorded on Shimadzu (QP 5000) spectrometer in CHCl₃, injected in HT-8 column, using helium as the carrier gas at a heating rate of 15 °C/min and scan rate of 1 scan/sec in a scan range of 100-500 Delton.

Preparation of extract stock solutions

A stock solution of the total sapogenin (50 mg/ml) was prepared in dimethyl-sulfoxide (DMSO) and the volume made up to 1000 ml (to obtain a 100 µg/ml concentration) with RPMI media (for blood culture) or Eagle's Modified MEM (for melanoma cell culture). It was then sterilized by filtration through a 0.2 µm membrane filter. Other concentrations, ranging from 40 – 75 µg/ml, were prepared by dilution. The final DMSO concentration of extract media combination was 0.25 %v/v and at this level, no growth inhibitory effects were observed.

***In-vitro* antioxidant activities**

In-vitro antioxidant activities of the total sapogenin were measured in a concentration range of 50 – 1000 µg/ml, using various methods.

Reducing power assay

The reducing power assay (RPA) involves the oxidation-reduction reaction of potassium ferricyanide and ferric chloride in the presence of the antioxidants. The reducing power of 1 mg sample is equivalent (E) to reducing power of 1 nM ascorbic acid (AS) expressed as ASE/mg assayed following the method of Oyaizu [11]. Different concentrations of the sapogenin and 1nM of ascorbic acid in a final volume of 1ml were taken in different test tubes and mixed with 500 µl of potassium ferricyanide (1%). The mixture was incubated at 50 °C for 20 min. TCA (10%, 50 µl) was added and centrifuged at 600 rpm for 10 min. To the supernatant, 0.5 ml of ferric chloride (0.1%) was added and the absorbance of the resultant violet coloured solution was measured at 700 nm.

Hydrogen-donating ability

Hydrogen-donating ability (HDA) was measured as the amount of total sapogenin required for inhibiting the formation of 1,1-diphenyl –2-picryl hydrazyl (DPPH) radical by

50 % (IC₅₀ value) according to the method of Hatano et al [12]. Antioxidants react with methanolic DPPH (100 µM) and convert it to 1,1-diphenyl –2-picryl hydrazine. Different concentrations of the sapogenin (200 µl) was added to 3 ml of methanolic DPPH; 20 min later, the amount of DPPH remaining was measured spectrophotometrically at 520 nm against blank.

Hydroxyl radical scavenging ability

Hydroxyl radical scavenging ability (HRSA) was measured by evaluating the competition between deoxyribose and the test compounds for hydroxyl radicals generated from the Fe³⁺/ascorbate/EDTA/H₂O₂ system using the method of Elizabeth and Rao [13]. The reaction mixture contained 100 µl of deoxyribose, 50 µl of FeCl₃, 50 µl of EDTA, 100 µl of H₂O₂ and 100 µl of ascorbic acid. Different concentrations of the sapogenin were added to KH₂PO₄/KOH buffer (20 mM, pH 7.4) to make it up to 1 ml. Incubation for 1 h at 37 °C resulted in the formation of thiobarbituric acid reactive substances (TBARS) measured spectrophotometrically at 532 nm following the method of Ohkawa et al [14]. The results were expressed as percent inhibition of TBARS and the amount of sample (µg/ml) producing 50 % antilipid peroxidation was determined.

Superoxide radical production

The effect of total sapogenin on superoxide radical production (SRP) was evaluated following the nitroblue tetrazolium (NBT) reduction method of McCord and Fridovic [15]. The reaction mixture containing 0.5 mM xanthine as substrate (300 µL), 1 mM EDTA in phosphate buffer (100 µL), 0.05 mM sodium cyanide (100 µl), 0.5 mM xanthine oxidase (20 µL), different concentrations of the sapogenin (20 µl) and 0.1 mM cytochrome C (300 µL) was placed in a 1 cm cuvette and the rate of increase in absorbance at 550 nm was recorded every minute for 5 min. The results were calculated

as amount of dismutase required to inhibit the rate of reduction of cytochrome C by 50 % (i.e., to a rate of 0.0125 absorbance unit per minute which is defined as 1 unit of activity) and the amount of sample (µg/ml) producing 50 % reduction of cytochrome C was determined.

Effect on radiation-induced chromosomal aberrations in cultured human lymphocytes

Fresh whole blood (1.5 ml) was taken in different culture flasks and treated with 40, 55, 75 and 100 µg/ml of total sapogenin in four different sets. One flask in each set was taken as vehicle control and treated with 75 µl of vehicle. All the culture flasks were incubated for 30 min and then each separate set of culture flasks was exposed to 1, 2, 3 and 4 Gy of γ-radiation. Triplicate cultures for each individual flask were set up by mixing 0.5 ml of blood with 4.5 ml of RPMI 1640 media. The culture flasks were coded and incubated at 37 °C in a humidified atmosphere of 5 % CO₂. The cultures were harvested at 72 h for chromosomal studies. Air-dried preparations of hypotonically treated lymphocytes were made using routine techniques for chromosomal analysis as described by Moorhead et al [16]. Slides were prepared by air drying method and chromosomal aberrations were scored in conventional Giemsa-stained preparations as described by Lloyd et al [17].

Media for blood culturing

Blood samples were collected in heparinised sterile glass vials from the median cubital vein of a non-smoking healthy female donor of approximately 25 years. The whole blood (0.5 ml) was incubated with 5 ml of RPMI 1640 media containing 5 % heat-inactivated foetal calf serum, penicillin (100 U/ml), streptomycin (100 µg/ml) and L-glutamine (1.5 mg/ml). An optimum concentration of phytohaemagglutinin 5 µg/ml was used to stimulate the lymphocytes to transform and divide in culture [18].

In-vitro radiation

Theratron 780C cobalt teletherapy unit (Theratronics Limited, Canada) was used for radiating whole human blood samples at a dose rate of 1 Gy/min in culture flasks (25 cm³), in a field size of 7 × 20 cm² which can accommodate 5 flasks at a time at a S.S.D (source surface distance) of 90 - 95 cm.

In-vitro cytotoxicity assay of the extract on murine F₁ B16 melanoma cell line

Murine F₁ B16 melanoma cell line was used to study the *in-vitro* anti-cancer activity of total sapogenin. Male C 57 BL/6 mice were used for *in-vivo* maintenance of cell line. *In-vitro* harvesting of cells was carried out from a full-grown melanoma site (2.5 to 3 cm) by aseptic transfer to collagen-coated culture flasks in Modified Eagle Minimum Essential Media containing NaHCO₃ (220 mg/100ml), which was supplemented with 10 % heat-inactivated faetal calf serum, L-glutamine (300 µg/ml) and 48 mg gentamicin at pH 7.4. The cells were expended in 75 cm² tissue culture flasks at 37 °C at an atmosphere of 5 % CO₂ in air (100 % humidity). A confluent monolayer was detached with 0.1 % trypsin containing 0.02 % EDTA in Ca²⁺ and Mg²⁺ free PBS (pH 7.4, 0.01 M) and dissociated into a single-cell suspension for further cell culture. Serially cultured cell lines at P₃ to P₄ stages were used for cytotoxicity assay.

The assay was performed in collagen-coated radiation sterilised cell culture dishes (2 ml capacity, growth surface 60 × 15 mm). An amount of the cell culture (2 × 10⁴ cells/ml) was seeded and allowed to adhere by incubating for 6 h at 37 °C in 5 % CO₂. This method was standardised previously in the laboratory following the method of Umadevi et al [19]. After 6 h, the cells were exposed to different concentrations of total sapogenin for a further 6 h. The range of tested concentrations was from 10 to 500 µg/ml for total sapogenin and from 10 to 500 ng/ml for vincristin as positive control. Triplicate dishes were incubated for 72 h and cell viability was

checked every 24 h. The cells were detached by rinsing twice with trypsin, and then palleted by centrifugation; cell viability was measured by tryphan blue dye exclusion test and counted using WBC counting chamber of Neubauer's chamber. The IC₅₀ (concentration at which cellular growth is inhibited by 50 %) was determined at 72 h. The IC₅₀ values were estimated from a plot of log total sapogenin concentration against percent cell viability.

Ethical considerations

The experimental protocol for animal studies was approved by the Institutional Animal Ethical Committee (ref no. Animal Eths. Comm./DB/304) prior to carrying out the experiments and the animals were handled as per the 'WHO guidelines for the care and use of animals in scientific research'.

Statistical analysis

All data are presented as mean ± SEM. Experimental data were analysed using one-way ANOVA followed by Student's *t*-test. P < 0.05 was considered significant. Graph Pad Prism Version 3.02 software was used for statistical calculations.

RESULTS

Identification of compound

The leaf extract (yield: 10.8 %) was positive for reducing sugar, tannins, flavonoids, alkaloids, and triterpenoidal saponin but negative for glycoside and fixed oil. Hydrolysis of the extract followed by extraction with water-saturated n-butanol gave a brown coloured crude sapogenin mixture with a yield 2.4 %. Chromatographic elution of crude sapogenin with chloroform, chloroform/ethyl acetate (80: 20, 60: 40, 40: 60 and 20: 80), ethyl acetate and methanol produced five fractions. The fractions obtained were: Fraction-1 (mixture), Fraction-2 (ENS-1) with Rf value of 0.385, Fraction-3 (ENS-2) with Rf value of 0.360, Fraction-4

(no spot) and Fraction-5 (ENS-3) with Rf value of 0.314.

Fraction 3 was positive for Salkowski and Noller's test and negative for Libermann Burchard test, indicating the presence of triterpene. The residue obtained was carefully crystallized on methanol gives a solid, white crystal (232 mg), m.p. 116 °C. UV δ_{\max} 270 nm; IR (KBr, cm^{-1}): 3400 & 1030 (3- β -OH), 2923 & 2854, 1637 (-C=C-), 1461 & 1376, 925, 862, 802 & 723; ^1H NMR (CDCl_3 300 MHz): δ 5.43 (1H, t, H-24), 3.29 (1H, m, H-3 β) 1.62 (Me-26), 1.77 (Me-27), 0.74 (Me-18), 0.85 (Me-19), 0.91 (Me-28), 1.16 (Me-29), 1.21 (Me-30), 1.12 (Me-21); ^{13}C NMR (CDCl_3 75 MHz): δ 81.01 (C-3), 135.62 (C-8), 37.26 (C-20), 129.20 (C-24), 20.18 (C-21), 17.52 (C-18), 21.14 (C-19), 18.72 (C-26), 27.32 (C-27), 17.78 (C-28), 26.56 (C-29), 30.14 (C-30); EIMS m/z (%): 426 (M^+), 408 ($\text{M}^+ - \text{H}_2\text{O}$), 297 0.24 % $\text{M}^+ - \text{H}_2\text{O} - \text{C}_8\text{H}_{15}$) and other fragments suggesting the fragmentation of side chain 111 (16.42 % C_8H_{15}), 97 (25.08 % C_7H_{13}), 83 (42.96 % C_6H_{11}), 69 (70.93 % C_5H_9), 55 (66.15 % C_4H_7) and 41 (100 % C_3H_5). All the data's were compared with published data for Euphol from Kansui Radix as reported by Lin et al [20]. Thus fraction 3 was characterised as euphol (8, 24-Euphadien- 3 beta-ol; m.p. 116 °C; yield 0.0232%, fig. 1) based on IR and mass spectrometry.




Fig. 1: 8, 24-euphadien- 3 beta-ol

Antioxidant activity

The *in vitro* antioxidant activities of the total sapogenin are depicted in Table 1.

Sapogenin, in the amounts used, showed moderate to high free radical scavenging activity and good hydrogen donating ability in relation to α -tocopherol.

Table 1: Antioxidant activity of total sapogenin fraction of *E. neriifolia* leaf

Treatment	RPA ASE/mg	IC ₅₀ ($\mu\text{g/ml}$)		
		HDA (DPPH assay)	HRSA	SRP
α -tocopherol	447.70 \pm 6.21	38.6 \pm 0.7	34.2 \pm 0.4	59.6 \pm 1.0
Total sapogenin	190.36 \pm 2.86	440.6 \pm 42.3	806.8 \pm 57.9	471.3 \pm 38.2

n = 9 (experiments were repeated thrice, each time in triplicate). The IC₅₀ value was defined as the concentration ($\mu\text{g/ml}$) of the total sapogenin required for inhibiting the formation of free radicals by 50 %. Each experiment was carried out in triplicate and deviation of the absorbance value was less than 10 %.

Effect of the extract on radiation-induced chromosomal aberrations

Table 2 shows the effect of the extract on radiation-induced chromosomal aberrations of cultured human lymphocytes.




Fig 2: Dose-response curve for total chromosomal aberrations (%) in cultured human peripheral lymphocytes treated with radiation alone and after pre-treatment with 75 $\mu\text{g/ml}$ of *E. neriifolia* leaf total sapogenin (■ = radiation alone; ● = radiation + total sapogenin)

Table 2: Effect of total sapogenin of *E. neriifolia* leaf on radiation induced chromosomal aberrations in cultured human lymphocytes

Radiation dose (Gy)	Treatment ($\mu\text{g/ml}$)	No. of cells scored	Chromosomal aberrations (%)						
			Dicentric			Centric rings	Excess acentrics	Chromatid	
			Fragments	Without fragments	Inter-change			breaks	gaps
-	Control	470	1.4	0.0	0.0	0.0	2.5	0.0	0.4
-	75	500	1.6	0.2	0.0	0.0	2.8	0.0	0.5
1	Control	532	6.7	0.3	2.3	1.4	5.2	0.2	1.3
	40	586	5.2	0.3	2.1	1.2	4.6	0.2	1.2
	55	725	3.7	0.2	1.4	1.0	3.5	0.0	1.2
	75	658	3.2	0.0	0.9	0.6	3.1	0.0	0.8
	100	564	3.5	0.0	1.1	0.8	3.2	0.0	0.8
2	Control	536	9.4	0.6	4.5	2.8	8.5	0.6	1.9
	40	513	8.3	0.6	4.2	2.0	6.0	0.4	1.8
	55	672	5.5	0.4	3.2	1.7	5.3	0.2	1.4
	75	651	4.1	0.2	2.9	1.3	4.2	0.2	1.3
	100	543	4.2	0.2	3.2	1.3	4.4	0.2	1.4
3	Control	995	18.1	1.6	12.4	5.8	12.8	0.8	2.7
	40	578	12.3	1.4	9.8	4.1	10.4	0.5	2.2
	55	733	10.0	1.1	7.8	3.5	8.7	0.5	1.8
	75	680	9.9	0.7	7.0	2.6	8.0	0.3	1.5
	100	570	10.3	0.8	7.0	2.5	8.3	0.3	1.6
4	Control	520	24.2	2.7	18.6	7.9	15.1	1.8	4.3
	40	544	20.6	2.2	12.3	6.4	12.9	1.4	3.4
	55	659	13.6	1.5	9.0	4.1	8.3	0.9	2.3
	75	712	10.2	1.2	8.7	3.5	6.9	0.6	1.8
	100	375	12.3	1.3	8.6	3.7	7.2	0.6	2.0

Control samples were incubated without any treatment. These data show the effect of the leaf total sapogenin treatment (40 -100 $\mu\text{g/ml}$) as well as the effect of different doses of radiation (1 - 4 Gy) on chromosomal aberrations in cultured normal non-malignant cells (human lymphocytes).

Pre-treatment with 75 $\mu\text{g/ml}$ of total sapogenin fraction reduced total chromosomal aberrations to 33.5 % compared to 71.5 % for radiation treatment (RT) alone at 4 Gy. The slope of the linear dose-response curve for total sapogenin treatment was 7.83, which is significantly ($p < 0.001$) lower than that of the radiation only treated group (see Figure 2).

Cytotoxicity assay on murine F₁ B16 Melanoma cell line

In-vitro testing of total sapogenin against the murine F₁ B16 Melanoma cell line showed 76.6 % cell viability at 10 $\mu\text{g/ml}$ compared to

13.6 % at 500 $\mu\text{g/ml}$ of total sapogenin, with control as 100 % cell viability, as shown in Figure 3(A). Figure 3(B) shows the plot of log total sapogenin concentration against cell viability (probit scale) with a best-fit linear regression curve (slope: 0.9415) superimposed. The assay data show that the IC₅₀ (over a period of 72 h) concentration of total sapogenin that inhibited growth of mouse melanoma cells by 50 % was 173.78 $\mu\text{g/ml}$ compared to 120 ng/ml for vincristin.

DISCUSSION

Plants are valuable sources of novel biologically active molecules. Saponins are

high molecular weight compounds comprising glycosides with a sugar moiety linked to a triterpene or steroid aglycone. Triterpene saponins, particularly, have been the subject of much interest because of their biological properties.

Free radical damage to biosystems is one of the major processes that contribute to degenerative diseases such as cancer and ageing. Free radical scavengers protect cellular DNA against indirect effects of ionizing radiation where hydroxyl radicals are believed to be the primary active species responsible for the damage [21]. The data obtained in this study demonstrate the antioxidant activity of the sapogenin isolated from the leaf extract of *E. neriifolia*. The good reducing power of sapogenin means that triterpenoidal compounds, especially euphol, are electron donors, and therefore, can act as antioxidants [22]. Hydrogen donating ability is an index of primary antioxidants. DPPH is known to abstract labile hydrogen and the ability to scavenge the DPPH radical is related to the inhibition of lipid peroxidation [23]. Total sapogenin inhibits oxygen derived free radicals such as superoxides and hydroxyl radical *in vitro* with a relatively moderate potency. Terpenes and bioflavones isolated from *Ginkgo biloba* inhibited lipid peroxidase and superoxide anion in hepatocytes that are generally implicated in cell damage [24]. There is a good correlation between antioxidant properties and radioprotection by flavonoids as they could prevent the accumulation of DNA damage induced by UV radiation. Castilla et al [25] has demonstrated the antioxidant as well as radioprotective effects of flavon-3-ol from grape seeds against chromosomal damage induced by x-rays.

Peripheral blood lymphocytes (PBL) are extensively used in biomonitoring of populations exposed to various mutagenic or

carcinogenic compounds because the sensitivity of this system in detecting chromosome damage induced by exposures of ionising radiation. γ -radiation produces morphological changes in lymphocytes by decaying their proliferation, which indirectly defines genomic instability. γ -rays generate hydroxyl radicals in cells and induce DNA damage that leads to mutations and chromosomal aberrations [26]. Total sapogenin at a concentration of 75 $\mu\text{g/ml}$ significantly decreased total chromosomal aberration. The results signify that the sapogenins reduced gamma radiation-induced genomic instability by reducing chromosomal aberrations due to the presence of antioxidants.

Total sapogenin exhibits cytotoxic activity on murine F₁ B16 Melanoma cell line. It showed apparently high IC₅₀ as the cells were exposed to extracts only for a short duration of 6 h. Using conventional cytotoxic tests, the cells were exposed to cytotoxic chemical for 72 h where cytotoxicity may be partly due to cumulative accumulation of the drugs in the culture media.

Saponins have many kinds of biological activities such as anti-bacterial, anti-viral, anti-tumour, anti-fertility, anti-inflammatory, anti-hyperlipidemic, anti-hypertension, anti-hyperglycaemic and immunoregulatory, etc. Furthermore, triterpenoids exert physiological activities in the cardiovascular, nervous and, adrenocortical systems as well as enzymatic activity, and therefore, are frequently a subject of research in natural medicine. The total triterpenoidal sapogenin extracted from bamboo (*Phyllostachys Sieb. et Zucc*) were reported to exhibit pharmacological activities such as anti-free radical, anti-oxidation, anti-tumour and anti-hypertension [27]. Triterpene alcohols such as ursolic and oleanolic acids are said to exert antitumour effect on lung,




Figure 3: Viability (%) of F1 B16 melanoma cells after 72 h of *in-vitro* culture growth with different concentrations of *E. neriifolia* leaf total sapogenin (A); Inhibitory profile of *E. neriifolia* leaf total sapogenin treatment on *in-vitro* growth of F1 B16 melanoma cells after 72 h of culture (B). IC₅₀ = 173.78 µg/ml (antilog of 2.24).

breast and colon tumours [28]. Topical application of euphol isolated from roots of *Euphorbia kansui* markedly suppressed the tumour-promoting effect of 12-O-tetradecanoylphorbol-13-acetate in two-stage carcinogenesis in mouse skin initiated with 7, 12-dimethylbenz[a] anthracene [3].

CONCLUSION

E. neriifolia Leaf is rich in crude sapogenin, and euphol (0.023 %) was identified as a major constituent. The sapogenin fraction showed antioxidant, radioprotective and cytotoxic activity against malignant melanoma cells. Our study supports the use of *E. neriifolia* as an antitumour herbal remedy in Indian traditional medicine. This study reports for the first time the potential anticancer properties of *E. neriifolia* triterpenes. However, the sapogenin content needs to be studied further to isolate other active constituents and to elucidate its *in-vivo* anticancer activity profile; this further work is in progress in our laboratory.

ACKNOWLEDGEMENT

The authors are grateful to All India Council of Technical Education, New Delhi, for financial assistance through the award of a National Doctoral Fellowship to one of the authors (PB) (grant no. FD/NDFS/2003-04). We also express our gratitude to Drs P Uma Devi and N Ganesh, and all the research staff of Jawaharlal Nehru Cancer Hospital and Research Center, Bhopal, for their assistance, support and valuable suggestions.

REFERENCES

1. Anonymous. *The Wealth of India: Raw Material*. Vol. III (D-E), New Delhi, Council of Scientific and Industrial Research Publication, 1952, pp 226.
2. Kupchan SM, Uchida I, Branfman AR, Dailey RG Jr, Fei BY. Antileukemic principles isolated from euphorbiaceae plants. *Science* 1976; 191(4227): 571-572.
3. Yasukawa K, Akihisa T, Yoshida ZY. Inhibitory effect of euphol, a triterpene alcohol from the roots of *Euphorbia kansui*, on tumor promotion by 12-o-tetradecanoylphorbol-13-acetate in two-stage carcinogenesis in mouse skin. *J Pharm Pharmacol.*, 2002; 54(1): 119-124.

4. Bigoniya P, Rana AC. Psychopharmacological profile of hydro-alcoholic extract of *Euphorbia neriifolia* leaves in mice and rats. *Indian J Exp Biol.*, 2005; 43: 859-862.
5. Bigoniya P, Rana AC. Wound healing activity of *Euphorbia neriifolia* leaf extract. *J Nat Remedies* 2007; 7(1): 94-101.
6. Bigoniya P, Rana AC. Immunomodulatory activity of *Euphorbia neriifolia* leaf hydro-alcoholic extract in rats. *Ind Drugs* 2008; 45(2): 90-97.
7. Bigoniya P, Rana AC. Hemolytic and In-vitro Antioxidant Activity of Saponin Isolated from *Euphorbia neriifolia* Leaf. In: *Recent Progress in Medicinal Plants, Natural Products – II*, vol 2, Chapter-20, 2006, pp 359-376.
8. Brain KR, Turner TD. *The Practical Evaluation of Phytopharmaceuticals*. Bristol, Wright-Scientenica, 1975, pp 153.
9. Shellard EJ. *Practical Plant Chemistry for Pharmacy Students*. London, Pitman Medical Publishing Co. Ltd., 1957, pp 34-80.
10. Hawk PB, Oster BL, Summerson WH. *The Practical Physiological Chemistry*. 13th ed, New York, McGraw Hill Book Co., 1954, pp 111.
11. Oyaizu M. Studies of products browning reaction: Antioxidative activity of products of browning reaction prepared from glucosamine. *Jpn J Nutr.*, 1986; 44: 307-315.
12. Hatano T, Kagawa H, Yasuhara T, Okuda T. Two new flavonoids and other constituents in cicore root: their relative astringency and radical scavenging effects. *Chem Pharm Bull.*, 1988; 36: 2090-2097.
13. Elizabeth K, Rao MNA. Oxygen radical scavenging activity of curcumin. *Int J Pharm.*, 1990; 58: 237-240.
14. Ohkawa H, Ohishi N, Yagi K. Assay for lipid peroxidation in animal tissue by thiobarbituric acid reaction. *Anal Biochem.*, 1979; 95: 351-358.
15. McCord JM, Fridovich I. Superoxide dismutase: An enzymic function for erythrocyte (hemocuprein). *J Biol Chem.*, 1969; 244: 6049-6055.
16. Moorhead PS, Nowell PC, Meliman WJ, Battip PMC, Hungerford DA. Chromosome preparations of leukocytes cultivated from human peripheral blood. *Exp Cell Res.*, 1960; 20: 613-617.
17. Lloyd DC, Purrott RJ, Prosser JS. The study of chromosomal aberration yield in human lymphocytes as an indicator of radiation dose: Revised Technique, National Radiological Protection Board. Technical Memorandum NRPB-M70: Chilton, Didcot, Oxon, Great Britain, 1982.
18. Fenech M, Morley AA. Cytokinesis-block micronucleus method in human lymphocytes: effect of in vivo ageing and low dose x-irradiation. *Mutation Res.*, 1986; 161: 193-198.
19. Uma Devi P, Nagarathnam A, Rao BSS. *Introduction to radiation biology*. New Delhi, B.I Churchill Livingstone Pvt. Ltd., 2000, pp 226-227.
20. Lin JH, Ku YR, Lin YZ, Teng SF, Wen KC, Liao CH. Preparative isolation and gas chromatography-mass spectrometry analysis of triterpenoids in *kansui radix*. *J Food Drug Anal.*, 2000; 8(4): 278-282.
21. Sasaki MS, Matsubara S. Free radical scavenging in protection of human lymphocytes against chromosome aberration formation by gamma-ray irradiation. *Int J Radiat Biol.*, 1977; 32: 439-445.
22. Yen GC, Chen HY. Antioxidant activity of various tea extracts in relation their antimutagenicity. *J Agric Food Chem.*, 1995; 43: 27-32.
23. Matsubara N, Fuchimoto S, Iwagaki H, Nonaka Y, Kimura T, Kashino H, Edamatsu R, Hiramatsu M, Orita K. The possible involvement of free radical scavenging properties in the action of cytokines. *Res Commun Chem Pathol Pharmacol.*, 1991; 71: 239-242.
24. Joyeux M, Lobstein A, Anton R, Mortier F. Comparative antilipidperoxidant, antinecrotic and scavenging properties of terpenes and bioflavones from Ginko and flavonoids. *Planta Med.*, 1995; 61: 126-129.
25. Castilla J, Benavente-García O, Lorente J, Alcaraz M, Redondo A, Ortuno A, Del-Río JA. Antioxidant activity and radioprotective effects against chromosomal damage induced in vitro by X-rays of flavan-3-ols (procyranidins) from grape seeds (*Vitis vinifera*): comparative study versus other phenolic and organic compounds. *J Agri Food Chem.*, 2000; 48: 1738-1745.
26. Riley PA. Free radicals in biology: oxidative stress and the effects of ionizing radiation. *Int J Radiat Biol.*, 1994; 65: 27-33.
27. Ying Z, Xiaoqin W, Zhuoyu Y, Yunlong Z, Lingen C, Shenggen L. Composition containing total triterpenoid saponins extracted from bamboo, and the preparation method and use thereof, United States Patent 20060148733, 2006. <http://www.freepatentsonline.com/y2006/0148733.html>. Accessed 2nd May, 2007.
28. Haridas V, Gutterman JU. Triterpene compositions and methods for use thereof, United States Patent 6962720, 2002. <http://www.freepatentsonline.com/6962720.html>. Accessed 7th August, 2007.