

## COMBINING ABILITY FOR QUALITY OF MAIZE FLOUR IN A 6-PARENT DIALLEL CROSS

J.E. ALIKA

Department of Crop Science, Faculty of Agriculture  
University of Benin, Benin City, Nigeria

(Received 4 October 1993 ; accepted 4 January 1994 )

### ABSTRACT

Whole maize (*Zea mays L.*) grains from 15 F<sub>1</sub> hybrids obtained from a diallel cross of six maize cultivars consisting of one waxy and five non-waxy were ground into flour and used to prepare “tuwo”, a thick porridge. “Tuwo” was prepared by heating a mixture of 10 g flour in 30 ml of water and evaluated for firmness, stickiness and time of gelatinization. Mean square for general combining ability (GCA) was highly significant for stickiness and firmness of “tuwo”. Gelatinization time was highly significant for specific combining ability (SCA). Additive gene effects were substantially more important for firmness and stickiness, while gelatinization time was considerably influenced by non-additive genes. The waxy maize contributed more unfavourable GCA effects for softness and stickiness of “tuwo” than the remaining five cultivars.

**Key Words:** Firmness, gelatinization time, general combining ability, specific combining ability, stickiness, “tuwo”.

### RÉSUMÉ

Les grains complets du maïs (*Zea mays L.*) provenant de 15 hybrides F<sub>1</sub> obtenus par croisement dialèle de 6 variétés dont un non farineux et 5 farineux sont réduits en farine pour la confection du “tuwo”, une bouillie épaisse. Le tuwo, préparé en chauffant la mixture de 10 g de farine dans 30 ml d'eau a été évalué pour la fermeté, la viscosité et le temps de gelatinisation. La carré moyenne pour une aptitude de combinaison générale (GCA) a été hautement significative quant à la viscosité et la fermeté de “tuwo”. Le temps de gelatinisation a été hautement significatif pour une aptitude de combinaison spécifique (SCA). Les effets des gènes additifs sont plus importants pour la fermeté et la viscosité, tandis que le temps de gelatinisation a été largement influencé par les gènes non additifs. Le maïs non farineux a contribué d'une manière défavorable aux effets GCA pour la douceur et la viscosité du “tuwo” que les 5 autres cultivars non farineux.

**Mots Clés:** fermeté, temps de gelatinisation, aptitude de combinaison générale, aptitude de combinaison spécifique, viscosité, tuwo

## INTRODUCTION

“Tuwo” is a local name to a dry milled cereal flour gel product which is a common food with only slight regional modifications from Senegal to Ethiopia (Scheuring *et al.*, 1982). In Nigeria, “tuwo” made from sorghum has been consumed more than that made from maize among people living in the north of the country, until recently when maize production acquired more popularity and acceptance.

Consumer acceptability of “tuwo” depends largely on its texture, i.e., firmness, stickiness and colour (Bello *et al.*, 1990). Since “tuwo” is eaten with hand dipped in a sauce, it is important that it should not be sticky to avoid clogging to the roof of the mouth and secondly that it should not be very soft to enable the ease of manipulation of “tuwo” between the thumb and the fore-fingers (Da *et al.*, 1982). Murty *et al.* (1982) observed that waxy grain samples produce fluid gels with low consistency, while grain flours from floury endosperm kernel have been reported to produce sticky “tuwo”. When “tuwo” is stored over-night as commonly done by “tuwo” consumers, the textural behaviour of the porridge tends to be influenced by the cultivar used for its preparation (Da *et al.*, 1982).

The nature of gene action influencing the texture of thick porridges of maize has received little attention. Kaw and Cruz (1991) studied the degree of heterosis for flour gel consistency in rice using  $F_1$  hybrids obtained from crossing 17 females with six male parents. The authors reported that a large number of the  $F_1$  hybrids expressed considerable heterosis when compared with the mid- and -high parent values and concluded that non-additive gene effects were most predominant.

This study was designed to investigate the nature of gene action on the rheological property of maize “tuwo”, a thick porridge flour meal.

## MATERIALS AND METHODS

Six open-pollinated maize cultivars, namely Pool 16 SR, TZESR-W, BC 63, DMR-ESR-W, Pop. 49 SR (EV 8349-SR) and Celaya waxy were crossed in all possible combinations excluding reciprocals. A set of the 15  $F_1$ 's without parents and reciprocals were grown in the University of Benin Teaching and Research Farm during the

1991 crop year. The entries were grown in a randomized complete block design with three replicates. The plot size was a single row plot of 5 m length. The rows were spaced 75 cm apart. Plants in each row were spaced 50 cm with two plants hill<sup>-1</sup>. Weeds were controlled by the application of 5.0 l ha<sup>-1</sup> of primextra (atrazine + metolachlor) as a pre-emergence herbicide, and fertilizers were applied at the rate of 120 kg ha<sup>-1</sup> N and 50 kg ha<sup>-1</sup> P<sub>2</sub>O<sub>5</sub> and K<sub>2</sub>O, respectively.

When the plants were fully matured, the ears were harvested on row basis. The kernels were shelled and dried at 48°C ± 2°C. A 100 g sample of whole grain was taken from each hybrid, ground with a kitchen blender with two vertically placed carborundum grinding stones. The ground flour was sifted through a sieve of 0.35 mm mesh (BSS 410).

**Preparation of “tuwo”.** “Tuwo” was prepared by the procedure described by Da *et al.*, (1982) with some modifications. Briefly, 10.0 g flour (dwb) was made into a slurry using 20 ml tap water. Ten ml of water was measured into a 150 ml beaker and heated over an electric burner until boiling. The flour slurry was poured into the boiling water and cooked for about 2–3 mins. The cooked “tuwo” was shared into two 10 ml beakers. “Tuwo” from the first beaker was evaluated for firmness after 1 hour as fresh “tuwo” and the second after 24 hr of storage at room temperature as stale “tuwo”. “Tuwo” quality was evaluated for firmness, stickiness and time of gelatinization. Firmness and stickiness were subjectively evaluated on a scale of 1–5. In the case of firmness 1 = very firm and 5 = very sticky. Both variables were measured by taking a small portion of the prepared “tuwo” and manipulating it into a ball with the forefingers and thumb. The degree of adherence of the “tuwo” to the fingers in the case of stickiness and the ease to form a ball in the case of firmness were scored. Time (minutes) of gelatinization was recorded as the time between the onset of cooking of the slurry to the onset of gelatinization.

Data were analysed as a randomized complete block design. Genotypic mean squares were partitioned into general combining ability (GCA) and specific combining ability (SCA) according to the procedure of Griffing (1956) using one set of  $F_1$  hybrids without parents and reciprocals.



unacceptable to consumers. A cross of line 3x5 produced "tuwo" with the least stickiness. DMR-ESR-W contributed the most favourable gene effect to non-sticky "tuwo" by contributing the least GCA effect (-0.83).

The array means for gel time revealed that the average effect of line 2 across the hybrids was the least and closely followed by line 6 (Table 5). The GCA effects ranged from -1.78 for line 2 to 1.89 for line 3. Line 3 contributed most unfavourable to delay in gel time while line 2 contributed most favourably to quicker gelatinization time. Line 2 could be considered useful for the production of gel products where instant cooking is desirable. Line 6, waxy maize with a GCA value of -0.36 for gelatinization time appears to agree with the finding of Murty *et al.* (1982) who reported that waxy sorghum gelled earlier than non-waxy.

Hybrids between lines (1x4) and (2x3) expressed the fastest gelatinization time (Table 6). Delayed gel time was observed in two hybrids of lines (1x3) and (3x4), respectively.

The preponderance of the variation due to GCA (Table 1) for firmness of stale "tuwo" indicates that additive genetic variance was more important than the non-additive gene effects. Similarly the significant mean squares due to GCA for firmness and stickiness of fresh "tuwo" suggest that the inheritance of both characters is controlled primarily by additive gene effects. This is, however, contrary to the findings of Kaw and Crux (1991) who reported that gel consistency in rice was controlled predominantly by additive genes. Improvement of "tuwo" texture (firmness and stickiness) can therefore be achieved using simple recurrent selection or recurrent full sib

TABLE 4. Hybrid means, array of means and general combining ability effects for stickiness.

Parents	Parents						Array Means	GCA effect
	1	2	3	4	5	6		
1. Pool 16 SR		3.0	3.7	1.7	3.0	4.0	3.1	-0.08
2. TZESR-W			3.0	3.3	2.7	4.3	3.3	0.17
3. BC 63				2.0	1.7	5.0	5.0	-0.08
4. DMR-ESR-W					2.0	3.3	3.5	-0.83
5. Pop. 49 SR						4.3	2.7	-0.50
6. Waxy							4.2	1.33
Mean							3.2	

TABLE 5. Hybrid means, array of means, and general combining ability effects for gelatinization time

Parents	Parents						Array Means	GCA effect
	1	2	3	4	5	6		
1. Pool 16 SR		35.0	40.3	31.0	39.0	34.0	35.9	-0.03
2. TZESR-W			32.3	38.3	33.7	33.3	34.5	-1.78
3. BC 63				41.0	37.3	36.3	37.4	1.89
4. DMR-ESR-W					32.7	35.7	35.7	-0.28
5. Pop. 49 SR						39.0	36.4	0.56
6. Waxy							35.7	-0.36
Mean							35.9	

TABLE 6. Specific combining ability effects for gelatinization time

Parents	Parents					
	1	2	3	4	5	6
1. Pool 16 SR		0.85	2.52	-4.65	2.85	-1.57
2. TZESR-W			-3.75	4.43	1.07	0.48
3. BC 63				3.48	-1.07	1.15
4. DMR-ESR-W					-3.37	0.35
6. Waxy						

family selection. The ration of GCA: SCA for gelatinization time was less than unity suggesting a preponderance of dominance gene effects. The highly significant mean square for SCA indicates that genetic progress from selection for gel time might be achieved using hybridization reciprocal recurrent selection.

#### REFERENCES

- Alika, J.E. 1991. Variation in starch yield and sensory characteristics of ogi among several maize (*Zeamays*, L.) hybrids. In: *Proceedings of National Conference on Maize Improvement, Production and Utilization in Nigeria*, IITA, Ibadan, 3–6 April. Fakorede, M.A.B. (Ed.). (In Press).
- Bello, A.B., Rooney, L.W. and Waniska. R.D. 1990. Factors affecting quality to a thick porridge. *Cereal Chemistry* 67: 20–25
- Da, S.J.O., Akingbala, L.W., Rooney, J.F., Scheuring, J.F. and Miller, F.R. 1982. Evaluation of quality in a sorghum breeding program. In: *Proceedings of the International Symposium on Sorghum Grain Quality*, 28–31 October, 1981, Patancheru, Andhra Pradesh, India. ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 406 pp.
- Griffing, B. 1956. Concept of general and specific combining ability in relation to diallel crossing systems. *Australian Journal of Biological Science* 9: 463–493.
- Kaw, R.N. and Cruz, N.M. 1991. Heterosis for grain quality in rice. *Indian Journal of Genetics* 51: 51–57.
- Murty, D. S, Patil, H.D. and House, L.R. 1982. Sankati quality evaluation of sorghum cultivars. In: *Proceedings of the International Symposium on sorghum Grain Quality*, 28–31, October 1981, Patancheru, Andhra Pradesh, India. ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 406 pp.
- Scheuring, J.F., Sidibe, S. and Kante, A. 1982. Sorghum alkali to: Quality considerations. *Proceedings of the International Symposium on Sorghum Grain Quality*, 28–31. October 1981, Pantacheru, Andhra Pradesh, India. ICRISAT (International Crops Research Institute for the Semi-Arid Tropics). 406 pp.

