

PERFORMANCE OF EDIBLE COATINGS FROM CARBOXYMETHYLCELLULOSE (CMC)  
AND CORN STARCH(CS) INCORPORATED WITH *Moringa oleifera* EXTRACT ON  
*CITRUS SINENSIS*

\*Adetunji C. O.<sup>1</sup>, Fawole O.B.<sup>2</sup>, Arowora K.A.<sup>1</sup>, Nwaubani S.I.<sup>1</sup>, Oloke J.K.<sup>4</sup>, Adepoju A.O.<sup>4</sup>  
Adetunji J.B.<sup>3</sup>, Ajani A.O.<sup>1</sup>

<sup>1</sup>Nigerian Stored Product Research Institute, Km 3 Asa dam road, P.M.B. 1489, Ilorin,

<sup>2</sup>University of Ilorin, Department of Agronomy, P.M.B.1515, Ilorin, Kwara State.

<sup>3</sup>University of Ilorin, Department of Biochemistry, P.M.B. 1515, Ilorin, Kwara State.

<sup>4</sup>Ladoke Akintola University of Technology, Department of Pure and Applied Biology P.M.B  
4000, Ogbomoso, Oyo State.

<sup>2</sup>University of Ilorin, Department of Agronomy, P.M.B.1515, Ilorin, Kwara State.

\*corresponding author: e-mail: charlignitar@yahoo.com, Telephone: +2348039120079

ABSTRACT

*Two different coatings were developed from Corn Starch (CSME) and Carboxymethylcellulose (CMCME) and their effects were investigated on the quality and storability of orange fruits. The two experimental coatings were: CSME and CMCME both mixed with 75mg/ml of crude extract of Moringa Oleifera. The following parameters were measured: Weight loss, ascorbic acid content and firmness. Four hundred and eighty (480) orange fruits were stored for seven weeks at ambient temperature. The overall result showed that polysaccharides coating from (CSME) and (CMCME) on orange fruits when compared to untreated in the following order: (CSME) > (CMCME) > Control in extending its shelf life.*

Keywords: Edible Coatings, *Moringa Oleifera*, Orange, Corn Starch and Carboxymethylcellulose.

INTRODUCTION

*Moringa oleifera* is considered one of the world's most useful trees, as almost every part of the tree can be used for food or has some other beneficial property. Moringa is a special food for the tropics, because the tree is in full leaf at the end of scarce (Iwu, 1993). It is available all year round. Almost all the parts of *Moringa Oleifera* are used as food and forage for livestock (Ram, 1994). The part (leaves, fruits, flowers and immature pods) are edible and form part of traditional diet in many countries of the tropics and subtropics (Odee, 1998).

The sweet orange (*Citrus sinensis* L. Osbeck) is the most commonly grown tree fruit in the world (Morton, 1987). Citrus fruits are produced all around the world and world citrus production in selected major producing countries in 2005/2006 is 72.8 million metric tons. Citrus fruits are said to be the first crops in the international trade in terms of values (CIAC, 2002).

Edible films and coatings are an environmentally-friendly alternative method to extend the

postharvest life of fresh and minimally processed fruits and vegetables (Baldwin, 1994; Olivas *et al.*, 2008; Pérez-Gago *et al.*, 2005; Vargas, Pastor, Chiralt *et al.*, 2008). They form a semi-permeable barrier to gas and water vapor that reduce respiration and weight loss. In addition, edible films and coatings may help in maintaining firmness and provide gloss to coated fruits.

According to their components, edible films and coatings can be divided into three categories: hydrocolloids (proteins and polysaccharides), lipids, and composites. Antioxidants, flavors and pigments, vitamins, and antimicrobial agents can be successfully incorporated into edible coatings to improve their functional properties. In the literature, several reviews reported on the efficacy of films and coatings containing antimicrobials to control microbial growth on fruits and vegetables (Ayala-Zavala *et al.*, 2008; Cagri, Ustunol, & Ryser, 2004).

Carboxymethyl cellulose (CMC) is a linear, long-chain, water-soluble, anionic polysaccharide.

Purified CMC is a white to cream-coloured, tasteless, odourless, free-flowing powder. Keller, J.D. (1986) and Hattori, K., *et al.* (2004).

From the literature reviews, corn starch (CS) appears to be an interesting alternative for edible films due to its abundance, cheap price, being biodegradable as well as edible (Biliaderis, C.G *et al.* (1999), Wing, R. E., *et al.* (1988), Bertuzzi, M.A., *et al.* (2007).

The present study was aimed at investigating the suitability of *Moringa oleifera* extract incorporated into CS and CMC as an edible coating to extend the shelf-life of orange fruits during ambient storage.

## MATERIALS AND METHODS

**Source of Materials:** Freshly harvested oranges were procured from Ipata market in Ilorin, Kwara state, Nigeria. They were selected on the basis of size, color and absence of external injuries. Fresh leaves of *Moringa oleifera* were obtained from Nigeria Stored Products Research Institute (NSPRI) garden at Ilorin, Kwara state Nigeria.

### Preparation of *Moringa oleifera* extract

*Moringa oleifera* leaves were dried in an open laboratory and ground into very fine powder using an electric blender (supermaster<sup>®</sup>, Model SMB 2977, Japan). The powder was further sieved to pass through 1 mm<sup>2</sup> perforations. The powder was then packed in plastic containers with tight lids and stored in a refrigerator at 4°C prior to use. Fifty grams of the dried *Moringa Oleifera* leaves was weighed and introduced into a conical flask containing 250ml of distilled water. The conical flask was then covered with aluminum foil and placed on a mechanical shaker. The suspension was then shaken for 48 hours at 190 rev. per min. The extract was decanted and passed through a clean muslin cloth and later filtered with Whatman filter paper. The filtrate obtained was evaporated to dryness at 50°C and the residues obtained are kept in an aluminum foil.

### Reconstitution and sterilization of extract

The dried residue was weighed into McCartney bottles and appropriate volume of distilled water was added to make a stock solution of 75mg/ml, for example 750mg in 10mls of distilled water. The

stock solution was then sterilized using 0.65 membrane filter by suction pump. The sterilized extract was stored inside McCartney bottle and kept in a refrigerator.

**Surface preparation of the Oranges:** The primary purpose of surface preparation was to remove all contaminants that would hinder proper coating adhesion and to render a sound clean substrate suitable for firm bonding. The surface should be in paint ready condition. Mold, mildew and/or algae were removed and sterilized with a 25% hypochlorite solution (1 gallon household bleach to 3 gallons water). The oranges were soaked in the 25% hypochlorite solution for two minutes.

**Preparation of edible coatings:**

1. Carboxy Methyl Cellulose (CMC) coatings (3%) was prepared by dissolving 6.0 g of methyl cellulose powder (Hangzhou Hongbo Chemical Co. Ltd, China) in 200 ml of water ethyl alcohol mixture (3:1L/L) at 80°C and stirred for 10 min by using magnetic stirrer. Ethyl alcohol was used to reduce drying time and obtain a transparent and shinny coating. 2% propylene glycol was also added in the formulation as plasticizer.

2. Corn Starch films were prepared as follows. Aqueous suspension of 3g CS/100 g with plasticizers was preheated at 100°C for 30 min in boiling water-bath, gelatinized at 160±5 °C in an oil bath for 30 min, cooled to 80°C, cast on the round polypropylene (PP) plate (300mm ID, 305mm OD and 10mm depth), and then dried at 50°C for 24 h in an oven. Glycerol was added to 100g of corn starch as plasticizers. In order to cook the Corn starch at 160±5°C, special stainless-steel high-pressure container (10mm thick) was then designed.

**Treatments**

T<sub>0</sub> (control):- T<sub>0</sub> was selected as the control (untreated oranges)

T<sub>1</sub> Oranges was coated with Corn Starch containing 70% *Moringa oleifera* (CSME)

T<sub>2</sub> Oranges was coated with Carboxymethylcellulose containing 70% *Moringa oleifera* (CMCME)

The treated and untreated oranges were packed in small plastic basket and each basket contains 20 orange fruits. The basket was stored at ambient temperature (27±2°C) and at 50-60% relative humidity. Physiochemical analysis was then carried out from 1-7weeks of coating.

**Firmness:** - Firmness was measured as the maximum penetration force (N) reached during tissue breakage, and determined with a 5 mm diameter flat probe. The penetration depth was 5 mm and the cross-head speed was 5 mm s<sup>-1</sup> using a TA-XT2 Texture Analyzer (Stable Micro Systems, Godalming, UK), MA. Oranges were sliced into halves and each half was measured in the central zone.

**Ascorbic acid:** - Ascorbic acid content was measured using 2, 5-6 dichlorophenol indophenols' method described by A.O.A.C 1994.

Percentage weight loss: - The weight loss of the orange fruit was determined using the Equation (1).  
 Weight loss (%)  $100 \times \frac{(M_1 - M_2)}{M_1}$

Where:  $M_1$  = Mass of sample before drying in g.  $M_2$  = Mass of sample after drying, in g.

Statistics

All results are means  $\pm$  S.E., SPSS software (version 12.0, SPSS Inc., US) was used for all statistical analysis for Analysis of variance. The significance level used was 0.05.

RESULTS AND DISCUSSION




Fig 1. Effect of edible coatings from CS and CMC with Moringa extract on firmness of orange at ambient temp

The mean $\pm$ SE value for the Firmness on coatings from CSME and CMCME were 3978.71 $\pm$ 451.14 lb/kg/N and 3635.71 $\pm$  563.17 lb/kg/N respectively while the mean $\pm$ SE value for the uncoated was 2904 $\pm$ 777.60 lb/kg/N.

During storage, the texture of the fruits is likely to soften due to several factors, including loss in cell turgidity pressure, loss of extracellular and vascular air and the degradation of the cell wall and consequent loss of water by the cell breakdown (Martínez-Ferrer, Harper, Pérez-Munóz, & Chaparro, 2002; Somogyi, Hui, & Barret, 1996).

Lerdthanangkul and Krochta (1996) also made similar observations and concluded that coatings and/or films significantly affected firmness of fruits in storage. The softening process in orange has been reported to be dependent on the increase in polygalacturonase,  $\beta$ galactosidase and pectinmethylesterase activities (Batisse *et al.*, 1996; Gerardi *et al.*, 2001; Remon *et al.*, 2003), being responsible for fruit quality loss. CSME and CMCME treatment significantly reduced the firmness losses (more than 50%) during ambient storage compared with control fruits. In addition, CSME and CMCME probably had some effects on the reduction of cell wall degrading-enzymes responsible for cucumber softening. These results show beneficial effects of the CSME and CMCME coatings on increasing the cucumber shelf life, since it has been postulated that fruit softening and texture changes during orange storage determine fruit storability and shelf life, as well as reduced incidence of decay and less susceptibility to mechanical damage (Batisse *et al.*, 1996; Vidrih *et al.*, 1998).

#### Ascorbic acid




Fig 2: Effect of edible coatings from CS and CMC with Moringa extract on Ascorbic acid content of orange at ambient temp

The mean $\pm$ SE value for ascorbic of the coatings from CSME and CMCME were 44.21 $\pm$ 0.49 mg/100g and 38.66 $\pm$  0.81 mg/100g respectively while the mean $\pm$ SE value for the uncoated was 30.32 $\pm$ 1.06 mg/100g.

Ascorbic acid content: Ascorbic acid is lost due to the activities of phenoloxidase and ascorbic acid oxidase enzymes during storage (Salunkhe *et al.*, 1991). Weichmann *et al.* (1985), while studying green bean, spinach and broccoli, postulated that the lower the oxygen content of the storage atmosphere, the smaller is the loss of ascorbic acid. The claim was that the oxidation of Vitamin C was mainly regulated by ascorbic acid oxidase and other oxidases, most of which had a

low affinity for oxygen. Ascorbic acid content decreased for cherries stored at both ambient temperature and cold temperature. CSME and CMCME coatings were effective in reducing the ascorbic acid loss under the storage conditions (Fig. 2). At the ambient temperature, the ascorbic acid contents of CSME and CMCME coated cucumber were significantly different from the control orange. The reduction of ascorbic acid loss in coated orange was due to the low oxygen permeability of CSME and CMCME coating which lowered the activity of the enzymes and prevented oxidation of ascorbic acid. The effect of low temperature significantly reduced the ascorbic acid loss. This shows the effect of temperature on the activities of the related enzymes.

Percentage weight loss




Fig 3: Effect of edible coatings from CS and CMC with Moringa extract on %Weight loss of orange at ambient temp

The coatings (CSME and CMCME) caused a significant ( $p=0.05$ ) decrease in weight loss compared with control sample as shown in fig 3 above. The mean $\pm$ SE value for the weight loss of CSME and CMCME were  $524.25\pm 4.52g$  and  $510.19\pm 5.60 g$  while the mean $\pm$ SE value for the weight loss of uncoated orange was  $351.81\pm 8.39g$ .

These results were in agreement with those of Mahmoud and Savello (1992) and Avena-Bustillos *et al.* (1997) who concluded that coatings and/or films significantly conserved water content. Post harvest weight changes in fruits and vegetables are usually due to the loss of water through transpiration. This loss of water can lead to wilting and shriveling which both reduce a commodity marketability. Edible films and coatings can also offer a possibility to extend the shelf life of fresh-cut produce by providing a semi-permeable barrier to gases and water vapor and therefore, they can reduce respiration, enzymatic browning and water loss (Gullbert, 1986; Baldwin & Nisperos-Carriedo Baker, 1995).

## CONCLUSION

Applications of Polysaccharide coatings to orange were shown to be beneficial in keeping the quality of the fruits in storage. Addition of *Moringa oleifera* extract into the Coatings slowed down the weight loss and reduced the ascorbic acid content. The shelf-life of orange was increased at an average temperature of  $27 \pm 2^\circ\text{C}$  and relative humidity 55-60%. Textural analysis showed that the coatings from CSME and CMCME could have a protective effect on orange, reflected by the greater firmness of coated samples during storage, which could reduce economic losses due to spoilage produced from mechanical damage during handling and transportation. Finally, the overall result showed that the coatings from CSME and CMCME is effective in extending the shelf-life of orange fruits when compared to untreated orange in the following order: CSME>CMCME>Control.

## REFERENCES

- AOAC, (1994). Official Methods of Analysis. *Association of Official Analytical Chemists*. 1111 North 19th Street, Suite 20, 16th Edi. Arlington, Virginia, USA. 22209.
- Avena-Bustillos R. J. Krochta J. M. and Saltveit M. E. (1997). Water vapor resistance of red delicious apples and celery sticks coated with edible caseinate-acetylated monoglyceride films. *Journal of Food Science*.62: 351-354.
- Ayala-Zavala, J. F., Del Toro-Sánchez, L., Alvarez-Parrilla, E., Soto-Valdez, H., Martín-Belloso, O. and Ruis-Cruz, S. (2008). Natural antimicrobial agents incorporated in active packaging to preserve the quality of fresh fruits and vegetables. *Steward Postharvest Review*, 3(9), 1-9.
- Baldwin, E. A. (1994). Edible coatings for fresh fruits and vegetables: past, present, and future. In J. M. Krochta, E. A. Baldwin, & M. O. Nisperos-Carriedo (Eds.), *Edible coatings and films to improve food quality* (pp. 25-64). Lancaster, PA, USA: Technomic Publishing Co. Inc.
- Baldwin, E. A., Nisperos-Carriedo, M. O. and Baker, R. A. (1995). Use of edible coatings to preserve quality of lightly (and slightly) processed products. *Critical Review Food Science Nutrition* 35, 509-524
- Batisse, C., Buret, M., Coulomb, P. J., (1996). Biochemical differences in cell wall of cherry fruit between soft and crisp fruit. *J. Agric. Food Chem.* 44, 453-457.
- Biliaderis, C. G., Lazaridou, A. and Arvanitoyannis, I. (1999). Glass transition and, physical properties of polyol-plasticised pullulan starch blends at low moisture. *Carbohydrate Polymers*, 40, 29-47.

- Bertuzzi, M. A., Castro Vidaurre, E. F., Armada, M., Gottifredi, J.C. (2007). Water vapor permeability of edible starch based films. *Journal of Food Engineering*, 80, 972-978.
- Cagri, A., Ustunol, Z., & Ryser, E. T. (2004). Antimicrobial edible films and coatings. *Journal of Food Protection*, 67(4), 833-848.
- CIAC, (2002). Strategic Investment Plan (Horticultural Australia). An *https documents available at www.ciac.org*.
- Gerardi, C., Blando, F., Santino, A., Zacheo, G., (2001). Purification and characterisation of a  $\alpha$ -glucosidase abundantly expressed in ripe sweet cherry (*Prunus avium* L.) fruit. *Plant Sci.* 160, 795805.
- Guilbert, S. (1986). Technology and application of edible protective films. In: Matholouthi, M. (Ed.), Food Packaging and Preservation: Theory and Practice. *Elsevier Applied Science Publishing Co.*, London, pp. 371-393.
- Hattori, K., Abe, E., Yoshida, T. and Cuculo, J. A. (2004). New solvents for cellulose II ethylenediamine/thiocyanate salt system. *Polymer Journal*, 36(2), 123-130.
- Iwu, M., 1993. Handbook of African Medical Plants. CRC press Florida, pp: 256-257.
- Keller, J.D. (1986). In: M. Gliksman, Editor, Sodium carboxymethylcellulose (CMC), Food Hydrocolloids, 3, *CRC Press, Boca Raton, Florida*, pp. 45-104.
- Pérez-Gago, M. B., Serra, M., Alonso, M., Mateos, M., & del Río, M. A. (2005). Effect of Whey protein- and hydroxypropyl methylcellulose-based edible composite coatings on color change of fresh-cut apples. *Postharvest Biology and Technology*, 36(1), 77-85.
- Ram, J., (1994). Moringa: A Highly Nutritious Vegetable Tree. Tropical Rural and Island. *A Toll Development, Experimental Station (TRIADES) Technical Bulletin No. 2*.
- Lerdthanangkul S. and Krochta J.M. (1996). Edible coating effects on post harvest quality of green bell peppers. *Journal of Food Science*. 61: 176-179.
- Morton, J., *Fruits of Warm Climates* (1987) Miami, FL, pp. 134-142. [Http://www.volusia.org/history/luegimgong.htm](http://www.volusia.org/history/luegimgong.htm)



- Martínez-Ferrer, M., Harper, C., Pérez-Munoz, F., & Chaparro, M. (2002). Modified atmosphere packaging of minimally processed mango and pineapple fruits. *Journal of Food Science*, 67(9), 3365-3371.
- Mahmoud R. and Savello P.A. (1992). Mechanical properties of and water vapor transferability through whey protein films. *Journal of Dairy Science*. 75: 942-946.
- Odee, D., 1998. Forest biotechnology research in dry lands of Kenya. The development of *Moringa* species. *Dryland Biodiversity*, 2: 7-12.
- Olivas, G. I., Dávila-Aviña, J. E., Salas-Salazar, N. A., & Molina, F. J. (2008). Use of edible coatings to preserve the quality of fruits and vegetables during storage. *Stewart Postharvest Review*, 3(6), 1-10.
- Remón, S., Venturini, M.E., López-Buesa, P., Oria, R., (2003). Burlat cherry quality after long range transport, I optimisation of packaging conditions. *Inno. Food Sci. Emerg. Technol.* 4, 425-434.
- Somogyi, L. P., Hui, Y. H., & Barret, D. M. (1996). Processing fruits: Science and technology (Vol 2). *Lancaster, Basel: Technomic Publishing Co.*
- Vargas, M., Pastor, C., Chiralt, A., McClements, D. J., & González-Martínez, C. (2008). Recent Advances in edible coatings for fresh and minimally processed fruits. *Critical Reviews in Food Science & Nutrition*, 48, 496-511.
- Vidrih, R., Zavrtnik, M., Hribar, J., (1998). Effect of low O<sub>2</sub>, high CO<sub>2</sub> or added acetaldehyde and ethanol on postharvest physiology of cherries. *Acta Hort.* 2, 693-695.
- Wing, R. E., Maiti, S. and Doane, W. M. (1988) Amylose content of starch controls the release of encapsulated bioactive agents, *Journal of Controlled Release*, 7 33-37.