

MRSA: the leading pathogen of orthopedic infection in a tertiary care hospital, South India

Thimmappa Latha¹, Bhat Anil², Hande Manjunatha², Mukhopadhyay Chiranjay²,
Devi Elsa¹, Nayak Baby¹, George Anice¹

1. Manipal College of Nursing Manipal, Manipal Academy of Higher Education Manipal.
2. Kasturba Medical College Manipal, Manipal Academy of Higher Education Manipal.

Abstract

Background: The rate of infection is high and heterogeneous in developing countries. This study aimed to find the rate and pattern of infection in a tertiary care hospital with a goal to improve the infection control practices.

Methods: The study was conducted in the orthopedic units of a multispecialty teaching hospital. Medical records of major orthopedic surgery adult patients without immunosuppression state were included. The bacterial culture report of the wound swabs were noted over a period of one year. The bacterial culture testing was performed by a recommended method.

Results: Among 2,249 orthopedic surgery patients, 83.7% were males, 49.1% had open wounds during admission and 32.2% patients were infected. Majority (64.2%) of the injuries were in the lower limb with 19.4% patients having undergone multiple surgeries during hospitalization. A total of 946 pathogens were grown from 725 specimens. *Staphylococcus aureus* was the maximum (48.4%) followed by *Pseudomonas aeruginosa* (26.3%) and *E coli* (16.7%). Among them, 57.3% were Methicillin Resistant *Staphylococcus aureus* (MRSA) and was the leading pathogen causing infection among orthopedic patients.

Conclusion: MRSA infection was high. Consequent to this, an interventional program entitled 'Extended Infection Control Measures' was designed to reduce the burden of infection.

Keywords: MRSA, orthopedic, infection, South India.

DOI: <https://dx.doi.org/10.4314/ahs.v19i1.12>

Cite as: Latha T, Anil B, Manjunatha H, Chiranjay M, Elsa D, Baby N, et al. MRSA: the leading pathogen of orthopedic infection in a tertiary care hospital, South India. *Afri Health Sci.* 2019;19(1). 1393-1401. <https://dx.doi.org/10.4314/ahs.v19i1.12>

Introduction

Health care associated infection is an important unwanted event in health care delivery system which accounts for 7/100 infection in the developed countries and 10/100 in the developing countries¹. Global Antibiotic Resistance Partnership (GARP) – India working group and Centre for Disease Dynamics, Economics and Policy (CDDEP) have reported that the hospital incidence of acquired infection in India is relatively high. The rate of Vancomycin

resistant Enterococcus and Methicillin Resistant *Staphylococcus aureus* (MRSA) infections have significantly increased in India². *Staphylococcus aureus*, coagulase negative *Staphylococcus*, *E coli*, *Klebsiella pneumoniae*, *Enterobacter species*, *Citrobacter species*, *Pseudomonas aeruginosa* are the frequently noted pathogens causing infection in the hospital³⁻⁵. These micro-organisms are commonly prevalent in the air, contaminated food and water, medication, equipment, soiled linen, hospital waste and contaminated wounds⁵. Patient's flora, cross-infection from other patients and health care personnel also contribute to the source of infection⁵. Open fractures, crush injuries, chronic osteomyelitis, chronic ulcers, instrumentation, arthrotomies, amputation and implants cause a variety of infection among orthopedic patients^{3,5,6}. However, the rate of orthopedic infection is not homogenous^{4,5}.

Corresponding author:

Bhat Anil,
Kasturba Medical College Manipal,
Manipal Academy of Higher Education Manipal.
Email: anilkbhat@yahoo.com

The infection increases the hospital stay, mortality and the overall cost of health care^{1,7,8}. Therefore, this study was aimed to find out the infection rate and pattern among orthopedic surgery patients in a tertiary care hospital with a view to develop an interventional program to reduce the incidence of such infections. Though all the hospitalized patients have a risk to acquire the infection, orthopedic patients have a greater risk due to prolonged hospitalization, nature of surgeries, open wounds and implants.

Methods

This study was conducted in an orthopedic unit of a tertiary care hospital, in South India. The hospital has more than 2032 beds with 150 beds for department of orthopedics. The data was collected from medical records of orthopedic surgery patients for one year in this retrospective observational study. After obtaining permission from institutional ethical committee, medical records of all the orthopedic patients, who underwent both elective and emergency orthopedic surgery were reviewed. Medical records of patients with compromised immunity status (HIV/AIDS, on immunosuppression therapy such as post-transplant patients) were excluded. Patients who required a bacterial culture test for a suspected or an obvious infection in the wound were included. The asymptomatic carrier state was not investigated. The wound swabs or tissue were collected during the surgery or in the post-operative period. The swabs were sent to microbiology department immediately for bacterial culture and

sensitivity testing. The culture and antibiotic sensitivity testing was done by standard recommended method⁹.

In one year, 4,382 orthopedic surgeries were performed. A total of 2,282 patients had undergone major orthopedic surgeries and 2,249 patients were eligible as per the inclusion criteria. The surgeries included in the study were major wound debridement, implant and prosthesis insertion, joint replacement surgeries, major graft / flap reconstructions, and spinal surgeries. The data was collected using the proforma which consisted of age, gender, presence of open wound at the time of admission, location of injury, type of surgery and date of the surgery, pathogen grown in the culture test and date of the culture tested were included in the proforma. Two investigators independently collected the data. The reliability of the data collection was assessed by inter-rater reliability and found reliable (97%).

Results

A total of 725 orthopedic surgery patients were found to be infected during the study period. The mean age of the patients was 40.3 ± 17.3 years and 83.7% were males. Nearly half (49.1%) of the patients had open wounds during admission. The patients had injuries at different areas and the details are given in figure 1. All patients (100%) had surgery done once, 31.9% twice and 9% had three surgeries done (table 1). The most of common surgical procedure performed was wound debridement (table 1).



Fig 1: Location of injury among orthopedic surgery patients (total patients 725)

Table 1: The type of surgery among orthopedic surgery patients n=725

Name of the surgery	First time surgery		Second time surgery		Third time surgery	
	f	%	f	%	f	%
Wound debridement	235	32.4	64	8.8	11	1.5
Flap reconstruction/ division	13	1.8	8	1.1	6	0.8
External Fixator application	24	3.3	5	0.7	-	-
Implant removal	107	14.8	15	2.1	3	0.4
Split skin graft	7	1.0	24	3.3	11	1.5
Implant insertion	80	11.0	38	5.2	7	1.0
Replacement (Knee, Hip)	35	4.8	-	-	-	-
Bone graft	16	2.2	5	0.7	-	-
Incision and Drainage	14	1.9	2	0.3	-	-
Spine Surgeries	17	2.3	-	-	-	-
Exploratory Biopsy	9	1.2	-	-	-	-
Amputation	12	1.7	3	0.4	1	0.1
Osteotomy, arthroscopic debridement, corticotomy	15	2.1	-	-	-	-
Multiple surgeries	141	19.4	67	9.2	26	3.6
Total	725	100	231	31.9	65	9.0

f= frequency

Bacterial culture testing report of 725 (32.2%) specimens showed a growth of 946 micro-organisms among 2249 eligible orthopedic surgery patients. Some patients re-

vealed the presence of monomicrobial and some patients had polymicrobial growth in their bacterial culture testing, making a total of 946 isolates (table 2).

Table 2: The rate of Gram positive and Gram negative microbes among orthopedic surgery patients n= 725 patients

	Frequency	Percentage
Gram positive Micro-organism (Total 382 Organisms)		
<i>Staphylococcus Aureus</i>	351	48.4
• MRSA	201	27.7
• MSSA	150	20.7
<i>Enterococcus species</i>	31	4.3
Gram negative Micro-organism (Total 564 organisms)		
<i>E coli</i>	121	16.7
<i>Proteus vulgaris</i>	18	2.5
<i>Acinetobacter</i>	63	8.7
<i>Pseudomonasaeruginosa</i>	191	26.3
<i>Klebsiella pneumonia</i>	106	14.6
<i>Enterobacter species</i>	29	4
<i>Proteus mirabilis</i>	11	1.5
<i>Klebsiella oxytoca</i>	10	1.4
<i>Citrobacter species</i>	15	2.1

Note: Polymicrobial growth was seen in some sample and a total growth of organisms was 946 among 725 patients.

Staphylococcus aureus was seen among 351 isolates, amounting to 48.4% compared to the other organisms. Among the 351 *Staphylococcus aureus*, 201 (57.3%) were MRSA and 150 (42.7%) were Methicillin sensitive *Staphylococcus aureus* (MSSA).

The calculated rate of infection among orthopedic surgery patient shows that MRSA was the leading cause

of infection (27.7%), followed by *Pseudomonas aeruginosa* (26.3%) and MSSA (20.7%). The infection rate during the study was 32.2% (725/2249).

The pattern of infection observed over one year period shows no uniformity. The monthly distribution of infection was suggestive that it was higher during June, July and August and significantly less in December and January (table 3).

Table 3: Monthly distribution of microbes among orthopedic surgery patients n=946

Month (2012-2013)	<i>E coli</i>	<i>Proteus vulgaris</i>	<i>Acinetobacter species</i>	<i>Pseudomonas aeruginosa</i>	MSSA	MRSA	<i>Klebsiella pneumonia</i>	<i>Enterobacter species</i>	<i>Citrobacter species</i>	<i>Enterococcus species</i>	<i>Proteus mirabilis</i>	<i>Klebsiella oxytoca</i>	Total
June	25	1	3	21	24	21	9	9	1	-	-	-	114
July	14	3	4	14	12	34	10	-	3	1	-	-	95
August	16	-	2	29	19	18	13	6	2	1	3	1	110
September	5	4	6	15	12	9	9	2	-	-	-	-	62
October	8	3	3	17	10	8	9	3	1	-	3	2	67
November	6	3	5	16	16	9	8	1	2	1	1	1	69
December	6	3	3	7	9	21	1	4	1	-	-	1	56
January	5	-	5	13	8	18	5	-	-	4	-	1	59
February	6	1	6	15	10	17	8	1	2	8	-	-	74
March	9	-	6	17	12	15	13	-	1	7	2	2	84
April	9	-	9	18	11	14	11	2	1	7	1	-	83
May	12	-	11	9	7	17	10	1	1	2	1	2	73
Total	121	18	63	191	150	201	106	29	15	31	11	10	946

Discussion

The infection among orthopedic patients is very common. In the present study, the infection was higher among males (83.7%). A similar result, 72.3%¹¹, 75.5 %¹², 81.4 %¹³ of infection among males is seen in previous reports. The mean age of the present study population was 40.3+17.3 years which is in par with the earlier study showing the maximum infection in the age group of 31-40 years¹². Motor vehicles and road traffic accidents, fall from height and certain risk taking behaviors are more common among men could be the cause of the high prevalence of infection among them.

Infection of the leg and foot was 81.6%¹², 64%⁴ and the upper limbs 12.1%¹². In consensus with the prior studies, this study found that the location of injury was more in lower limbs (right lower – 37% and left lower – 27.2%). Possibly lower limbs, especially right side, is more prone to get injured as most of the people are right side dominant. Also the Indian vehicles have the driving seat on the right hand side which also could explain the reason for the predominance of the injury on the right side.

Wound infection results in increased morbidity and healthcare cost^{1,7,8}. Developing infection in the hospital is one of the adverse events. *Staphylococcus aureus* infection, 48.4% (351/725) is the major threat for orthopedic patients. The results are concurrent with the studies worldwide (table 4). The rate of orthopedic surgical site infection however, differed from hospital to hospital ranging from 12%¹⁵ to 40%⁵. Nevertheless, the much higher infection rate of 68% of infection is published among orthopedic patients in general and *Pseudomonas aeruginosa* 18 (26.4%) was the predominant isolate followed by *Staphylococcus aureus* 17 (25%)⁶.

The presence of MRSA (27.7%) among orthopedic surgery patient is alarmingly high in the present study. This study also noted that among the *Staphylococcus aureus* infections, more than half of (57%) was methicillin resistant. Table 4 summarizes the rate of infection from different studies. Added to that, Jain and Banerjee has observed 39.6% (21/53) *Staphylococcus aureus* as the most prevalent infection at surgical site among orthopedic surgery patients¹⁷.

Table 4: Rate of orthopedic infections in the previous studies

	Sample	<i>Staphylococcus aureus</i>	<i>Enterococcus species</i>	<i>E coli</i>	<i>Proteus vulgaris</i>	<i>Acinetobacter</i>	<i>Pseudomonas aeruginosa</i>	<i>Klebsiella pneumoniae</i>	<i>Enterobacter species</i>	<i>Klebsiella oxytoca</i>	<i>Citrobacter species</i>
Present study	Wound/ tissue of all major orthopedic surgery patients with ≥18 years (725 patients)	37	3.3	12.7	1.9	6.6	20.1	11.1	3.1	1.1	1.6
Agrawal et.al, 2008 ⁵	Fracture, bed sore, wound of all orthopedic patients (Jan 2003-March 2004)	18.9	NA	NA	6.31	NA	26.13	8.11	NA	8.11	NA
Dhawan et al., 2005 ⁴	Wound swab of ortho patients – hospitalized (Jan 2000-June 2003)	40	4	11	1	12	12	8	2	NA	NA
Al-Mulhim et. al., 2014 ¹⁹	Wounds swabs of hospitalized orthopedic surgery patients	29.1	17.7	3.8	NA	21.5	19	3.8	NA	NA	NA
Dash et.al., 2016 ²⁰	Wound swab from compound fractures and post-operative orthopedic patients	26.9	4	1.8	3.4	8.8	11.4	10.8	NA	NA	5
Chandrasekar, 2016 ¹²	Wound swabs of hospitalized orthopedic patients	48.7	NA	22	3.3	3.3	25.4	15.3	22	NA	8.4
Agaja, 2008 ¹⁵	Patients with osteomyelitis underwent surgery	37.6	NA	9.3	6.5	NA	6.5	9.3	NA	NA	NA
Muley, L., 2015 ²¹	Hospitalized orthopedic patients	40.9	12	15.9	2.3	NA	13.6	7.9	NA	2.3	NA
Das, 2015 ¹⁶	Orthopedic surgical site infection	24.8	NA	18.9	5.4	2.7	18.9	8.1	8.1	NA	NA
Vasundhara Devi, 2017 ¹⁷	Wound swab of orthopedic patients	25	NA	7.3	3	1.5	26.4	6	1.5	NA	7.3

NA: Not Available

A report conveyed that 53% Gram positive and 47% Gram negative infections were observed among orthopedic in-patients⁴, whereas another study says 51.3% infection was Gram negative¹². In addition to this, a study has estimated 56.1% of Gram positive and 43.9% Gram negative bacilli infection¹¹. The present study shows 40.6% Gram positive and 59.4% Gram negative microbes.

The present study has seen 57.3% of *Staphylococcus aureus* as MRSA. The rate of MRSA is dissimilar between the hospitals ranging from 12%²⁰, 23.5%¹⁸, 50%¹², to 63%³. A study has reported 800 (38%) positive cultures and *Staphylococcus aureus* (32%) was the most common pathogen causing surgical site infection. MRSA was isolated in 44% of *Staphylococcus aureus* infections. But, the rate of MRSA was not similar in different surgical specialty such as neurosurgery (26%), orthopedics (24%), pediatric (17.8%), cardiothoracic (14%), and general surgery (7.1%)²¹. In a four year study on the prevalence of MRSA infec-

tion among both out-patient and in-patient department, 38.44% MRSA infection was documented²².

One of the causes for higher rate of infection is performing emergency surgeries¹⁸. The present study has noticed that many patients were admitted from the trauma center and have undergone emergency surgeries. Usually, emergency surgeries are not well prepared like elective surgeries. Patients with an open fracture tend to be infected more (14.7%) than a closed fracture (4.2%)⁹. Having an open wound at the time of admission increases the risk of infection. Patients with open fracture and fracture fixation have a high rate of infection²³. Possibly open wounds (49.1%) were one of the causes of high infection rate among our patients. Also, repeated surgeries are common among orthopedic patients. many a times, wound debridement is performed as an emergency procedure for severe traumatic wounds. Later, the implant is used for internal fixation of fracture. Covering the wound with

skin graft or flap is also a frequently performed surgery. Therefore, repeated exposure of orthopedic patients to operation room is unavoidable.

The growth of polymicrobial in an orthopedic wound is not uncommon¹⁸. Open fractures have a high risk of polymicrobial growth (5.8%) due to exposure of the wounds and contamination than that of closed fracture (1%)¹⁰ and many of our patients had polymicrobial infections; i.e. 946 microbes were grown from 725 patients.

We have observed that the infection rate was higher in the months of June, July and August and relatively lesser during December and January. In the months of June, July and August it is rainy season and December and January is winter in the present study setting. However, we could not trace any literature in relation to seasonal variation of bacterial infection among orthopedic surgery patients.

The wound swab was sent either during or after the surgery and the data was collected from the records. The investigators, therefore, were unable to differentiate community acquired infection and hospital acquired infection. Initial screening of pre-operative patients is not a routine practice in this study setting. We lacked data on antibiotic/s usage among the study population. Hence, additional research is needed to determine the rate of community acquired and hospital acquired infection and the causes of higher infection rate among the orthopedic cohort.

GARP – India working group has reported that the unreasonable use of antibiotic is the major threat for increasing multidrug resistant organism. In India, the abuse of antibiotics is not a new concept. Irrational or inappropriate prescription such as incorrect dose, frequency or duration; redundant or not considering the adverse interaction or reaction with other medications are observed in many parts of India. Prescribing the excess or unnecessary antibiotics has been documented in public and private hospitals, clinics and pharmacies²⁴. The studies also have explored the antibiotic usage for simple viral problems such as upper respiratory infection, diarrhea, etc.^{25,26}.

Certainly, these factors will increase the burden of the antibiotic resistant pathogen. The antibiotics are sold in many pharmacies without a proper prescription. In some

places, it is available in general stores as well. The current study has been carried out in a tertiary care referral hospital. Hence, the usage of antibiotics by the primary physician or in the local hospital cannot be denied. Often, many orthopedic patients visit or get admitted to the hospital and undergo invasive procedures such as implant insertion or implant removal. Thus, the risk of MRSA increases among orthopedic surgery patients.

Coia et al. have given a systematically prepared guideline for prevention of MRSA infection²⁷. The guideline explains about the different measures to be instigated in a hospital setting and feasibility of adopting this guideline in our setting need to be explored. However, executing a self-prepared guideline would give a better control of infection compared to adopting the guidelines from developed countries, as there are differences in hospital setting, resource availability and utilization, acceptance by the employees, practicability in implementing the measures and the causes of infection.

This study was conducted to develop an interventional program to reduce the infection rate. There are multiple interventions that have tried to decrease MRSA infection such as hand hygiene, nasal decolonization, pre-operative screening, isolation, cohort/barrier nursing, floor and environmental cleaning, etc. Unfortunately, no single intervention is found to be effective in reducing MRSA infection rate. Therefore, we have planned an intervention to find out the effectiveness of Extended Infection Control Measures (EICM) to reduce MRSA infection among orthopedic surgery patients. EICM includes routine infection control policies of the hospital along with other interventions such as pre-operative screening, treating high risk patients, isolation, education and frequent reinforcement of infection control ensures to health care providers, screening of healthcare providers to identify and treat carrier status.

Conclusion

Infection among the orthopedic surgery patients can have devastating consequences and hence newer measures are needed to combat this. Many hospitals in India have infection control committees which are working very hard to bring down the infection rates. Yet, the infection rate in several hospitals is high. The current study shows a

high rate of MRSA infection. Often, traumatic injury patients are encountered with contaminated wounds and if not treated immediately, may end up with wound infection. Further, these patients may receive an improper or no first aid after the traumatic injury. Moreover, they may require emergency surgery and very little time is available for patient preparation. Some of the orthopedic patients undergo repeated surgeries such as debridement, skin or bone grafting, instrumentation and removal, open reduction and internal fixations etc. Hence, frequent hospitalization or visit to an out-patient department is directly proportional to the rate of infection. Though we cannot avoid some of these events, we certainly can reduce the infection rates in the present setting.

Acknowledgement

We acknowledge the staffs of Medical Record Department (MRD) for providing us the necessary files for smooth data collection process.

References

1. World Health Organisation. Health care-associated infections Fact sheet. World Heal Organisation [cited 2016 November 22]. 2015;4. Available from: http://www.who.int/gpsc/country_work/gpsc_ccisc_fact_sheet_en.pdf
2. Hospital-acquired infections high in India. 22 September 2011. [cited 2017 January 21]. Available from: <http://www.indiamedicaltimes.com/2011/09/22/hospital-acquired-infections-high-in-india-study/India Medical Times>.
3. Patted SM, Chinagudi S, Soragavi VR, Bhavi SB. The prevalence of MRSA infection in orthopaedic surgery in a Medical College Hospital : A 2-year analysis. *Biomed Res-India*. 2013;24(1):33–5.
4. Dhawan B, Mohanty S, Das BK, Kapil A. Bacteriology of orthopaedic wound infections in an Indian Tertiary Care Hospital. *Indian J Med Res*. 2005 April;121:784–5.
5. Agrawal AC, Jain S, Jain RK, Raza HKT. Pathogenic bacteria in an orthopaedic hospital in India. *J Infect Developing Countries*. 2008;2(2):120–3.
6. Mathur P, et al. Implementation of a short course of prophylactic antibiotic treatment for prevention of post-operative infections in clean orthopaedic surgeries. *Indian J Med Res*. 2013 January;137(1):111–6.
7. Zimlichman E, Henderson D, Tamir O, Franz C. Health Care–Associated Infections A Meta-analysis of

Costs and Financial Impact on the US Health Care System. *JAMA Intern Med*. 2013;173(22):2039–46.

8. Yalcin AN. Socioeconomic burden nosocomial of nosocomial infections. *Ind J Med Sci*. 2003;57(10):6–9.
9. O’Leary WM. Practical Handbook of Microbiology [Internet]. Practical Handbook of Microbiology 1989;109–11. [cited 2017 January 15] Available from: <http://books.google.com/books?hl=de&lr=&id=erL29sLL5nEC&pgis=1>
10. Oliveira PR, Carvalho VC, da Silva Felix C, de Paula AP, Santos-Silva J, Lima ALLM. Incidence and microbiological profile of surgical site infections following internal fixation of closed and open fractures. *Rev Bras Ortop*. 2016;51(4): 396–9.
11. Naik TB, Wadekar MD, Upadhya A, Mane V. Cross sectional analysis of aerobic bacteria and their antibiotic susceptibility pattern among orthopedic wound infections at a tertiary care hospital in Karnataka. *Indian J Microbiol Res*. 2016;3(1):58. Available from: <http://www.indian-journals.com/ijor.aspx?target=ijor:ijmr1&volume=3&issue=1&article=015>
12. Chandrashekhara B, Sangeetha S. Isolation Identification and its Drug Susceptibility Pattern in Orthopaedic Wound Infections. *Int J Curr Microbiol App Sci*. 2016;5(9):213–8.
13. Akinkunmi EO, Adesunkanmi AR, Lamikanra A. Pattern of pathogens from surgical wound infections in a Nigerian hospital and their antimicrobial susceptibility profiles. *Afr Health Sci*. 2014;14(4):802–9.
14. Agaja SB, Ayorinde RO. Chronic osteomyelitis in Ilorin, Nigeria. *S Afr J Surg*. 2008;46(4):116–8.
15. Das R, Singh A, Srivastava P, Pradhan S, Murthy R. Microbial Profile and Antibiotic Susceptibility Pattern of Surgical Site Infections in Orthopedic Patients at a Tertiary Hospital in Bilaspur. *International Journal of Scientific Study*. 2015 June;3(3):43–7.
16. Devi PV, Reddy S, Shabnum M. Microbial profile and antibiotic susceptibility pattern of orthopedic infections in a tertiary care hospital: A study from South India. *Int J Med Sci Public Heal*. 2017;6(4):1–4.
17. Jain B, Banerjee M. Surgical site infections and its risk factors in orthopaedics: a prospective study in teaching hospital of central India. *Injury*. 2013;2(1):110–3.
18. Al-Mulhim FA, Baragbah MA, Sadat-Ali M, Alomran AS, Azam MQ. Prevalence of surgical site infection in orthopedic surgery: A 5-year analysis. *Int Surg*. 2014;99(3):264–8.

19. Dash A, et al. Surveillance of antibiotic sensitivity and resistance pattern of bacteria isolated from orthopaedic wound discharge. *Int J Pharm Sci Rev Res.* 2016;36(1):208–11.
20. Muley VA, Ghadage DP, Bhore A V. Bacteriological Profile of Neonatal Septicemia in a Tertiary Care Hospital from Western India. *J Glob Infect Dis.* 2015;7(6):75–7.
21. Tyagi A, Kapil A, Singh P. Incidence of Methicillin resistant *Staphylococcus aureus* (MRSA) in pus samples at a tertiary care hospital, AIIMS, New Delhi. *Journal, Indian Acad Clin Med.* 2008;9(1):33–5.
22. Tiwari HK, Sapkota D, Sen MR. High prevalence of multidrug-resistant MRSA in a tertiary care hospital of northern India. *Infect Drug Resist.* 2008;1:57–61. PubMed.
23. Willey M, Karam M. Impact of Infection on Fracture Fixation. *Orthop Clin North Am. Orthop Clin N Am.* 47 (2016) 357–4.
24. Global Antibiotic Resistance Partnership-India (GARP-India). *Situation Analysis: antibiotic Use and Resistance in India.* 2011;(March):1–83.
25. Sharma M, Eriksson B, Marrone G, Dhaneria S, Stålsby Lundborg C. Antibiotic prescribing in two private sector hospitals; one teaching and one non-teaching: A cross-sectional study in Ujjain, India. *BMC Infect Dis.* 2012;12(155):1–9. PubMed.
26. Kotwani A, Holloway K. Antibiotic prescribing practice for acute, uncomplicated respiratory tract infections in primary care settings in New Delhi, India. *Trop Med Int Health.* 2014;19(7):761–8.
27. Coia JE, et al. Guidelines for the control and prevention of methicillin-resistant *Staphylococcus aureus* (MRSA) in healthcare facilities. *J Hosp Infect.* 2006;63(sup 1):1–44.