

Cyclosporiasis: an emerging public health concern around the world and in Africa

Robert M. Karanja¹, Wangeci Gatei^{2*} and Njeri Wamae²

¹Centre for Biotechnology Research & Development (CBRD), Kenya Medical Research Institute, P.O. Box 54840 City Square, Nairobi, Kenya;

²Centre for Microbiology Research (CMR), Kenya Medical Research Institute, P.O. Box 19464-00200, Nairobi, Kenya;

Background: Cyclosporiasis is an emerging gastro-enteric disease caused by the coccidia protozoan *Cyclospora cayetanensis*. It is associated with diarrhoea among children in developing countries, in the Americas where *C. cayetanensis* is endemic, traveller's diarrhoea and/or food and waterborne outbreaks in the developed countries.

Objectives: The aim of this review is to highlight cyclosporiasis and its relevance to public health in East Africa and Africa at large.

Methods: All literature on *Cyclospora*, *C. cayetanensis*, cyclosporiasis in Africa, and endemic cyclosporiasis was searched from libraries, colleagues and internet but only literature on its history, clinical presentation, epidemiology in endemic settings, and occurrence in Africa were scrutinised.

Results: In Sub Saharan Africa, cyclosporiasis has been reported in at least 3 countries, including Tanzania, in East Africa, occurring in both immunocompromised and immunocompetent patients. Zoonotic species of *Cyclospora* have also been identified in East African primates, indicating likely endemicity of this little reported disease in the region. This can be attributed to lack of awareness in the public and medical profession concerning the disease, and therefore not routinely checked at the health centres. Cyclosporiasis is characterized by intermittent diarrhoea, and secondary conditions or sequelae such as reactive arthritis syndrome (Reiter's syndrome), have been associated with progression of the disease. Its management is based on antibiotics, an unusual scenario for a protozoa.

Conclusions: Although many aspects of this disease and its transmission remain an enigma, the situation has been rapidly changing since the disease first came to medical attention in the 1970s.

African Health Sciences 2007; 7(2) 62-67

Introduction

Cyclosporiasis and Africa – should we be concerned?

Cyclosporiasis is a diarrhoeal disease caused by the protozoan pathogen *Cyclospora cayetanensis*¹. It is considered an emerging disease of public health concern primarily in the developed countries where it has been identified as the cause of several outbreaks in North America and Europe, and with traveller's diarrhoea^{2,6}. In these countries, cyclosporiasis transmission has primarily been linked to foods imported from developing countries adding impetus to the review of food import regulations with far reaching implications on developing country economies that depend on North-South trade in agricultural produce^{4,9}. The transmission of the disease in the developing South where *C. cayetanensis* is endemic, remains little understood but has been associated with water and sanitation in Nepal, Guatemala, Nigeria and Egypt^{7,10,11}.

Ashford is credited with having first identified *C. cayetanensis*¹² as a human pathogen in 1979 in Papua New Guinea. Thereafter, the parasite remained largely uninvestigated and it was not until 1994, when the first detailed morphological description and naming of *Cyclospora cayetanensis* was done¹. Few studies have been carried out since then to determine the transmission and epidemiology of cyclosporiasis in Africa and its public health impact on the continent. The disease has been reported in Nigeria, Tanzania and Egypt¹³⁻¹⁵. Although it is thought to be self-limiting in immuno-competent individuals, cyclosporiasis can cause prolonged diarrhoea that could be life threatening in immuno-compromised patients. Moreover, like other coccidian parasites, *C. cayetanensis* has been associated with various sequelae including biliary disease, acalculous cholecystitis, Guillain-Barré syndrome and reactive arthritis syndrome following prolonged infection¹⁶⁻¹⁹.

The study of cyclosporiasis is rife with difficulties that have largely contributed to its underreporting in Africa and little public awareness²⁰. A key contributor to this is the specialised staining methods for diagnosis that are not routinely carried out in clinical

Correspondence

Robert M. Karanja

Centre for Biotechnology Research & Development (CBRD),

Kenya Medical Research Institute,

P.O. Box 54840, Nairobi, Kenya

Email: RKaranja@kemri.org

laboratories²⁰⁻²². Additionally, efforts to elucidate its transmission and biology in endemic areas is impeded by lack of sensitive and specific methods for *Cyclospora* detection in the environment, agricultural produce and water bodies and further compounded by the lack of a suitable animal-model²³⁻²⁶. Cyclosporiasis has therefore remained a little known disease albeit with potentially far reaching impact on the health and economy of developing countries due to its impact on the horticulture industry as a food borne pathogen. Several publications pertaining to the methods of diagnosis and detection exist and have been ably reviewed by others⁹. Our aim is to bring this disease to the attention of African health researchers and to highlight potential areas of study in our likely endemic setting.

History of Cyclosporiasis

According to Lainson²⁷, it was Eimer who as early as 1870 first described an organism in intestines of the mole *Talpa europaea* that was eventually characterized and named *Cyclospora caryolytica* by Schaudinn in 1902²⁸. But *C. glomericola*, observed in the millipede, was the first species to be classified as *Cyclospora* by Schneider who established the genus in 1881²⁹. So far *C. glomericola* is the only species out of a total of 19 *Cyclospora* species identified to date to be encountered in an invertebrate host, the rest being found in reptiles (snakes) and mammals (rodents and non-human primates)²⁷.

It was not until the 1970s that cyclosporiasis first came to medical attention when an un-described coccidian was associated with diarrhoea in Papua New Guinea^{4,12}. Thereafter, endemic cases were also reported in Haiti, Nepal, Peru and linked to Travellers Diarrhoea in the 1980s, where it was reported variously as *Cryptosporidium muris*-like, a flagellate, an unsporulated coccidian, a large *Cryptosporidium*, a blue-green alga (cyanobacterium-like body), or a coccidian-like body^{4,30}. By 1990 the new pathogen was associated with chronic diarrhoea in acquired immunodeficiency syndrome patients³¹. Nevertheless, it was not until the 1990s that food and water-borne outbreaks in North America grabbed public and scientific attention to the emergence of the disease, after a food or waterborne outbreak of 21 was reported in a Chicago teaching hospital in 1990³².

This fuelled efforts to definitively identify and characterise the pathogen, with the first breakthrough occurring in 1992 when Ortega *et al.*, reported they had sporulated and excysted the oocysts, thus ending speculation on the classification of the parasite and placing it in the genus *Cyclospora*³³. The same group finally identified and characterised the causative agent as the coccidian *Cyclospora cayetanensis* (Apicomplexa:

Eimeriidae) using classical morphological methods^{1, 30}. However, molecular techniques have placed the pathogen closer to *Eimeria* species, suggesting its classification as such^{34, 35}. Today, *C. cayetanensis* transmission in developed countries through food and water borne outbreaks has been well documented and understood, however the modes of transmission and risk factors in endemic areas remain poorly understood^{4, 36, 37}.

Life cycle and Biology

Cyclospora cayetanensis is an obligate intracellular parasite that is normally found in the jejunum. Oocysts found in the stool can be mistaken for the common *Cryptosporidium* spp. infecting humans, but can easily be distinguished by their larger size (8-10 μm) using an ocular micrometer³⁸. The life cycle begins with the ingestion of the sporulated oocyst in contaminated water or food. Unlike *Cryptosporidium*, this oocyst when freshly passed in the stool is not sporulated and therefore not infective^{1, 30}. Oocysts require a few days to weeks, depending on climatic factors, to develop and mature in the environment into the infective sporulated oocyst, thus precluding direct fecal-oral transmission^{1, 38}. Temperatures ranging between 25-30°C are most suitable for sporulation. Upon being ingested, the oocysts excyst in the gut, releasing the sporozoites, which proceed to invade the epithelial cells of the small intestine^{39, 40}. The sporozoites undergo two generations of asexual reproduction in the cell, whereby they form meronts that contain numerous merozoites⁴⁰. The first generation meronts have 8-12 merozoites, whilst the second have only 4 merozoites, which penetrate new cells to form gametes. Some of these gametes enlarge to form the female macrogametes and some microgametes that undergo meiosis to form numerous flagellated sperm-like microgametes. The mature microgametes leave the microgametocyte and migrate to fertilize the macrogamete. The development of a resilient oocyst wall around the zygote, which contains 2 sporocysts or small oocysts, makes up the unsporulated oocyst. Eventually the oocyst is passed out with stool to begin the process of sporulation. Sporogony commences in the presence of higher atmospheric oxygen concentrations and is complete between 7-12 days⁴⁰⁻⁴³.

Clinical aspects of *C. cayetanensis* infection

Presentation:

The clinical presentation of *C. cayetanensis* may include gastrointestinal (GI) symptoms such as loose or watery diarrhoea, nausea, vomiting, abdominal cramps, loss of appetite, or unintentional weight loss; or constitutional symptoms such as fever, chills, muscle

aches, joint aches, generalized body aches, headache, or fatigue. Although asymptomatic infections are known to occur, the onset of symptoms in naïve populations observed in outbreaks is 1-14 days post exposure and is often accompanied by a characteristic waxing and waning of symptoms^{6, 40, 41, 44}. In endemic countries, symptoms begin approximately 5-8 days after and may persist for a month or more. However, infection without watery diarrhoea is a common occurrence and has been observed in Haiti among other areas⁴⁵.

Diagnosis

Detection is based on the identification of oocysts in stool specimens using modified acid fast staining, or hot safranin test²¹. Where cyclosporiasis is suspected, up to 3 stool specimens taken 2 days apart should be tested to rule out the parasite. A rapid method of identification is possible on wet mounts using fluorescent microscopy employing a filter with a wavelength in the range of 340-380 nm, which reveals the bright, pale blue oocysts glow³⁰. Polymerase chain methods (PCR) have also been developed for diagnosis and detection in the environment, but the primers appears to cross react with *Eimeria* spp.^{26, 35, 46}.

Differential diagnosis of cyclosporiasis includes all other causes of diarrhoea. Infections with *Giardia*, *Cryptosporidium*, *Isospora*, *Toxoplasma* and *Microsporidia* should be suspected in cases of persistent diarrhoea that does not respond to the usual treatment^(6, 20, 31, 36, 41, 47). Cyclosporiasis can also present symptoms closely resembling celiac disease and irritable bowel syndrome⁴⁸.

Sequelae

Like cryptosporidiosis, *C. cayetanensis* infection has also been associated with the onset of various extra-intestinal complications that may accompany prolonged infection, especially in the HIV immunocompromised patients. These include acalculous cholecystitis¹⁹; biliary disease¹⁸, Guillian-Barré syndrome¹⁷, and Reiter or reactive arthritis syndrome¹⁶. One case of *C. cayetanensis* oocysts found in the sputum of 60-year-old HIV negative male with a history of successfully treated TB has been reported in Argentina⁴⁹, and a similar case has also been reported in Egypt where a 45 year-old HIV negative male with TB history also presented oocysts in the sputum accompanied with active TB infection⁵⁰. The latter has prompted calls for the inclusion of cyclosporiasis as a new causative agent of respiratory disease and as a differential diagnosis for TB. In both TB negative and positive cases, the patients presented with loss of weight, cough with expectoration of purulent sputum and

dyspnea. The implications of pulmonary infection to the biology, life cycle and transmission of *C. cayetanensis* remains to be evaluated, as does its interaction with TB, a disease currently on the increase in Africa due to the HIV/AIDS pandemic⁵¹.

Treatment

Symptoms resolve with the administration of Trimethoprim-Sulfamethoxazole, the standard treatment for cyclosporiasis at 160/800 mg oral dose taken twice a day for 7 days or 160/800 mg oral dose taken 4 times a day for 10 days in immunocompromised patients with AIDS often with symptoms resolving and passage of oocysts in the stool ceasing within 24-48 hr^{52, 53}.

Epidemiology in endemic setting

The first epidemiological studies conducted in an area endemic for *C. cayetanensis*, were carried out in 1997-1998 in Guatemala where raspberry exports to USA were linked to cyclosporiasis outbreaks¹⁰. Children of age groups 1.5 to 4 years and 5 to 9 years were found to be 5 times more likely to be positive for *C. cayetanensis* in stool samples than adults. Overall infection did not differ significantly by sex. The infection rates were also found to fluctuate seasonally, with prevalence peaking in the rainy season (6.7%) and falling to undetectable levels in the dry seasons. Drinking of untreated water or swimming in rivers or springs, having a septic tank as opposed to municipal sewage, direct contact with the soil and ownership of dogs, chicken or other fowls, were all found to significantly increase risk of infection although cats and pigs did not¹⁰.

Subsequent epidemiological studies in endemic areas of Haiti, Nepal and Peru have corroborated higher infection rates in children, seasonal fluctuation of prevalence, and risk associated with water source and ownership of domestic animals, particularly fowl, guinea pigs and rabbits^{37, 54}. However no oocysts were detected in a survey for *C. cayetanensis* in domestic animals carried out for 1.5 years in an endemic area in Haiti, ruling out domestic animals as a reservoir²⁴. Studies in Peru have also shown that previous infection offers significant protection against subsequent exposures⁵⁴. This consolation is however short lived as clinical sequelae may still be linked with asymptomatic infection. Humans are therefore the only known host for *C. cayetanensis*, but the eating of raw or insufficiently cooked bivalves, which are filter feeders that concentrate pathogens from waters, has been shown to be a risk factor⁵⁵.

Reports of *Cyclospora* spp. infecting non-human primates in the East African countries of Uganda, Kenya, Tanzania and Ethiopia have led to efforts to conduct

studies of non-human primates in order to better understand the epidemiology and ecology of endemic cyclosporiasis^{25, 56-59}. Host specificity of *Cyclospora* species in the different non-human primates, geographic overlaps of both monkey and parasite species notwithstanding, corroborated the likelihood of humans as the sole definitive host of *C. cayetanensis*⁵⁸. However the collection of oocyst positive stools from the studied troops of vervet monkeys, baboons and colobus monkeys at all times of the year in Kenya, which is notable for extreme weather patterns, demonstrated a lack of seasonality in infection contrary to expectations⁵⁸.

Cyclospora the enigma

It is clear that the epidemiology of cyclosporiasis as we know it today is a study in contradictions, as the causative agent itself remains shrouded in mystery. How can a pathogen incapable of direct faecal-oral infection and lacking a zoonotic reservoir or ubiquitous presence in the environment³⁸, be responsible for so many widespread outbreaks of gastroenteritis all over the world? Trace amounts of oocyst contamination in food products indicate that the infective dose of *C. cayetanensis* is likely to be very low³⁸. However attempts at experimental infection in both humans and animals have all proved futile^{25, 60}. The lack of an animal model has therefore made it hard to determine the viability of oocyst exposed to varying environment conditions that may impact its transmission³⁸. While most outbreaks have been linked to food sources, just as many unexplained sporadic cases linked to water/sewage transmission have been reported in the developed countries^{32, 38, 61}. These cases may signify ubiquitous endemicity, remaining undetected due to seasonality of infection, but by what means the parasite survives from one infection season to the next remains unknown.

Even more puzzling is the dearth of *C. cayetanensis* reports in African populations and as a cause of traveller's diarrhoea in visitors returning from the continent. The screening of over 4,800 samples using modified acid fast staining, and other concerted efforts in previous years have all failed to determine the presence of *C. cayetanensis* in Kenya⁶². In addition to economic restraints for effective routine diagnosis in health centres and national/regional surveillance, it has been postulated that the amount of sulpha drugs administered in sub Saharan Africa as drugs of choice for malaria and other infectious diseases may suppress the parasite, humans being the only reservoir³⁶. Could sulfadoxine-pyrimethamine (SP), the now largely ineffective but still affordable treatment of malaria, be our unlikely saviour? Or is it the use of trimethoprim (Septrin), which is widely

used as a first line of antibiotics and in management of HIV before ARVs? Significantly higher HIV and malaria prevalence in sub-Saharan Africa may account for a higher drug pressure than other parts of the world, however, the withdrawal of SP as a first line antimalarial due to resistance and similar use of Septrin in *C. cayetanensis* endemic Peru and Haiti where 5-11% HIV prevalence have been documented, makes the dearth of reported African cyclosporiasis all the more mysterious.

The controversy and questions surrounding *C. cayetanensis* are no less intriguing. Whereas there is compelling molecular evidence for its reclassification as a mammalian *Eimeria* species, only *C. cayetanensis* DNA sequences and *C. colobi*, *C. cercopithecii* and *C. papionis* from East African non-human primates are available^{27, 34, 36, 57}. The way forward on the classification of this parasite in the light of these new developments remains uncharted waters. The biology of *C. cayetanensis* in the host is no less intriguing. The presence of oocysts in HIV negative TB patients strongly suggests the existence of an extra-intestinal parasite lifecycle and colonization in humans that is yet unknown^{49, 50}. Moreover, more studies are required to determine whether *C. cayetanensis* interacts with TB or has only been observed in TB patients because it is an acid-fast staining organism like the TB bacillus. Extra-intestinal stages in the life cycle of *Cyclospora talpae* in European moles have been demonstrated in the liver where sexual stages develop in the bile duct epithelial cells^{63, 64}. *C. cayetanensis* has similarly been shown to cause biliary disease in HIV/AIDS patients suggesting possible colonization of the liver^{18, 36}. Unfortunately, the lack of animal/experimental models remains an impediment to resolving these questions.

Conclusions and recommendations

Cyclosporiasis therefore presents an ever-broadening frontier cutting across multiple disciplines of research including clinicians, epidemiologists, parasitologists, veterinarians, environmental scientists among others. As a food-borne disease that impacts on horticulture (the backbone of most developing country economies) and primarily infects children and HIV/AIDS patients, routine surveillance in both the population and environment should be carried out. Surveillance in populations should include sputum samples regardless of TB or HIV status. The role of its interaction with TB and/or HIV should also be elucidated. Use of non-pathogen *Cyclospora* spp. such as non-human primates cyclosporiasis and *C. talpae* and mole hosts should be developed as alternative animal models for the study of extra-intestinal colonization and disease. Additionally,

the possible role of sulphur drugs in control of cyclosporiasis should be evaluated, in order to address the impact of the disease in Africa.

Acknowledgements

This work was funded by the Centre for Microbiology Research and is published with the permission of Director, Kenya Medical Research Institute.

References

- Ortega YR, Gilman RH, Sterling CR. A new coccidian parasite (Apicomplexa: Eimeriidae) from humans. *J. Parasitol.* 1994;80:625-629.
- Curry A, Smith HV. Emerging pathogens: *Isospora*, *Cyclospora* and *Microsporidia*. *Parasitol* 1998;117:S143-S159.
- Drenaggi D, Cirioni O, Giacometti A, Fiorentini A, Scalise G. Cyclosporiasis in a traveler returning from South America. *J. Travel Med.* 1998;5(3):153-155.
- Herwaldt BL. *Cyclospora cayetanensis*: a review, focusing on the outbreaks of cyclosporiasis in the 1990s. *Clin. Infect. Dis.* 2000;31(4):1040-1057.
- Ho AY, Lopez AS, Eberhart MG, Levenson R, Finkel BS, da Silva AJ, et al. Outbreak of cyclosporiasis associated with imported raspberries, Philadelphia, Pennsylvania, 2000. *Emerging Infect. Dis.* 2002;8(8):783-786.
- Mansfield LS, Gajadhar AA. *Cyclospora cayetanensis*, a food- and waterborne coccidian parasite. *Vet. Parasitol.* 2004;126(1-2):73-90.
- Alakpa GE, Clarke SC, Fagbenro-Beyioku AF. Cyclospora cayetanensis infection: vegetables and water as possible vehicles for its transmission in Lagos, Nigeria. *Br. J. Biomed. Sci.* 2003;60(2):113-114.
- Crist A, Morningstar C, Chambers R, Fitzgerald T, Stoops D, Deffley M, et al. Outbreak of cyclosporiasis associated with snow peas — Pennsylvania, 2004. *MMWR* 2004;53(37):876-878.
- Shields JM, Olson BH. *Cyclospora cayetanensis*: a review of an emerging parasitic coccidian. *Int. J. Parasitol.* 2003;33(4):371-391.
- Bern C, Hernandez B, Lopez MB, Arrowood MJ, de Mejia MA, de Merida AM, et al. Epidemiologic studies of *Cyclospora cayetanensis* in Guatemala. *Emerging Infect. Dis.* 1999;5(6):766-774.
- El-Karamany EM, Zaher TI, el-Bahnasawy MM. Role of water in the transmission of cyclosporiasis in Sharkia Governorate, Egypt. *J Egypt Soc Parasitol.* 2005;35(3):953-962.
- Ashford RW. Occurrence of an undescribed coccidian in man in Papua New Guinea. *Ann. Trop. Med. Parasitol.* 1979;73:497-500.
- Alakpa GE, Clarke SC, Fagbenro-Beyioku AF. *Cyclospora cayetanensis* infection in Lagos, Nigeria. *Clin Microbiol Infect.* 2003;9(7):731-733.
- Cegielski JP, Ortega YR, McKee S, Madden JF, Gaido L, Schwartz DA, et al. Cryptosporidium, enterocytozoon, and cyclospora infections in pediatric and adult patients with diarrhea in Tanzania. *Clin. Infect. Dis.* 1999;28(2):314-321.
- Nassef NE, el-Ahl SA, el-Shafee OK, Nawar M. Cyclospora: a newly identified protozoan pathogen of man. *J Egypt Soc Parasitol.* 1998;28(1):213-219.
- Connor B, Johnson E, Soave R. Reiter syndrome following protracted symptoms of *Cyclospora* infection. *Emerging Infect. Dis.* 2001;7(3):453-454.
- Richardson RF, Remler BF, Katirji B, Murad MH. Guillain-Barre syndrome after *Cyclospora* infection. *Muscle Nerve* 1998(669-671).
- Sifuentes-Osornio J, Porras-Cortes G, Bendall RP, Morales-Villarreal F, Reyes-Teran G, Ruiz-Palacios GM. *Cyclospora cayetanensis* infection in patients with and without AIDS: biliary disease as another clinical manifestation. *Clin. Infect. Dis.* 1995;29:613-616.
- Zar FA, El-Bayoumi E, Yungbluth MM. Histological proof of acalculous cholecystitis due to *Cyclospora cayetanensis*. *Clin. Infect. Dis.* 2001;33:e140-e141.
- Mohle-Boetani JC, Werner SB, Waterman SH, DJ V. The impact of health communication and enhanced laboratory-based surveillance on detection of cyclosporiasis outbreaks in California. *Emerging Infect. Dis.* 2000;6(2):200-203.
- Eberhard ML, Pieniazek NJ. Laboratory diagnosis of *Cyclospora* infections. *Arch. Pathol. Lab. Med.* 1997;121(8):792-797.
- Kimura K, Kumar Rai S, Takemasa K, Ishibashi Y, Kawabata M, Belosevic M, et al. Comparison of three microscopic techniques for diagnosis of *Cyclospora cayetanensis*. *FEMS Microbiol Lett.* 2004;238(1):263-266.
- Dowd SE, John D, Eliopolus J, Gerba CP, Naranjo J, Klein R, et al. Confirmed detection of *Cyclospora cayetanensis*, *Encephalitozoon intestinalis* and *Cryptosporidium parvum* in water for drinking. *J. Water Health* 2003;1(3):117-123.
- Eberhard ML, Nace EK, Freeman AR. Survey for *Cyclospora cayetanensis* in domestic animals in an endemic area in Haiti. *J. Parasitol.* 1999;85(3):562-563.
- Eberhard ML, Ortega YR, Hanes DE, Nace EK, Do RQ, Robl MG, et al. Attempts to establish experimental *Cyclospora cayetanensis* infection in laboratory animals. *J. Parasitol.* 2000;86(3):577-582.
- Shields JM, Olson BH. PCR-Restricted Fragment Length Polymorphism method for detection of *Cyclospora cayetanensis* in environmental waters without microscopic confirmation. *Applied and Environmental Microbiology* 2003;69(8):4662-4669.
- Lainson R. The Genus *Cyclospora* (Apicomplexa: Eimeriidae), with a description of *Cyclospora schneideri* n.sp. in the snake *Anilius scytale scytale* (Aniliidae) from Amazonian Brazil - a review. *Mem. Inst. Oswaldo Cruz* vol.100 no.2 2005;100(2).
- Schaudinn F. Studien über krankheitserregende Protozoen I. *Cyclospora caryolitica* Shaud., der Erreger der perniciosen Enteritis des Maulwurfs. *Arch. K. Gesundheitsamte* 1902;18:378-416.
- Schneider A. Sur les psorospermies oviformes ou coccidies. Espécies nouvelles ou peu connues. *Arch. Zool. Exp. Gen.* 1881;9:387-404.
- Ortega YR, Sterling CR, Gilman RH, Cama VH, Diaz F. *Cyclospora* species - a new protozoan pathogen of humans. *N. Engl. J. Med.* 1993;328:1308-1312.
- Hart AS, Ridinger MT, Soundarajan R, Peters CS, Swiatlo AL, FE K. Novel organisms associated with chronic diarrhoea in AIDS. *Lancet* 1990;335:169-170.
- Huang P, Weber JT, Sosin DM, Griffin PM, Long EG, Murphy JJ, et al. The first reported outbreak of a diarrheal disease associated with *Cyclospora* in the United States. *Ann. Intern. Med.* 1995;123:409-414.
- Ortega YR, Sterling CR, Gilman RH, Carna VA, Diaz F. *Cyclospora cayetanensis*: a new protozoan pathogen of humans. In: The 41st Annual Meeting of the American So-

- ciety of Tropical Medicine and Hygiene; 1992: J Am Soc Trop Med Hyg; 1992. p. 210.
34. Pieniazek NJ, Herwaldt BL. Reevaluating the molecular taxonomy: is human-associated *Cyclospora* a mammalian *Eimeria* species? *Emerg. Infect. Dis.* 1997;3:381-383.
 35. Relman DA, Schmidt TM, Gajadhar A, Sogin M, Cross J, Yoder K, et al. Molecular phylogenetic analysis of *Cyclospora*, the human intestinal pathogen, suggests that it is closely related to *Eimeria* species. *J. Infect. Dis.* 1996;173:440-445.
 36. Eberhard ML, Arrowood MJ. *Cyclospora* spp. *Curr. Opin. Infect. Dis.* 2002;15(5):519-522.
 37. Lopez SL, Bendik JM, Alliance JY, Roberts JM, daSilva AJ, Moura INS, et al. Epidemiology of *Cyclospora cayetanensis* and other intestinal parasites in a community in Haiti. *J. Clin. Microbiol.* 2003;41(5):2047-2054.
 38. Sterling CR, Ortega YR. *Cyclospora*: An Enigma Worth Unraveling. *Emerg. Infect. Dis.* 1999;5(1):48-53.
 39. Bendall RP, Lucas S, Moody A, Tovey G, Chiodini PL. Diarrhoea associated with cyanobacterium-like bodies: a new coccidian enteritis of man. *Lancet* 1993;341:590-592.
 40. Ortega Y, Nagle R, Gilman RH, Watanabe J, Miyagui J, Kanugusuku P. Pathologic and clinical findings in patients with cyclosporiasis and a description of intracellular parasite life-cycle stages. *J. Infect. Dis.* 1997;176:1584-1589.
 41. Connor BA, Reidy J, Soave R. Cyclosporiasis: clinical and histopathological correlates. *Clin. Infect. Dis.* 1999;28(12):1216-1222.
 42. Sun T, Ilardi CF, Asnis D, Bresciani AR, Goldenberg S, Roberts B, et al. Light and electronmicroscopic identification of *Cyclospora* species in the small intestine: evidence of the presence of asexual life cycle in human host. *Am. J. Clin. Pathol.* 1996;105:216-220.
 43. Tran Van Nhieu J, Nin F, Fleury-Feith J, Chaumette MT, Schaeffer A, Bretagne S. Identification of intracellular stages of *Cyclospora* species by light microscopy of thick sections using hematoxylin. *Hum. Pathol.* 1996;27:1107-1109.
 44. Ayala-Gaytan JJ, Diaz-Olachea C, Riojas-Montalvo P, Palacios-Martinez C. Cyclosporidiosis: clinical and diagnostic characteristics of an epidemic outbreak. *Rev. Gastroenterol. Mex.* 2004;69(4):226-229.
 45. Eberhard ML. *Cyclospora cayetanensis* infections in Haiti: a common occurrence in the absence of watery diarrhea. *Am. J. Trop. Med. Hyg.* 1999;60(4):584-586.
 46. Pieniazek NJ, Slemenda SB, da Silva AJ, Alfano EM, Arrowood MJ. PCR confirmation of infection with *Cyclospora cayetanensis*. *Emerg. Infect. Dis.* 1996;2:357-359.
 47. Rezk H, el-Shazly AM, Soliman M, el-Nemr HI, Nagaty IM, Fouad MA. Coccidiosis among immuno-competent and -compromised adults. *J Egypt Soc Parasitol.* 2001;31(3):823-834.
 48. Vidyut Pingé-Suttor, Chris Douglas, Antony Wettstein. *Cyclospora* infection masquerading as coeliac disease. *MJA* 2004;180(6):295-296.
 49. Di Gliullo AB, Cribari MS, Bava AJ, Cicconetti JS, Collazos R. *Cyclospora cayetanensis* in sputum and stool samples. *Rev Inst Med Trop Sao Paulo* 2000;42(2):115-117.
 50. Hussein EM, Abdul-Manaem AH, el-Attary SL. *Cyclospora cayetanensis* oocysts in sputum of a patient with active pulmonary tuberculosis, case report in Ismailia, Egypt. *J Egypt Soc Parasitol.* 2005;35(3):787-793.
 51. World Health Organization. *Global Tuberculosis Control: Surveillance, Planning, Financing.* WHO Report 2004. Geneva, Switzerland: WHO; 2004. Report No.: WHO/CDS/TB/2004.331.
 52. Hoge CW, Shlim DR, Ghimire M, Rabold JG, Pandey P, Walch A, et al. Placebo-controlled trial of co-trimoxazole for *Cyclospora* infections among travellers and foreign residents in Nepal. *Lancet* 1995;345:691-693.
 53. Madico G, Gimán RH, Miranda E, Cabrera L, Sterling CR. Treatment of *Cyclospora* infections with co-trimoxazole. *Lancet* 1993;342:122-123.
 54. Bern C, Ortega Y, Checkley W, Roberts JM, Lescano AG, Cabrera L, et al. Epidemiologic differences between cyclosporiasis and cryptosporidiosis in Peruvian children. *Emerging Infect. Dis.* 2002;8(6):581-585.
 55. Negm AY. Human pathogenic protozoa in bivalves collected from local markets in Alexandria. *J Egypt Soc Parasitol.* 2003;33(3):991-998.
 56. Ashford RW, Warhurst DC, Reid GD. Human infection with cyanobacterium-like bodies. *Lancet* 1993;341:1034.
 57. Eberhard ML, da Silva AJ, Lilley BG, Pieniazek NJ. Morphological and molecular characterization of new *Cyclospora* species from Ethiopian monkeys: *C. cercopitheci* sp.n., *C. colobi* sp.n., and *C. papioni* sp.n. *Emerging Infect. Dis.* 1999;5(5):651-658.
 58. Eberhard ML, Njenga MN, DaSilva AJ, Owino D, Nace EK, Won KY, et al. A survey for *Cyclospora* spp. in Kenyan primates, with some notes on its biology. *J. Parasitol.* 2001;87(6):1394-1397.
 59. Smith HV, Paton CA, Girdwood RW, Mtambo MM. *Cyclospora* in non-human primates in Gombe, Tanzania. *Vet. Rec.* 1996;138(21):528.
 60. Alfano-Sobsey EM, Eberhard ML, Seed JR, Weber DJ, Won KY, Nace EK, et al. Human challenge pilot study with *Cyclospora cayetanensis*. *Emerging Infect. Dis.* 2004;10(4):726-729.
 61. Wurtz R. *Cyclospora*: a newly identified intestinal pathogen of humans. *Clin Infect Dis* 1994;18:620-623.
 62. Gatei W, Wamae CN, Mbae C, Waruru A, Mulinge E, Waithera T, et al. Cryptosporidiosis: prevalence, genotype analysis, and symptoms associated with infections in children in Kenya. *Am J Trop Med Hyg* 2006;75(1):78-82.
 63. Mohamed HA, Molyneux DH. Developmental stages of *Cyclospora talpae* in the liver and bile duct of the mole *Talpa europaea*. *Parasitol* 1990;101:345-350.
 64. Pellérdy L, Tanyi Z. *Cyclospora talpae* sp.n. (Protozoa: Sporozoa) from the liver of *Talpa europaea*. *Folia Parasitol (Praha)* 15: 275-277 1968;15:275-277.