

## CLINICAL PROFILE AND PATTERN OF DEMENTIA IN A GERIATRIC CENTRE

O.O. Elugbadebo<sup>1</sup>, T.H. Farombi<sup>2</sup>, O.A. Afolabi<sup>3</sup>, L.A. Adebusoye<sup>2</sup>

1. Department of Psychiatry, College of Medicine, University of Ibadan, Nigeria.
2. Chief Tony Anenih Geriatric Centre, University College Hospital, Ibadan, Nigeria.
3. Department of Psychiatry, University College Hospital, Ibadan, Nigeria.

*Correspondence:*

**Dr. O. Elugbadebo**

Department of Psychiatry,  
College of Medicine,  
University of Ibadan,  
Ibadan, Nigeria.  
E-mail: fisayoelu08@yahoo.com

*Submission Date: 16th April, 2023*

*Date of Acceptance: 1st April, 2024*

*Publication Date: 30th April, 2024*

### ABSTRACT

**Background:** Knowledge on the clinical presentation of dementia is essential for appropriate care, especially in Low-and-Middle-Income Countries where these cases are on a sharp rise and can also aid early detection of other underlying conditions.

This study sought to provide a broad and updated socio-demographic, clinical profile, pattern of diagnosis and treatment features of people diagnosed with dementia in this setting.

**Method:** A retrospective cohort study which reviewed the medical case records of all older adults with dementia receiving treatment at the psychogeriatric and the neurology clinic of the Geriatric Centre (N=192). A proforma was designed to collect information from the case records.

**Results:** The mean ( $\pm$ SD) age of the participants was 74.0( $\pm$ 7.2) years, 97.9% lived with other persons, 50.0% had at least one comorbidity and 52.6% presented late for treatment. Overall, hypertension (64.1%) and diabetes (22.4%) were the most common comorbidity, 55.2% had complaints bordering on behavioural problems; irrational speech (31.3%) being the most common, while 91.7% had forgetfulness as a cognitive symptom.

**Conclusions:** A high rate of comorbidities, as well as late presentation was common among the participants. Our findings appraise the clinical importance of detailed knowledge of the patterns and profiles of older adults with dementia for early presentation and treatment.

**Keywords:** Dementia, Older adults, Clinical profile, Clinical pattern, Comorbidity, Behavioural symptoms

### INTRODUCTION

The prevalence of dementia is rising globally with the increasing aging population.<sup>1,2</sup> According to a global report in 2019, about 57.4 million people were living with dementia, and this prevalence has been estimated to increase to about 152.8 million cases in 2050.<sup>3</sup> In Nigeria, a systematic review conducted by Adeloye *et al.* in 2019, showed that about 4.9% older adults were living with dementia and projected that by the year 2050, this would have increased to over 1.4 million cases.<sup>4</sup> Of no doubt, this will pose some level of care burden. Knowledge on the clinical presentation of the illness is essential for appropriate clinical care especially in Low-and-Middle-Income Countries where dementia cases are on a sharp rise.<sup>5,6</sup> Evidences have established that identifying the pattern of presentation of people diagnosed with dementia is crucial in making early diagnosis and developing strategies for effective treatment.<sup>7,8</sup> These patterns include the participants' sociodemographic, clinical and pathologic characteristics.<sup>9-11</sup>

To our knowledge, there is little evidence on the presentation and clinical pattern of dementia in Nigeria. Although, there exists some robust studies on its incidence, prevalence and risk factors,<sup>12-14</sup> there is still a dearth of information on the clinical course, pattern of presentation, and specific pointers to diagnosing its subtypes.<sup>15,16</sup>

Dementia awareness is still relatively low with few specialists available,<sup>16,17</sup> leading to a common trend of substantial time lag between identification of symptoms and presentation for treatment among people with dementia (PWD) residing in this setting.<sup>18,19</sup> With the growing number of persons with dementia in LMICs,<sup>9</sup> late identification, under diagnosis and misdiagnosis could become more challenging without adequate knowledge about dementia. Moreover, dementia is usually accompanied with some medical comorbidities<sup>20</sup>; explicit knowledge of the profile and presentation of the disease can aid early detection of other underlying conditions.

Therefore, this study sought to provide a broad and updated sociodemographic and clinical profile, patterns of diagnosis and treatment features of the people diagnosed with dementia in this setting.

## MATERIALS AND METHODS

### Study Design and Population

A retrospective cohort study conducted among PWD receiving treatment at the psychogeriatric and the neurology clinic of the Geriatric Centre. The Geriatric Centre is located within the University College Hospital (UCH), Ibadan, specially established for provision of care to older people. Psychogeriatric and neurological services such as in-patient and outpatient care, as well as daycare services are part of the specialist services rendered in the facility. The data were collected between 26<sup>th</sup> July 2021 and 16<sup>th</sup> September 2021 and the population consisted of older adults attending the Geriatric centre, UCH, Ibadan, Nigeria. Eligible participants were people diagnosed with dementia who had been attending either the psychogeriatric or neurology clinic of the Geriatric centre from 1<sup>st</sup> January 2013 to 31<sup>st</sup> December 2020. The participants included were those who had a dementia diagnosis based on either the International Classification of Diseases 10<sup>th</sup> revision (ICD-10) or Diagnostic Statistical Manual of Mental Disorders IV (DSM 4) criteria documented in their case records and whose hospital records were accessible.

### Data Collection

Medical case records of all older adults attending either of the two clinics (psychogeriatric or Neurology) located within the Geriatric Centre, who were diagnosed with dementia between 1st January 2013 and 31<sup>st</sup> December 2020 were identified. A total of 976 older adults attended both clinics, however only 220 had dementia diagnosis, out of which only 192 case records were available for review. The remaining 28 case records were missing. A proforma was adopted to collect the socio-demographic and clinical details of the participants. The socio-demographic features studied were age, sex, education, occupation,

living conditions, while the clinical details include, presenting complaints at first contact, duration of symptoms, medical history, family history, comorbidity profiles, diagnosis before and after examination, type of treatment and medications being used. The psychiatric diagnoses were based on International Classification of Diseases (ICD-10), Classification of Mental and Behavioural Disorders, Neuropsychological Inventory (NPI) and Mini-Mental State Examination (MMSE), Computed Tomography (CT)/ Magnetic Resonance Imaging (MRI) scans and other laboratory tests. A total sampling technique was used to recruit the participants from both clinics.

### Data Analysis

Descriptive statistics and charts were generated to evaluate distributions of all study variables and to generate summary statistics including means, standard deviations, frequencies, and percentages. These included the profile/pattern of presentation and diagnosis/treatment received. To determine the pattern of diagnoses, the investigations carried out by the participants were summarized into frequencies using descriptive statistics. The data were analysed using Statistical Package for the Social Sciences (SPSS) version 23.0. The value of  $p < .05$  was considered statistically significant.

### Ethical Approval

This study was performed in accordance with the principles stated in the declaration of Helsinki and was approved by the University of Ibadan/University College Joint Ethical Committee (UI/EC/23/0015).

## RESULTS

### Socio demographic and clinical characteristics of dementia

A total of 192 older adults with dementia were identified from the medical case records based on the eligibility criteria. Mean ( $\pm$ SD) age of the participants was 74.0 ( $\pm$ 7.2) years and 56.2% were males. Majority of the participants were educated (82.8%), 67.2% married, 97.9% lived with other persons and the



Figure 1: Duration of symptoms among behavioural and cognitive cases

majority were referred from the general outpatient unit of the Geriatric centre for cognitive evaluation (90.6%). (Table 1).

**Table 1:** Socio-demographic characteristics of participants (N=192)

Characteristics	N	%
<b>Age (Years)</b>		
<70	54	28.1
70-79	96	50.0
≥80	42	21.9
Mean(±S.D)	74.0(±7.2)	
<b>Sex</b>		
Male	108	56.2
Female	84	43.8
<b>Educational status</b>		
No formal education	33	17.2
Educated	159	82.8
<b>Living arrangement</b>		
Living alone	4	2.1
Living with others	188	97.9
<b>Marital status</b>		
Currently not married	63	32.8
Currently married	129	67.2
<b>Source of referral</b>		
<sup>a</sup> GOP Geriatric centre	174	90.6
<sup>b</sup> GOPD UCH (main hospital)	6	3.1
Private hospital	3	1.6
<sup>c</sup> Others	9	4.7

<sup>a</sup>General Outpatient unit of the Geriatric Centre

<sup>b</sup>General Outpatient Department, University College Hospital

<sup>c</sup>Others- Family Physician, (referral from doctors)


Over 76.0% of the participants had at least one comorbidity and 52.6% presented late for treatment (>1year). Overall, hypertension (64.1%) and diabetes (22.4%) were the most common comorbid illness and 24.0% had no comorbidity. Furthermore, 66.1% of the participants dropped out of treatment, and 25.5% received pharmacotherapy type of treatment. At the

end of this study period, 62.5% participants were alive and 37.5% dead. (Table 2).

Based on the presenting complaints, 55.2% of the participants had complaints bordering on behavioural problems; irrational speech (31.3%) was the most common behavioural symptom listed. Almost all

**Table 2:** Clinical characteristics and management of older people with dementia (N = 192)

Variables	N	%
<b>Presenting Complaints</b>		
Behavioural	12	6.3
Cognitive	68	35.4
Both	112	58.3
<b>Had Comorbidities</b>		
Yes	146	76.0
No	46	24.0
<b>List of comorbidities</b>		
None	46	24.0
Hypertension	123	64.1
Diabetics	43	22.4
Stroke	4	2.1
Glaucoma	9	4.7
Asthma	5	2.6
Others	13	6.8
<b>Type(s) of treatment received</b>		
Pharmacotherapy	49	25.5
Psychotherapy	11	5.7
Both	132	68.8
<b>Place of care</b>		
Outpatient	186	96.9
Inpatient	6	3.1
<b>Period of presentation to the clinic</b>		
Late presentation (>1year)	101	52.6
Early presentation (≤1year)	91	47.4
<b>Status</b>		
Alive	120	62.5
Dead	72	37.5
<b>Stay in treatment</b>		
Regular	65	33.9
Dropped out	127	66.1



**Figure 2:** Frequency of investigations carried out by the participants

participants had forgetfulness (91.7%) as a cognitive symptom (Table 3)

While majority of our participants had late onset dementia, a very few cases of young onset dementia were observed similar to reports from other studies.<sup>28,29</sup>

**Table 3:** Behavioural and psychological symptoms found among the participants

Presenting complaints	Yes, n (%)	No, n (%)
<b>Cognitive</b>		
Forgetfulness	176 (91.7)	16 (8.3)
Inability to carry out daily activities	30 (15.6)	162 (84.4)
<b>Behavioural</b>		
Irrational speech/Talking to self	60 (31.3)	132 (68.7)
Odd behaviour	30 (15.6)	162 (84.4)
Wandering	24 (12.5)	168 (87.5)
Sleeping disorders	19 (9.9)	173 (90.1)
Defecating/urinating on self	17 (8.9)	175 (91.1)
Delusion	14 (7.3)	178 (92.7)
Aggression	9 (4.7)	183 (95.3)
Hallucination	12 (6.3)	180 (93.7)
Mood change	5 (2.6)	187 (97.4)

Among those with behavioural complaints, late presentation and dropout rate was found at 57.3% and 59.3% respectively. The mean duration of symptoms presentation for behavioural is 27months and for cognitive, 24months (Fig 1).

In reference to the laboratory tests carried out, 49.0% performed Full Blood Count (FBC), 59.9% Urinalysis, 68.8% Fasting Lipid Profile, 55.2% Mini Mental State Examination and 31.3% Neuroimaging (Fig 2).

## DISCUSSION

### Profiles of older adults with dementia

A male predominance was noted in the study, which contradicts the findings of several demographic studies that have reported a high prevalence of dementia among the females.<sup>1,21,22</sup> Globally, women have a higher life expectancy than men<sup>23</sup> which contributes to the gender differences in dementia prevalence and incidence.<sup>23</sup> However, it is perceived that the life expectancy gap is gradually closing, and both genders may be at equal risk in some years ahead.<sup>24</sup> Alternatively, the high educational attainment observed among the male participants may have informed the high prevalence of the gender in this study. Consistently, past studies have also shown that highly educated individuals have positive health seeking behaviors<sup>25</sup> and are more inclined to seek specialized services especially for health-related issues than the less educated<sup>26</sup> which can explain our findings as we had more educated males in our study.

Majority of the PWD were between the ages of 70-79 years similar to the findings from other studies.<sup>4,27</sup>

Age has been reported to be the greatest risk factor for dementia<sup>30,31</sup> and people above 65 years are the most affected<sup>2</sup> as this age group have a high propensity for deterioration in intellectual functioning.<sup>32-35</sup> Reports from previous studies conducted have also shown that most people present or are mostly diagnosed with dementia at age 65 and over.<sup>36,37</sup> Another plausible explanation is that dementia symptoms are relatively mild at the initial stages and may be attributed to normal aging, hence, people seek help at the latter stages when the symptoms begin to have significant impact on their daily functioning and quality of life. Notably, over 97% of our participants live with other persons especially their family members who are mostly the caregivers. Dementia as a chronic condition that affects functioning<sup>18,38</sup> underscores the need for either an informal or formal care.<sup>39,40</sup> In Nigeria, the multigenerational living is still common<sup>41</sup> and has served the purpose of providing informal caregiving to PWD. This is possibly a reflection of the cultural beliefs in our society where filial relationship and respect for the elderly are considered among the highest virtues, and certain stigma are associated with moving older ones/parents to a nursing home; such children are perceived as irresponsible. Notably, Ogunniyi *et al.* in a study on caring for persons living with dementia in Nigeria, reported that most patients live with their family and relatives further affirming the prominent familial connections that exists in this part of the world.<sup>42</sup> However, the mass exodus of the younger generation to inner cities and more developed countries is gradually causing a shift from informal care to formal caregiving and institutionalized care.<sup>41</sup>

### Presenting complaints of participants

In the present study, forgetfulness was a common presenting complaint similar to results from some previous studies.<sup>43,44</sup> Dröes *et al.* in their study on Memory problems in dementia confirmed forgetfulness as one of the first identified symptoms in most forms of dementia.<sup>43,45</sup>

In addition to this finding, a larger percentage of our cohort presented with at least one behavioural and psychological symptom of dementia (BPSD) with irrational speech being the most common. Previous reports confirmed that about 90% of people living with dementia commonly present with behavioural problems at the onset and during the course of the illness.<sup>46,47</sup> This is very important in diagnosis, treatment and outcome as BPSD has been shown to be associated with high burden of care<sup>48</sup> and psychological problems in caregivers<sup>49</sup> with increased mortality rates<sup>50</sup> and reduced quality of life.<sup>51</sup> It also makes up a crucial aspect of dementia and a major reason for presentation in hospitals.<sup>47</sup>

### Comorbidities

In keeping with findings from literature, hypertension and diabetes were the most common comorbid medical conditions observed in our study. Hypertension and diabetes mellitus have been considered to be related to cognitive dysfunction and brain pathology<sup>52-54</sup> and are major risk factors for dementia.<sup>55</sup> Chen *et al.*, in a cross-sectional study on comorbidity and dementia among persons living with dementia in Taiwan found that hypertension and diabetes were the most common underlying conditions among persons with dementia. Similarly, Ogunniyi *et al.* found that most PWD are commonly diagnosed with hypertension, a comorbidity mostly attributed to old age.<sup>42</sup> A six-year follow-up study among PWD and people without dementia in Nigeria showed that the group with dementia exhibited a notably higher prevalence of hypertension compared to the group without dementia.<sup>56</sup> Notably, persons with these medical conditions are predisposed to cognitive decline and dementia.<sup>20</sup> Moreover, there is increasing evidence that dementia is associated with cardiovascular diseases.<sup>57</sup> Studies have shown that having a comorbidity often delays dementia diagnosis and significantly affects the outcomes of treatment<sup>58</sup>, increases the risk of dying,<sup>59</sup> reduces quality of life<sup>60</sup> and greater use of health care services. Primary family caregivers of older adults with dementia often experience high caregiver burden and significant decline in a range of health outcomes.<sup>61</sup>

### Time of presentation and treatment outcome

It is notable that a larger fraction of the participants presented symptoms late similar to what obtains in

previous studies.<sup>62,63</sup> A significant time gap of symptom presentation was found between cognitive and behavioral complaints. The time to presentation of participants with behavioural symptoms was shorter compared to those who had cognitive symptoms. This could be explained by the normalcy bias that forgetfulness in older people is a normal part of the aging process which in most cases lead to the dismissal of seeking care for the cognitive problems. Nakahori *et al.* similarly affirmed that the normalcy bias can lead both older adults and their family members to dismiss the possibility of a seeming case of dementia.<sup>43</sup> However, many begin seeking help when BPSD become disruptive and burdensome<sup>64</sup>. Furthermore, due to the progressive nature of the disease, dementia symptoms are usually mild and overlooked at the initial stages. The negative perceptions of dementia in our setting further contributes to the delay in seeking treatment.<sup>17</sup>

Although studies have shown that behavioral symptoms are one of the major reasons for seeking dementia care.<sup>46,47</sup> We found that most of the participants in our study, especially those with behavioral symptoms, dropped out after initial consultation. However, we can speculate that caregivers tend to withdraw from subsequent appointments after resolution of behavioural symptoms with treatment. Moreover, in some situations, this incidence may have occurred when caregivers become aware that the illness can only be managed and not cured. Family caregivers may be discouraged by the costs of medications, frequent consultations and additional costs of co-managing dementia with other comorbidities evidenced in the increased number of investigative procedures.<sup>65</sup> Majority of the medical costs in this setting are covered by the out-of-pocket system especially for those who are not under any insurance scheme.<sup>61</sup> In keeping with this finding, a study conducted by Elugbadebo *et al.* on accessibility and discontinuity of treatment among older people receiving care in a Geriatric facility, postulated that among several other constraints, older adults drop out after their first appointments due to financial constraints.<sup>66</sup>

The results of the clinical investigations showed a relative comprehensive approach to diagnosing dementia. Notably, over half of the patients had a comprehensive cognitive assessment necessary in diagnosing dementia. Similarly, a higher fraction of the participants had investigations to assess cardiovascular risk factors. Remarkably, only one-third of the cases underwent neuroimaging. While this is relatively low, neuroimaging such as CT scans and MRI, can be very expensive considering that treatment is

usually by out-of-pocket payment and health insurance does not cover full cost for the very few patients that are on the National Health Insurance Scheme.

### Limitation

This study was explorative; thus, conclusions have to be drawn cautiously. Secondly, the sample size was relatively small limiting further sub analysis of some important variables.

### CONCLUSION

The current study emphasises the range of symptoms that are often present among people living with dementia. Unlike several other studies, male gender was predominant in the sample and most participants live with others. A high rate of medical comorbidities (mostly hypertension) and dropouts, as well as late presentation was observed among the participants. Those who presented with late symptoms had cognitive complaints. Our findings appraise the clinical importance of detailed knowledge of the pattern and profile of older adults with dementia for early presentation and treatment.

### Recommendation

The study emphasises the need for the evaluation of the patterns and profile of presentation among a larger sample of older adults with dementia to achieve proper treatment of dementia; increasing the quality of life among the elderly residing in LMICs. There is a need for more dementia awareness with the aim of improving early presentation. We, therefore, recommend more wholistic care in managing dementia.

**Acknowledgement:** The authors would like to appreciate Chiamaka Okwudiri for helping with logistics for this study.

**Funding details:** The authors received no funding for this study

**Declaration of interest statement:** The authors declare no conflict of interest

### REFERENCES

1. **Prince M**, Bryce R, Albanese E, *et al.* The global prevalence of dementia: A systematic review and metaanalysis. *Alzheimers Dement* . 2013 ;9(1):63-75.e2.
2. **Prince M.** Epidemiology of dementia. *Psychiatry*. 2007 Dec 1;6(12):488–90. Available from: <https://www.sciencedirect.com/science/article/pii/S1476179307002133>
3. **Nichols E**, Steinmetz JD, Vollset SE, *et al.* Estimation of the global prevalence of dementia in 2019 and forecasted prevalence in 2050: an analysis for the Global Burden of Disease Study 2019. *Lancet Public Health*. 2022 Feb 1;7(2):e105–125.
4. **Adeloye D**, Auta A, Ezejimofor M, *et al.* Prevalence of dementia in Nigeria: a systematic review of the evidence. *J Glob Health Rep*. 2019;3:e2019014. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6746335/>
5. **Prince MJ**, Acosta D, Castro-Costa E, *et al.* Packages of Care for Dementia in Low- and Middle-Income Countries. *PLOS Med*. 2009 Nov 3;6(11):e1000176. Available from: <https://journals.plos.org/plosmedicine/article?id=10.1371/journal.pmed.1000176>
6. **Uwakwe R**, Prince M. Identifying Dementia And Depression In The Elderly: The Role of the Family. *Niger J Clin Pract*. 2005;7(2):82–87. Available from: <http://www.ajol.info/index.php/njcp/article/view/11188>
7. **Maxwell CJ**, Vu M, Hogan DB, *et al.* Patterns and Determinants of Dementia Pharmacotherapy in a Population-Based Cohort of Home Care Clients. *Drugs Aging*. 2013 Jul 1;30(7):569–585.
8. **Mughal AY**, Devadas J, Ardman E, *et al.* A systematic review of validated screening tools for anxiety disorders and PTSD in low to middle income countries. *BMC Psychiatry [Internet]*. 2020 Dec;20(1):338. Available from: <https://bmcp psychiatry.biomedcentral.com/articles/10.1186/s12888-020-02753-3>
9. **Amoo G**, Ogundele AT, Olajide AO, *et al.* Prevalence and pattern of psychiatric morbidity among community-dwelling elderly populations in Abeokuta, Nigeria. *J Geriatr Psychiatry Neurol [Internet]*. 2020 Nov 1;33(6):353–362.
10. **Jelastopulu E**, Giourou E, Argyropoulos K, *et al.* Demographic and Clinical Characteristics of Patients with Dementia in Greece. *Adv Psychiatry*. 2014 Nov 17;2014:e636151. Available from: <https://www.hindawi.com/journals/apsy/2014/636151/>
11. **Almeida OP**, Hill K, Howard R, *et al.* Demographic and clinical features of patients attending a memory clinic. *Int J Geriatr Psychiatry*. 1993;8(6):497–501.
12. **Hendrie HC**, Osuntokun BO, Hall KS, *et al.* Prevalence of Alzheimer's disease and dementia in two communities: Nigerian Africans and African Americans. *Am J Psychiatry*. 1995 Oct;152(10):1485–1492.
13. **Olayinka OO**, Mbuyi NN. Epidemiology of Dementia among the Elderly in Sub-Saharan Africa. *Int J Alzheimerx2019s Dis*. 2014 Aug

- 6;2014:e195750. Available from: <https://www.hindawi.com/journals/ijad/2014/195750/>
14. **Adeloye D**, Auta A, Ezejimofor M, *et al.* Prevalence of dementia in Nigeria: a systematic review of the evidence. *J Glob Health Rep.* 2019;3:e2019014. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6746335/>
  15. **Uwakwe R**. The pattern of psychiatric disorders among the aged in a selected community in Nigeria. *Int J Geriatr Psychiatry.* 2000 Apr;15(4):355–62.
  16. **Baiyewu O**, Bello B. Psychogeriatrics and Psychogeriatric Services in Nigeria. *Niger J Psychiatry.* 2015;13(2):2–6. Available from: <https://www.ajol.info/index.php/njpsyc/article/view/125671>
  17. **Adebiyi AO**, Fagbola MA, Olakehinde O, Ogunniyi A. Enacted and implied stigma for dementia in a community in south-west Nigeria. *Psychogeriatrics.* 2016;16(4):268–273.
  18. **Nguyen T**, Li X. Understanding public-stigma and self-stigma in the context of dementia: A systematic review of the global literature. *Dementia.* 2020 Feb 1;19(2):148–181.
  19. **Brecher DB**, West TL. Underrecognition and Undertreatment of Pain and Behavioral Symptoms in End-Stage Dementia. *Am J Hosp Palliat Med.* 2016 Apr 1;33(3):276–280.
  20. **Chen TB**, Yiao SY, Sun Y, *et al.* Comorbidity and dementia: A nationwide survey in Taiwan. *PLOS ONE.* 2017 Apr 12;12(4):e0175475. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0175475>
  21. **Cao Q**, Tan CC, Xu W, *et al.* The Prevalence of Dementia: A Systematic Review and Meta-Analysis. *J Alzheimers Dis.* 2020 Jan 1;73(3):1157–1166. Available from: <https://content.iospress.com/articles/journal-of-alzheimers-disease/jad191092>
  22. **Savva GM**, Arthur A. Who has undiagnosed dementia? A cross-sectional analysis of participants of the Aging, Demographics and Memory Study. *Age Ageing.* 2015 Jul 1;44(4):642–647.
  23. **Liu Y**, Arai A, Kanda K, *et al.* Gender gaps in life expectancy: generalized trends and negative associations with development indices in OECD countries. *Eur J Public Health.* 2013 Aug; 23(4): 563–568.
  24. **McIntyre N**. When will men live as long as women? By 2032, say experts. *The Guardian.* 2018 Feb 13; <https://www.theguardian.com/inequality/2018/feb/13/when-will-men-live-as-long-as-women-by-2032-say-experts>
  25. **Mackenzie CS**, Gekoski WL, Knox VJ. Age, gender, and the underutilization of mental health services: The influence of help-seeking attitudes. *Aging Ment Health.* 2006 Nov 1 ;10(6):574–582. Available from: <https://doi.org/10.1080/13607860600641200>
  26. **Schmidt H**, Wild EM, Schreyögg J. Explaining variation in health information seeking behaviour – Insights from a multilingual survey. *Health Policy.* 2021 May 1;125(5):618–26. Available from: <https://www.sciencedirect.com/science/article/pii/S0168851021000269>
  27. **Bader S**, Ellouz E, Abderrahim R, *et al.* Socio-demographic and clinical features of patients with dementia attended in the psychiatry department. *Eur Psychiatry.* 2021 Apr;64(S1):S744–S744. Available from: <https://www.cambridge.org/core/journals/european-psychiatry/article/sociodemographic-and-clinical-features-of-patients-with-dementia-attended-in-the-psychiatry-department/A010E5036D4AEBFB6BB8D628D56DD0EA>
  28. **Wong JFW**, Kirk A, Perlett L, *et al.* Characteristics of Young-Onset and Late-Onset Dementia Patients at a Remote Memory Clinic. *Can J Neurol Sci.* 2020 May;47(3):320–327. Available from: <https://www.cambridge.org/core/journals/canadian-journal-of-neurological-sciences/article/characteristics-of-youngonset-and-lateonset-dementia-patients-at-a-remote-memory-clinic/3540B81EFAB18492E0C098E65B6BBFCC>
  29. **Papageorgiou SG**, Kontaxis T, Bonakis A, *et al.* Frequency and Causes of Early-onset Dementia in a Tertiary Referral Center in Athens. *Alzheimer Dis Assoc Disord.* 2009 Dec;23(4):347–351. Available from: [https://journals.lww.com/alzheimjournal/Abstract/2009/1000/Frequency\\_and\\_Causes\\_of\\_Early\\_onset\\_Dementia\\_in\\_a.8.aspx](https://journals.lww.com/alzheimjournal/Abstract/2009/1000/Frequency_and_Causes_of_Early_onset_Dementia_in_a.8.aspx)
  30. **Mielke MM**. Sex and Gender Differences in Alzheimer's Disease Dementia. *Psychiatr Times.* 2018 Nov;35(11):14–17. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6390276/>
  31. **Abbott A**. Dementia: A problem for our age. *Nature.* 2011 Jul;475(7355):S2–4. Available from: <https://www.nature.com/articles/475S2a>
  32. Organization WH, International AD. Dementia: a public health priority. World Health Organization, World Health Organization; 2012 Jun. Available from: <https://apo.org.au/node/307503>
  33. **Farina N**, Suemoto CK, Burton JK, *et al.* Perceptions of dementia amongst the general public across Latin America: a systematic review. *Aging Ment Health.* 2021 May 4;25(5):787–96. Available from: <https://doi.org/10.1080/13607863.2020.1725738>
  34. **Wortmann M**. Dementia: a global health priority - highlights from an ADI and World Health Organization report. *Alzheimers Res Ther.* 2012

- Sep 21;4(5):40. Available from: <https://doi.org/10.1186/alzrt143>
35. **Perkins AJ**, Hui SL, Ogunniyi A, *et al.* Risk of mortality for dementia in a developing country: the Yoruba in Nigeria. *Int J Geriatr Psychiatry*. 2002;17(6):566–573. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1002/gps.643>
  36. **Cao Q**, Tan CC, Xu W, *et al.* The Prevalence of Dementia: A Systematic Review and Meta-Analysis. *J Alzheimers Dis*. 2020 Jan 1;73(3):1157–1166. Available from: <https://content.iospress.com/articles/journal-of-alzheimers-disease/jad191092>
  37. **Di Carlo A**, Baldereschi M, Amaducci L, *et al.* Incidence of Dementia, Alzheimer's Disease, and Vascular Dementia in Italy. The ILSA Study. *J Am Geriatr Soc*. 2002;50(1):41–8. Available from: <https://onlinelibrary.wiley.com/doi/abs/10.1046/j.1532-5415.2002.50006.x>
  38. **McParland P**, Devine P, Innes A, Gayle V. Dementia knowledge and attitudes of the general public in Northern Ireland: an analysis of national survey data. *Int Psychogeriatr*. 2012 Oct; 24(10):1600–13. Available from: <https://www.cambridge.org/core/journals/international-psychogeriatrics/article/abs/dementia-knowledge-and-attitudes-of-the-general-public-in-northern-ireland-an-analysis-of-national-survey-data/8EE9DFA07D85BEC3504D53B0078B5388>
  39. **Bieber A**, Nguyen N, Meyer G, Stephan A. Influences on the access to and use of formal community care by people with dementia and their informal caregivers: a scoping review. *BMC Health Serv Res*. 2019 Feb 1;19(1):88. Available from: <https://doi.org/10.1186/s12913-018-3825-z>
  40. **Hoe J**, Hancock G, Livingston G, Orrell M. Quality of life of people with dementia in residential care homes. *Br J Psychiatry*. 2006 May;188(5):460–464. Available from: <https://www.cambridge.org/core/journals/the-british-journal-of-psychiatry/article/quality-of-life-of-people-with-dementia-in-residential-care-homes/B3ED49D72E73F43A589EE6EECB355E7>
  41. **Baiyewu O**, Bella AF, Adeyemi JD, *et al.* Health problems and socio-demographic findings in elderly Nigerians. *Afr J Med Med Sci*. 1997 Jun; 26(1–2):13–17.
  42. **Ogunniyi A**, Hall KS, Baiyewu O, *et al.* Caring for individuals with dementia: the Nigerian experience. *West Afr J Med*. 2005 Sep;24(3):259–262.
  43. **Nakahori N**, Sekine M, Yamada M, *et al.* Discrepancy in the perception of symptoms of cognitive decline between older adults and their family members: results of the Toyama dementia survey. *BMC Neurol*. 2019 Dec 26;19(1):338. Available from: <https://doi.org/10.1186/s12883-019-1581-2>
  44. **Moylett S**, Price A, Cardinal RN, *et al.* Clinical Presentation, Diagnostic Features, and Mortality in Dementia with Lewy Bodies. *J Alzheimers Dis*. 2019 Jan 1;67(3):995–1005. Available from: <https://content.iospress.com/articles/journal-of-alzheimers-disease/jad180877>
  45. **Dröes RM**, van der Roest HG, van Mierlo L, Meiland FJM. Memory problems in dementia: adaptation and coping strategies and psychosocial treatments. *Expert Rev Neurother*. 2011 Dec;11(12):1769–81; quiz 1782.
  46. **Majer R**, Simon V, Csiba L, *et al.* Behavioural and psychological symptoms in neurocognitive disorders: Specific patterns in dementia subtypes. *Open Med*. 2019 Jan 1;14(1):307–16. Available from: <https://www.degruyter.com/document/doi/10.1515/med-2019-0028/html>
  47. **Cerejeira J**, Lagarto L, Mukaetova-Ladinska E. Behavioral and Psychological Symptoms of Dementia. *Front Neurol*. 2012;3:73. Available from: <https://www.frontiersin.org/article/10.3389/fneur.2012.00073>
  48. **Savva GM**, Zaccari J, Matthews FE, *et al.* Prevalence, correlates and course of behavioural and psychological symptoms of dementia in the population. *Br J Psychiatry*. 2009 Mar;194(3):212–9. Available from: <https://www.cambridge.org/core/journals/the-british-journal-of-psychiatry/article/prevalence-correlates-and-course-of-behavioural-and-psychological-symptoms-of-dementia-in-the-population/5C45CC9AA7FB7AEBC9B8DA4C7664BEBE>
  49. **Kim B**, Noh GO, Kim K. Behavioural and psychological symptoms of dementia in patients with Alzheimer's disease and family caregiver burden: a path analysis. *BMC Geriatr*. 2021 Mar 5;21(1):160.
  50. **Bränsvik V**, Granvik E, Minthon L, *et al.* Mortality in patients with behavioural and psychological symptoms of dementia: a registry-based study. *Aging Ment Health*. 2021 Jun;25(6):1101–9.
  51. **Poon E**. A Systematic Review and Meta-Analysis of Dyadic Psychological Interventions for BPSD, Quality of Life and/or Caregiver Burden in Dementia or MCI. *Clin Gerontol*. 2022 Sep;45(4):777–797.
  52. **Nagai M**, Hoshida S, Kario K. Hypertension and Dementia. *Am J Hypertens*. 2010 Feb 1;23(2):116–124.
  53. **Biessels GJ**, Staekenborg S, Brunner E, *et al.* Risk of dementia in diabetes mellitus: a systematic review. *Lancet Neurol*. 2006 Jan 1;5(1):64–74.



Available from: <https://www.sciencedirect.com/science/article/pii/S1474442205702842>

54. **Deckers K**, Schievink SHJ, Rodriquez MMF, *et al.* Coronary heart disease and risk for cognitive impairment or dementia: Systematic review and meta-analysis. *PLOS ONE*. 2017 Sep 8;12(9):e0184244. Available from: <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0184244>
55. **Faraco G**, Iadecola C. Hypertension. *Hypertension*. 2013 Nov 1;62(5):810–7. Available from: <https://www.ahajournals.org/doi/full/10.1161/HYPERTENSIONAHA.113.01063>
56. **Ogunniyi A**, Lane KA, Baiyewu O, *et al.* Hypertension and incident dementia in community-dwelling elderly Yoruba Nigerians: Hypertension and dementia in elderly Yoruba. *Acta Neurol Scand*. 2011 Dec;124(6):396–402. Available from: <https://onlinelibrary.wiley.com/doi/10.1111/j.1600-0404.2011.01491.x>
57. **Bunn F**, Burn AM, Goodman C, *et al.* Comorbidity and dementia: a scoping review of the literature. *BMC Med*. 2014 Oct 31;12(1):192. Available from: <https://doi.org/10.1186/s12916-014-0192-4>
58. **Schubert CC**, Boustani M, Callahan CM, *et al.* Comorbidity Profile of Dementia Patients in Primary Care: Are They Sicker? *J Am Geriatr Soc*. 2006;54(1):104–109.
59. **Rajamaki B**, Hartikainen S, Tolppanen AM. The effect of comorbidities on survival in persons with Alzheimer's disease: a matched cohort study. *BMC Geriatr*. 2021 Mar 9;21(1):173. Available from: <https://doi.org/10.1186/s12877-021-02130-z>
60. **Nelis SM**, Wu YT, Matthews FE, *et al.* The impact of co-morbidity on the quality of life of people with dementia: findings from the IDEAL study. *Age Ageing*. 2019 May;48(3):361–367. Available from: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC6503940/>
61. **Uwakwe R**, Ibeh CC, Modebe AI, *et al.* The Epidemiology of Dependence in Older People in Nigeria: Prevalence, Determinants, Informal Care, and Health Service Utilization. A 10/66 Dementia Research Group Cross-Sectional Survey. *J Am Geriatr Soc*. 2009;57(9):1620–1627.
62. **Lim J**, Goh J, Chionh HL, Yap P. Why do patients and their families not use services for dementia? Perspectives from a developed Asian country. *Int Psychogeriatr*. 2012 Oct;24(10):1571–1580.
63. **Parker M**, Barlow S, Hoe J, Aitken L. Persistent barriers and facilitators to seeking help for a dementia diagnosis: a systematic review of 30 years of the perspectives of carers and people with dementia. *Int Psychogeriatr*. 2020 Feb 6;1–24.
64. **Shaji KS**, George RK, Prince MJ, Jacob KS. Behavioral symptoms and caregiver burden in dementia. *Indian J Psychiatry*. 2009 Mar;51(1):45–49. Available from: [https://journals.lww.com/indianjpsychiatry/Fulltext/2009/51010/Behavioral\\_symptoms\\_and\\_caregiver\\_burden\\_in.10.aspx](https://journals.lww.com/indianjpsychiatry/Fulltext/2009/51010/Behavioral_symptoms_and_caregiver_burden_in.10.aspx)
65. **Annear MJ**, Lucas P. Dementia in a regional hospital setting: contextual challenges and barriers to effective care. *Int J Ageing Later Life*. 2018 Aug 13;12(1):91–119. Available from: <https://ijal.se/article/view/1293>
66. **Elugbadebo O**, Ojagbemi A, Adefolarin A, Gureje O. Access and Discontinuity of Care at an Outpatient Mental Health Service for Older People in South Western Nigeria. *Community Ment Health J*. 2021 Nov;57(8):1518–1524.