



# ANATOMICAL INVESTIGATION ON THE CRANIOFACIAL AND OCULAR MORPHOMETRICS OF THE MALE WESTERN CATTLE EGRET (*Bubulcus ibis*)

Azeez Idris Ayodeji<sup>1\*</sup>, Ekeolu Oyetunde Kazeem<sup>2</sup>, Usende Ifukibot Levi<sup>3</sup>, Adejumobi Olumuyiwa Abiola<sup>4</sup> and Adetogun Adefunke<sup>5</sup>

<sup>1</sup>Department of Veterinary Anatomy, University of Jos, Nigeria; <sup>2</sup>Department of Veterinary Anatomy, University of Benin, Nigeria; <sup>3</sup>Department of Veterinary Anatomy, University of Abuja, Nigeria; <sup>4</sup>Department of Veterinary Medicine, University of Ibadan, Nigeria; <sup>5</sup>Department of Veterinary Anatomy, University of Ibadan, Nigeria.

Correspondence to: Dr. Azeez Idris Ayodeji; Department of Veterinary Anatomy, University of Jos, PMB 2084, Zip Code 930001, Plateau State, Nigeria. Email: [idris.azeez505@gmail.com](mailto:idris.azeez505@gmail.com); Phone no: +2348034493343.

## ABSTRACT

The heron species, Cattle egret (*Bubulcus ibis*) can be found all over the world. In the South-Western region of Nigeria, they are colonial, white in colour, and are seen in open lands and around water bodies. Morphometric data are important parameter in the assessment of functional morphology, hence we here present for the first time, data on the craniofacial indices of cattle egret. The cranium, nasal, orbital and beak morphometric indices and correlation data of the cattle egret were investigated on the fresh specimen and macerated skull of the male cattle egrets, using the GraphPad Prism. The height of the cranium was  $33.92 \pm 0.970$  cm, while the cranium length and breadth were  $52.92 \pm 1.800$  and  $34.48 \pm 1.714$ , respectively. The right eye socket length,  $36.17 \pm 5.636$  and breadth,  $31.67 \pm 2.160$  were slightly greater than the left eye socket length,  $35.92 \pm 4.128$  and breadth,  $31.00 \pm 2.757$ . There was a strong positive correlation between the right eye socket breadth and the left eye socket breadth ( $r=0.940$ ), and also, a very strong positive correlation was found between the right and the left eye socket length ( $r=0.981$ ). Results showed the foramen magnum length and breadth to be  $19.17 \pm 0.753$  and  $18.83 \pm 0.753$ , respectively, however, a low positive correlation was recorded between the foramen magnum length and breadth ( $r=0.059$ ). A strong positive correlation was observed between the upper beak length and the lower beak length ( $r=0.901$ ), while similar values were recorded for the nasal breadth and length, respectively. Data generated from this study will prove useful in comparative, regional and clinical anatomy and could also help in identifying archaeological remains of the cattle egrets.

**Keywords:** cattle egret, cranium, orbit, beak, nasal, morphometrics, craniofacial indices

DOI: <https://dx.doi.org/10.4314/aja.v11i1.2>

## INTRODUCTION

The cattle egret was first described by Carl Linnaeus in his *Systema naturae* (Linnaeus 1758). Cattle egret (*Bubulcus ibis*) is a cosmopolitan species of heron and is present in worldwide (Rezk 2015). Geographically, the cattle egret has two races which are classified as, the western cattle egret, *B. ibis*, and eastern cattle egret, *B. coromandus*, and these two

forms were described by McAllan et al. (1988). In the South Western part of Nigeria, cattle egrets are colonial, white in colour, and are found in open lands and also, around water bodies. The cattle egret has widely been described based on the color of their plumage, habitat, and feeding habits (Hasan 2015). This bird is of huge economic importance in the

control of ectoparasites of cattle as well as land pests (Hussein and Rezk 2016). While detailed account of the fowl skeleton (Getty 1975) and wings and pelvic bones anatomy of emu (Kumar and Singh 2014), buzzard (Atalar et al. 2007) and kit (Sharma and Dubal 2018; Tiwari et al. 2011) have been described, studies on the gross morphometry of the skeleton of cattle egret are scarce and the scarcity applied to evolutionary study of the bird morphology (Ekeolu et al. 2016). The descriptive anatomy on its appendicular (Rezk 2015a) and axial skeleton (Rezk 2015b) has been reported. Also, Ekeolu et al., (2016) reported data on comparative gross morphometrics of the forelimb and hind limb

skeleton. More recently, Sasan et al. (2019) published data on the gross and morphometrical studies on the humerus of cattle egret. Morphometrical studies of the skull not only reflect the genetic and eco-phenotypic variation of individuals and animals but also provide foundations for clinical and surgical stereotaxic practices (Wehausen and Ramey 2000), hence, we here provide for the first time, baseline information on gross morphometric data on the cranium, nasal and orbital bones and the beak of *B. ibis*, thereby adding to the limited body of knowledge on its anatomy.

## MATERIALS AND METHODS

Six (6) apparently healthy adult male migrant cattle egrets line-trapped at the University of Ibadan Teaching and Research Farm were used for this study. The birds were euthanized by lethal injection using a combination of xylazine (10 mg/kg) and ketamine (100 mg/kg). They were then decapitated at the level of the atlanto-occipital joint. The organs in the head were removed. The tongue was carefully dissected from the mouth and eyes were removed from the socket by stitching the upper eye lid with the lower one and then pulling the eye by the thread as described by Olopade et al. (2011), then carefully dissect around the rim of the socket. The heads were then de-fleshed as much as possible using a scalpel blade. Cold water maceration method of Ekeolu et al. (2016) was then used to prepare the skull and the mandible. Briefly, the egrets were soaked in cold water with ammonium solution and sodium hydroxide overnight to remove grease and soften the connective tissues; meninges (dura mater) of the brain, and muscular attachment on the bones of the head. The solution was changed daily, removing the dissolved brains and tissues. This was repeated for a week and extraneous tissues on the bones were picked using hand (thumb) forceps, after brushing the muscle fibers and connective tissues attached to the bones with sponges. Then, each skull and mandible were then washed in clean water and

air dried. Digital Vernier caliper was engaged in taking the linear measurement of the various bones of the skull, the volume was measured by filling the cranium up with grains. The quantities of grains that fill up each cranium were quantified in a measuring cylinder in millimeters. Figures 1 and 2 show the *B. ibis* skull with illustrations of methodologies employed in the measurement, and Tables 1-6 show the morphometric indices of measured parameters. Measured parameters are concisely defined below.

### ***Definition of Parameters: Cranium and foramen magnum***

1. Cranium top length (**CTL**) - dorsal cranial length, anterior tip of the occipital condyle to the craniofacial hinge (beginning of the upper jaw).
2. Cranium top breadth (**CTB**) - dorsal cranial breadth, the distance between the lateral orbital borders.
3. Cranium back (**CB**) - length from the lateral sides of the occipital crest
4. Whole length of cranium to beak length (**LCBL**) - dorsal anterior tip of occipital condyle to the anterior tip of premaxillary (beak)
5. Height of cranium (**CH**) - dorsal frontal surface to the post orbital process

6. Cranium flat top (**CFT**) - dorsal frontal length
7. Foramen magnum length (**FML**)
8. Foramen magnum breadth (**FMB**)

**Eye socket parameters**


9. Left eye socket length (**ESLL**) - orbital length of the left eye
10. Left eye socket breadth (**ESLB**) - orbital breadth of the left eye
11. Right eye socket length (**ESRL**) - orbital length of the right eye
12. Right eye socket breadth (**ESRB**) - orbital breadth of the right eye
13. Eye socket and beak breadth (**ESBB**) - highest point of orbit to articular surface of the lower jaw
14. Eye socket and upper beak length (**ESBL**) – post-frontal process of orbit to tip of premaxillary of upper jaw
15. Eye socket to nose length (upper) (**ESNL**) – post-frontal process of orbit to nasal length end

**Beak and nose parameters**


16. Beak whole length- upper jaw (**BLU**) - craniofacial hinge to tip of premaxillary.

17. Beak whole length- lower jaw (**BLL**) - articular to tip of lower jaw
18. Beak breadth lower beak- lower jaw (**BBL**) - height of dentary.
19. Gap between lower beak (**BLG**) - posterolateral width at the jaw articulation and the width between the middle of the lower jaw.
20. Upper beak breadth (**UBB**) - breadth of craniofacial hinge to tip of upper jaw.
21. Beak flat top (**BFT**) - dorsal surface of the craniofacial hinge and dorsal beak width.
22. Nose breadth (**NB**) - nasal breadth
23. Nose length (**NL**) - nasal length
24. Nose to end of the upper beak (**NUB**) - nasal length to end of premaxillary of the upper jaw

All numerical data generated from the morphometric studies were subjected to statistical analysis using Student's t-test and correlation analysis with the use of GraphPad Prism (GraphPad Prism 5.04, GraphPad Software, Inc., La Jolla, CA, USA) and expressed as mean±standard deviation (SD). Values of  $p < 0.05$  were considered statistically significant



**Fig. 1:** dorsal view of the skull and beak of the cattle egret with illustrations of some of the measured parameters: Beak whole length- upper jaw (**BLU**) - craniofacial hinge to tip of premaxillary, whole length of cranium to beak length (**LCBL**) - dorsal anterior tip of occipital condyle, upper beak breadth (**UBB**) - craniofacial hinge to tip of upper jaw, cranium top- length (**CTL**) - dorsal cranial length, anterior tip of occipital condyle to the craniofacial hinge (beginning of the upper jaw), cranium top-breadth (**CTB**) - dorsal cranial breadth, distance between the lateral orbital borders, Cranium back (**CB**) - length from the lateral sides of the occipital crest



**Fig. 2:** Left lateral view of the skull and beak of the cattle egret with illustrations of some of the measured parameters: Upper beak length (**BLU**) - craniofacial hinge to tip of upper jaw; nose length (**NL**) - nasal length; nose to end of upper beak (**NUB**) – nasal length to end of premaxillary of the upper jaw; eye socket left length (**ESLL**) - orbital length; eye socket left breadth (**ESLB**) - orbital breadth; height of cranium (**CH**) - dorsal frontal surface to post orbital process; Beak whole length- lower jaw (**BLL**) - articular to tip of lower jaw

## RESULTS

### Cranium and foramen magnum

The skull of the cattle egret has the splanchnocranium comprising mainly the bones of the face and the neurocranium in which the brain is lodged. A prominent orbital cavity clearly demarcates the two parts. As shown in Table 1, the cranium breadth (CTB) which is a measure of the distance between the lateral orbital borders is lower than the cranium length (CTL), measured from the tip of the occipital condyle to the craniofacial hinge ( $34.48 \pm 1.714$  vs  $52.92 \pm 1.800$ ). The foramen magnum length

(FML) is slightly higher than the breadth (FMB) ( $19.17 \pm 0.753$  vs  $18.83 \pm 0.753$ ). The height of cranium (CH) value ( $33.92 \pm 0.970$ ) was similar to cranium flat top (CFT) ( $32.32 \pm 1.933$ ). The whole length of cranium to beak length (**LCBL**) measured ( $112.17 \pm 2.463$ ) while the length of the cranium from the lateral sides of the occipital crest (CB) measured  $31.83 \pm 0.753$ . With Pearson’s correlation, there was a strong positive correlation of CTL vs LCBL (0.940) and CTL vs FML (0.824), as shown in Table 2.

**Table 1:** Mean±SD of cranium indices (n=6)

Parameters	Mean±SD
CTL	52.92±1.800
CTB	34.48±1.714
CB	31.83±0.753
LCBL	112.20±2.463
CH	33.92±0.970
CFT	32.32±1.933
FML	19.17±0.753
FMB	18.83±0.753

**Table 2:** Pearson's correlation coefficient values (r) of cranium indices (n=6)

Parameters	CTL	CTB	CB	LCBL	CH	CFT	FML	FMB
<b>CTL</b>		0.576	-0.307	0.940	0.024	0.635	0.824	-0.307
<b>CTB</b>	0.576		0.199	0.387	0.131	0.665	0.111	-0.654
<b>CB</b>	-0.307	0.199		-0.575	-0.434	0.497	-0.647	-0.412
<b>LCBL</b>	0.940	0.387	-0.575		0.070	0.392	0.953	-0.090
<b>CH</b>	0.024	0.131	-0.434	0.070		-0.186	0.023	-0.434
<b>CFT</b>	0.635	0.665	0.497	0.392	-0.186		0.259	-0.740
<b>FML</b>	0.824	0.111	-0.647	0.953	0.023	0.259		0.059
<b>FMB</b>	-0.307	-0.654	-0.412	-0.090	-0.434	-0.740	0.059	

### Eye socket

Eye socket indices and Pearson's correlation coefficients are shown in Table 3 and 4, respectively. The eye sockets of the cattle egrets are lodged in the splanchnocranium. Right eye socket length (**ESRL**) was slightly larger than the left eye socket length (**ESLL**) ( $36.17 \pm 5.636$  vs  $35.92 \pm 4.128$ ) while value of the left eye socket breadth (**ESLB**) was very similar right eye socket breadth (**ESRB**) ( $31.00 \pm 2.757$  vs  $31.67 \pm 2.160$ ). With Pearson's correlation, there

was a strong positive correlation between **ESLB** and **ESRB** (0.940), and also a strong positive correlation was found between **ESLL** and **ESRL** (0.981). The **ESBB** had a negative correlation with the **ESRB** (-0.133) and a very low negative correlation with the **ESRL** (-0.078), while a low positive correlation was recorded with the **ESLB** and **ESLL** (0.148 vs 0.060). **ESBL** was negatively correlated with **ESLB** (-0.474) and **ESRB** (-0.521) while it has strong positive correlations with **ESLL** (0.874) and **ESRL** (0.817).

**Table 3:** Mean $\pm$ SD of eye socket indices (n=6)

Parameters	Mean $\pm$ SD
<b>ESLB</b>	31.00 $\pm$ 2.757
<b>ESLL</b>	35.92 $\pm$ 4.128
<b>ESRL</b>	36.17 $\pm$ 5.636
<b>ESRB</b>	31.67 $\pm$ 2.160
<b>ESBB</b>	36.33 $\pm$ 1.966
<b>ESBL</b>	96.50 $\pm$ 2.757
<b>ESNL</b>	58.00 $\pm$ 2.074

**Table 4:** Pearson's correlation coefficient values (r) of eye socket indices (n=6)

Parameters	ESLB	ESLL	ESRL	ESRB	ESBB	ESBL	ESNL
<b>ESLB</b>		-0.782	-0.862	0.940	0.148	-0.474	-0.560
<b>ESLL</b>	-0.782		0.981	-0.845	0.060	0.874	0.888
<b>ESRL</b>	-0.862	0.981		-0.865	-0.078	0.817	0.830
<b>ESRB</b>	0.940	-0.845	-0.865		-0.133	-0.521	-0.625
<b>ESBB</b>	0.148	0.060	-0.078	-0.133		0.037	0.184
<b>ESBL</b>	-0.474	0.874	0.817	-0.521	0.037		0.971
<b>ESNL</b>	-0.560	0.888	0.830	-0.625	0.184	0.971	

## Beak and nose

Beak and nose indices and Pearson's correlation coefficients are shown in Table 5 and 6, respectively. There was a low positive correlation between the upper beak length (BLU) and the upper beak breadth (UBB) (0.290). Similar values were recorded for nasal breadth (NB) ( $17.50\pm 0.894$ ) and nasal length (NL) ( $22.83\pm 1.472$ ), with moderate positive

correlation (0.632). The NL was higher than the NB ( $22.83\pm 1.472$  vs  $17.50\pm 0.8944$ ) with a positive correlation. Meanwhile, very similar values were recorded between the BLG and BBL ( $20.70\pm 0.7014$  vs  $20.43\pm 0.5989$ ) with low insignificant correlations (0.224). Also, the NB was recorded to have strong positive correlations with the BLU (0.822) and BLL (0.867), respectively.

**Table 5:** Mean $\pm$ SD of nose and beak indices (n=6)

Parameters	Mean $\pm$ SD
<b>BLU</b>	73.67 $\pm$ 1.633
<b>BLL</b>	102.6 $\pm$ 3.353
<b>BBL</b>	20.43 $\pm$ 0.599
<b>BLG</b>	20.70 $\pm$ 0.701
<b>UBB</b>	20.85 $\pm$ 1.350
<b>BFT</b>	21.53 $\pm$ 0.963
<b>NB</b>	17.50 $\pm$ 0.894
<b>NL</b>	22.83 $\pm$ 1.472
<b>NUB</b>	63.83 $\pm$ 3.296

**Table 6:** Pearson's correlation coefficient values (r) of nose and beak indices (n=6)

Parameters	BLU	BLL	BBL	BLG	UBB	BFT	NB	NL	NUB
<b>BLU</b>		0.901	-0.518	-0.733	0.290	0.187	0.822	0.555	0.322
<b>BLL</b>	0.901		-0.345	-0.591	0.604	0.371	0.867	0.672	0.581
<b>BBL</b>	-0.518	-0.345		0.357	0.109	0.005	-0.597	-0.424	0.398
<b>BLG</b>	-0.733	-0.591	0.357		0.042	0.021	-0.335	-0.562	-0.476
<b>UBB</b>	0.290	0.604	0.109	0.042		0.872	0.629	0.659	0.503
<b>BFT</b>	0.187	0.371	0.005	0.021	0.872		0.534	0.725	0.220
<b>NB</b>	0.822	0.867	-0.597	-0.335	0.629	0.534		0.684	0.153
<b>NL</b>	0.555	0.672	-0.424	-0.562	0.659	0.725	0.684		0.426
<b>NUB</b>	0.322	0.581	0.398	-0.476	0.503	0.220	0.153	0.426	

## DISCUSSION

The nomenclature here adopted in this study for the anatomical descriptions was as described in the *Nomina Anatomica Avium* by Baumel et al. (1993). While there is information on the gross morphological descriptions skulls of the cattle

egrets (Rezk 2015), there is presently a dearth on information concerning morphometric parameters of the skull. For this purpose, some of the craniofacial measurements were investigated. This study also shows the

correlative relationship of cranial, orbital, beak and nasal parametric indices.

Morphometric data (i.e. absolute or relative size of particular interest) are useful barometers of functional morphology (Saber and Gummow 2014; Oyelowo et al. 2017), and we present for the first time, data on the craniofacial indices of cattle egret. In the morphology of avian skulls, their striking spectrum of shapes and relative sizes of the facial skeleton are very evident (Zusi 1993). Thin plates of bone that are formed from connective tissue or cartilaginous templates form the head (König, Korbelt, and Hans-Georg 2016). Birds differ critically in the way the neurocranium is structured which is most visible in the extreme inclination of the nuchal plane in any species (Marugán-Lobón and Buscalioni 2004; 2006).

The frontal and maxillary processes of the premaxillary bone form the boundary of the bony nostril, and the delicate nasal bone, while the caudodorsal angle of the nostril is formed by the premaxillary and maxillary processes of the nasal bone (König, Korbelt, and Hans-Georg 2016). The nasal bones, which are thin flat bones form the dorsolateral boundary of the nasal cavity and form part of the upper beak (Rezk 2015).

In the present study, we showed the cranium breadth and length of the male *B. ibis* to be  $34.48 \pm 1.714$  and  $52.92 \pm 1.800$ , respectively. Darwish et al. (2006) report the length and height of the skull of *E. ibis* as 10.6 and 2.4; *C. coturnix* as 4.1 and 1.4; *M. gallopavo* as 10.1 and 3.2; *A. anser* as 12.5 and 4.4 and *A. atthis* as 9.1 and 1.9, respectively. It is highly expected that the length and breadth of the skull will affect the cranial capacity *vis a vis* the brain capacity. Brain capacity/size in relationship to intelligence in school children (Estabrooks 1928) and animals (Hiecks and Dougherty, 2013) have been studied and in wombats and wallabies could reflect their intelligence for getting food and water, managing territory, offences and in defense (Saber and Gummow 2014) which could also be the case with cattle egret.

Concerning the foramen magnum, our report showed the length and breadth to be  $19.17 \pm 0.753$  and  $18.83 \pm 0.753$ , respectively. Burdan et al. (2012) reported the mean values of the foramen length for Eastern-European adult males versus female (human) ( $37.06 \pm 3.07$  vs.  $35.47 \pm 2.60$  mm), breadth ( $32.98 \pm 2.78$  vs.  $30.95 \pm 2.71$  mm) were significantly higher in males than in females with a significant, positive correlation between foramen length and breadth. The foramen magnum is an important anatomical opening in the base of the skull through which the posterior cranial fossa communicates with the vertebral canal and it also related to a number of pathological conditions including tumors, and occipital dysplasia.

It has been reported that avian species have a very large bony orbit that lies between the neurocranium and splanchnocranium (Nickel, Schummer, and Seiferle 1977; Dyce, Sack, and Wensing 2010) and the position and size varying within the various species reflecting the divergence of their behavior and mode of locomotion (Wiedersheim 1909; Darwish et al. 2006). In this present study, we showed that the mean length and breadth of the right orbit of cattle egret to be  $36.17 \pm 5.636$  and  $31.67 \pm 2.160$ , respectively and the mean length and breadth of the left orbit to be  $35.92 \pm 4.128$  and  $31.00 \pm 2.757$ , respectively. Earlier, Darwish et al. (2006) reported same parameters in various avian species; *E. ibis* as  $2.3 \pm 0.03$  and  $1.7 \pm 0.07$ ; *C. coturnix* as  $1.3 \pm 0.05$  and  $1.0 \pm 0.07$ ; *M. gallopavo* as  $3.6 \pm 0.07$  and  $2.3 \pm 0.07$ ; *A. anser* as  $3.0 \pm 0.01$  and  $2.6 \pm 0.07$  and *A. atthis* as  $2.1 \pm 0.04$  and  $1.4 \pm 0.07$  for length and height of the orbit, respectively.

Evidence had shown that the relative visual field depends on the position and direction of the orbit, the degree of projection of the eye ball beyond the orbital rim and the movement of the animal head (Ibrahim et al. 1990, 1992). Specifically, Ibrahim et al. (1992) mentioned that the relative height of the orbit to the total height of the skull is more dorsal in cattle (63.4-100%)

compared to rabbit (39.4-100%), dog (56-92%) and sheep (46.95-82.6%).

We suggest that the position, height and length of the orbit in relation to the beak in cattle egret strongly influences its vision and feeding habits. This is so as Wiedersheim (1909) had shown a relationship between the size and position of the eyes as well as the orbit and the shape of the beak.

Avian beaks are remarkable for their diversity in shape and size and provides elegant illustrations of the process and power of natural selection (Darwin 1859; Badyaev 2010; Badyaev et al. 2008). Even small differences in beak morphology including its length and width can critically affect what foods are accessible to an individual (Boag and Grant 1981; Benkman and Lindholm 1991; Temeles and Kress 2003) and how they can defend themselves against ectoparasites (Badyaev et al. 2008). Because beaks are under such strong selection pressure, recently, great interests in understanding the mechanisms that control their morphology have been on the increase (Handel et al. 2010) with less attention on its morphometrics especially in cattle egret.

The conformation of the facial skeleton is influenced considerably by the shape and mobility of the beak (König, Korbelt, and Hans-Georg 2016). Changes in key morphometric proteins can result to a broad diversity of beak shapes (Schneider and Helms 2003; Wu et al. 2006; 2004; Campàs et al. 2010), thus affecting its morphometrics. Beak morphometrics can be used to characterize normal and abnormal beak

and also to monitor the development of beak deformities in individual birds through time (Handel et al. 2010). Important information that can be obtained from beak morphometrics could include crossed bite which occurs when the upper and lower beaks are laterally offset from each other and overbite which occurs when upper beak is longer than the lower beak (Handel et al. 2010) among others. In this present study, we showed strong correlations between the ESLL and ESRL; BLU and BLL; BLU and NB; BLL and NB, respectively. This could have potential clinical implications. Earlier, Handel et al., (2010) showed that in adult Black capped chickadees, the upper beak ranged from 6.0 to 8.5mm from nares to tip and average slightly longer in males than in females and the gonyes of the lower beak ranged from 6.0 to 7.5mm not differing by sex, similar to our findings herein reported.

Recent studies of the genetic control of beak development have found that two signaling molecules (bone morphogenic protein 4 and calmodulin) play key roles in determining the depth, width and length of beak (Wu et al., 2004, Abzhanov et al., 2006, Grant et al., 2006). These proteins are active during embryonic craniofacial development and control the outgrowth of the beak primordia (Handel et al. 2010).

In conclusion, our findings have provided baseline information on the Craniofacial Indices of the Male Cattle Egret (*Bubulcus ibis*), which would be of benefit in understanding morphofunctional and paleontological studies of this bird.

**CONFLICT OF INTEREST:** The authors declare that there is no conflict of interest with regards to the publication of this article.

## REFERENCES

1. Atalar, Ömer, İbrahim Kürtül, Derviş Özdemir, Fırat Üniversitesi, Veteriner Fakültesi, Anatomi Anabilim, Dalı Elazığ-Türkiye, et al. 2007. "Morphological and Morphometric Approach to the Bones of the Wings in the Long-Legged Buzzard (*Buteo Rufinus*)."  
<http://www.fusabil.org>.
2. Badyaev, Alexander V. 2010. "The Beak of the Other Finch: Coevolution of Genetic Covariance Structure and Developmental Modularity during Adaptive Evolution." *Philosophical Transactions of the Royal Society B: Biological Sciences*. <https://doi.org/10.1098/rstb.2009.0285>.

3. Badyaev, Alexander V., Rebecca L. Young, Kevin P. Oh, and Clayton Addison. 2008. "Evolution on a Local Scale: Developmental, Functional, and Genetic Bases of Divergence in Bill Form and Associated Changes in Song Structure between Adjacent Habitats." *Evolution* 62 (8): 1951–64. <https://doi.org/10.1111/j.1558-5646.2008.00428.x>.
4. Baumel, Julian J., Anthony S King, James E Breazile, Howard E Evans, and James C. Vanden Berge. 1993. *Handbook of Avian Anatomy: Nomina Anatomica Avium, Second Edition. Publications of the Nuttall Ornithological Club*. <https://doi.org/10.2307/1369201>.
5. Benkman, Craig W., and Anna K. Lindholm. 1991. "The Advantages and Evolution of a Morphological Novelty." *Nature*. <https://doi.org/10.1038/349519a0>.
6. Boag, Peter T., and Peter R. Grant. 1981. "Intense Natural Selection in a Population of Darwin's Finches (Geospizinae) in the Galápagos." *Science*. <https://doi.org/10.1126/science.214.4516.82>.
7. Burdan, F., J. Szumito, J. Walocha, L. Klepacz, B. Madej, W. Dworzański, R. Klepacz, A. Dworzańska, E. Czekajska-Chehab, and A. Drop. 2012. "Morphology of the Foramen Magnum in Young Eastern European Adults." *Folia Morphologica (Poland)*.
8. Campàs, O., R. Mallarino, A. Herrel, A. Abzhanov, and M. P. Brenner. 2010. "Scaling and Shear Transformations Capture Beak Shape Variation in Darwin's Finches." *Proceedings of the National Academy of Sciences of the United States of America*. <https://doi.org/10.1073/pnas.0911575107>.
9. Darwin, Charles. 1859. *On the Origin of the Species, or, the Preservation of Favoured Races in the Struggle for Life*. Darwin. [https://doi.org/10.1016/S0262-4079\(09\)60380-8](https://doi.org/10.1016/S0262-4079(09)60380-8).
10. Darwish et al. 2006. "Some Morphometric Studies on the Orbit of Some Different Birds." *The Egyptian Journal of Experimental Biology (Zoology)* 2 (0): 241–56.
11. Dyce, K. M. (Keith M.), W. O. (Wolfgang O.) Sack, and C. J. G. (Cornelis Johannes Gerardus) Wensing. 2010. *Textbook of Veterinary Anatomy*. Saunders/Elsevier.
12. Ekeolu et al. 2016. "International Journal of Veterinary Science." *International Journal of Veterinary Science* 5 (4): 285–89.
13. Estabrooks, G. H. 1928. "The Relation between Cranial Capacity, Relative Cranial Capacity and Intelligence in School Children." *Journal of Applied Psychology*. <https://doi.org/10.1037/h0070481>.
14. Getty, R. 1975. "Sisson and Grossman's The Anatomy of the Domestic Animals." In *Sisson and Grossman's The Anatomy of the Domestic Animals*.
15. Handel, Colleen M., Lisa M. Pajot, Steven M. Matsuoka, Caroline Van Hemert, John Terenzi, Sandra L. Talbot, Daniel M. Mulcahy, Carol U. Meteyer, and Kimberly A. Trust. 2010. "Epizootic of Beak Deformities Among Wild Birds in Alaska: An Emerging Disease in North America?" *The Auk*. <https://doi.org/10.1525/auk.2010.10111>.
16. Hasan, Mohammad Wasiul. 2015. "Study on Status of Cattle Egret in Bangladesh." Chittagong Veterinary and Animal Sciences University Khulshi, Chittagong-4225, Bangladesh. <http://101.2.160.165:8080/handle/123456789/704>.
17. Hussein, Shaymaa, and Hamdy Rezk. n.d. "Macro and Microscopic Characteristics of the Gastro-Intestinal Tract of the Cattle Egret (Bubulcus Ibis)." *Int J Anat Res* 2016 (2): 2162–74. Accessed February 12, 2020. <https://doi.org/10.16965/ijar.2016.169>.
18. Ibrahim, I.A., M. Taha, A.A. Mansour, and A.T.G. (Assiut Univ. (Egypt). Faculty of Veterinary Medicine) Ahmed. 1992. "Some Anatomical Studies on the Orbit in Dog, Sheep, Cattle and Rabbit." *Benha Veterinary Medical Journal (Egypt)*.
19. König, HE, R Korb, and L Hans-Georg. 2016. *Avian Anatomy Textbook and Colour Atlas*. 5M Publishing.
20. Kumar, Pawan, and Gurdial Singh. 2014. "Gross Anatomy of Wing and Pelvic Limb Bones in Emu (Dromaius Novaehollandiae)."
21. Linnaeus, Carolus. 1758. *Caroli Linnaei Systema Naturae per Regna Tria Naturae: Secundum Classes, Ordines, Genera, Species, Cum Characteribus, Differentiis, Synonymis, Locis. Editio Duodecima. 1. Regnum Animale. 1 & 2 Holmiae, Laurentii Salvii. Holmiae [Stockholm], Laurentii Salvii*.
22. Marugán-Lobón, Jesús, and Ángela D. Buscalioni. 2004. "Geometric Morphometrics in Macroevolution: Morphological Diversity of the Skull in Modern Avian Forms in Contrast to Some Theropod Dinosaurs." In *Morphometrics*, 157–73. Springer Berlin Heidelberg. [https://doi.org/10.1007/978-3-662-08865-4\\_12](https://doi.org/10.1007/978-3-662-08865-4_12).

23. Marugán-Lobón, Jesús, and Ángela D. Buscalioni. 2006. "Avian Skull Morphological Evolution: Exploring Exo- and Endocranial Covariation with Two-Block Partial Least Squares." *Zoology* 109 (3): 217–30. <https://doi.org/10.1016/j.zool.2006.03.005>.
24. McAllan, Ian., Murray D. Bruce, Biocon Research Group., and New South Wales Bird Atlassers. 1988. *The Birds of New South Wales: A Working List*. Biocon Research Group in association with the New South Wales Bird Atlassers.
25. Nickel, R. (Richard), A. (August) Schummer, and E. (Eugen) Seiferle. 1977. *Anatomy of the Domestic Birds*. Parey.
26. Olopade et al. 2011. "Morphometric Studies of the Eyeball and Orbital Region of the Nigerian Local Pig (Sus Scrofa)." *Tropical Veterinarian* 29 (1): 34–40. [https://www.researchgate.net/publication/271457959\\_Morphometric\\_studies\\_of\\_the\\_eyeball\\_and\\_orbita\\_l\\_region\\_of\\_the\\_Nigerian\\_local\\_pig\\_Sus\\_scrofa](https://www.researchgate.net/publication/271457959_Morphometric_studies_of_the_eyeball_and_orbita_l_region_of_the_Nigerian_local_pig_Sus_scrofa).
27. Oyelowo et al. 2017. "Morphological and Craniometric Features of the Skull of African Savanna Hare (Lepus Microtis) Found in North-Central Nigeria." *Journal of Veterinary Anatomy* 10 (2): 85–107. <https://doi.org/10.21608/jva.2017.45447>.
28. Rezk, Hamdy Mahmoud. 2015a. "Anatomical Investigation on the Appendicular Skeleton of the Cattle Egret (Bubulcus Ibis)." *Journal of Experimental and Clinical Anatomy* 14 (1): 5–12. <https://doi.org/10.4103/1596-2393.158919>.
29. Rezk, Hamdy Mahmoud. 2015b. "Anatomical Investigation on the Axial Skeleton of the Cattle Egret, Bubulcus Ibis." *Assiut Vet. Med. J.* 61 (145): 12-21
30. Rezk, Hamdy Mahmoud. 2015. "Anatomical Investigation on the Appendicular Skeleton of the Cattle Egret (Bubulcus Ibis)." *Journal of Experimental and Clinical Anatomy*. <https://doi.org/10.4103/1596-2393.158919>.
31. Saber, Ashraf, and B. Gummow. 2014. "Morphometric Studies on the Skull in Three Marsupial Species (Koala, Wombat, Wallaby)." *Journal of Veterinary Anatomy*. <https://doi.org/10.21608/jva.2014.44816>.
32. Sasan, Jasvinder Singh, Kamal Sarma, Shalini Suri, and Neelofar Nabi. 2019. "Gross and Morphometrical Studies on Humerus of Cattle Egret (Bubulcus Ibis)." *Exploratory Animal and Medical Research*.
33. Schneider, R. A., and J. A. Helms. 2003. "The Cellular and Molecular Origins of Beak Morphology." *Science*. <https://doi.org/10.1126/science.1077827>.
34. Sharma, Aarti, and S. C. Dubal. 2018. "Gross Anatomical Study of Pectoral Girdle of Small Indian Kite (Milvus Migrans Govinda)." *Veterinary Practitioner*.
35. Temeles, Ethan J., and W. John Kress. 2003. "Adaptation in a Plant-Hummingbird Association." *Science*. <https://doi.org/10.1126/science.1080003>.
36. Tiwari, Yogita, A Pandey, A B Shrivastav, M P S Tomar, and Rakhi Vaish. 2011. "Gross Morphometrical Studies on Pectoral Limb of Pariah Kite (Milvus Migrans)." *Research Article Annual Review & Research in Biology*. Vol. 1. [www.sciencedomain.org](http://www.sciencedomain.org).
37. Wehausen, J. D., and R. R. Ramey. 2000. "Cranial Morphometric and Evolutionary Relationships in the Northern Range of Ovis Canadensis." *Journal of Mammalogy*. <https://doi.org/10.1093/jmammal/81.1.145>.
38. Wu, Ping, Ting Xin Jiang, Jen Yee Shen, Randall Bruce Widelitz, and Cheng Ming Chuong. 2006. "Morphoregulation of Avian Beaks: Comparative Mapping of Growth Zone Activities and Morphological Evolution." *Developmental Dynamics*. <https://doi.org/10.1002/dvdy.20825>.
39. Wu, Ping, Ting Xin Jiang, Sanong Suksaweang, Randall Bruce Widelitz, and Cheng Ming Chuong. 2004. "Molecular Shaping of the Beak." *Science*. <https://doi.org/10.1126/science.1098109>.
40. Zusi, Richard L. 1993. "Patterns of Diversity in the Avian Skull." In *The Skull 2*.