

Full Length Research Paper

Evaluation of rice genotypes for brown planthopper (BPH) resistance using molecular markers and phenotypic methods

Sai Harini A.^{1*}, Sai Kumar S.¹, Padma Balaravi², Richa Sharma¹, Ayyappa Dass M.¹ and Vinay Shenoy¹

¹Barwale Foundation, Hyderabad, Andhra Pradesh, India.

²Department of Genetics, Osmania University, Hyderabad, Andhra Pradesh, India.

Accepted 11 April, 2013

Twenty eight (28) rice genotypes were used to evaluate the genetic variability based on known BPH resistant loci spread through most of the genome (chromosomes 2, 3, 4, 5, 6, 8, 10, 11 and 12), using closely linked simple sequence repeat (SSR) markers and by different phenotypic screening methods. A total number of 155 alleles were detected by 30 polymorphic markers with an average of 4.6 per locus. The genetic diversity, polymorphic information content (PIC) ranged from 0.15 to 0.89 and 0.13 to 0.88, respectively and the allele frequency ranged from 0.21 to 0.89. These microsatellite markers linked to BPH resistance loci classified rice genotypes into three clusters with additional sub groups and sub sub groups. Our study reveals high genetic variation and clear genotypic relationship for BPH resistance based on BPH resistance linked markers and known phenotypic screening methods such as standard seedbox screening technique, honey dew test and nymphal survival method. Phenotypic evaluation showed clear distinction between resistant and susceptible type by clearly revealing moderately resistant types as well. Combined use of phenotypic and genotypic evaluation methods can improve the efficiency of marker assisted selection and utilization of resistant genotypes for crop improvement by rice breeders.

Key words: *Nilaparvata lugens*, microsatellite markers, polymorphism, genetic diversity.

INTRODUCTION

Rice productivity is adversely impacted by numerous biotic and abiotic factors. An approximate 52% of the total global production of rice is lost annually owing to the damage caused by biotic factors, of which nearly 21% is attributed to the attack of insect pests (Brookes and Barfoot, 2003). Among the biotic stresses, the brown planthopper (BPH) *Nilaparvata lugens* (Stal.) is one of the most destructive monophagous insect pest, one of

the main biotic constraint of rice productivity causing huge yield losses every year in rice grown throughout tropical, subtropical and temperate areas in Asia (Park et al., 2008).

These insects draw nutrients from the phloem of rice plants. High BPH populations can destroy a plant in a short period of time (Huang, 2001, Yang et al., 2002). Large number of plant hoppers causes the infested plants

*Corresponding author. E-mail: saiharini@barwalefoundation.org.

Abbreviations: BPH, Brown planthopper; cM, Centimorgan; PIC, polymorphic information content; GD, genetic diversity; AF, allele frequency.

to become brown and dry. The condition is called hopperburn. Even if the planthopper population is not high enough to kill the plants, BPH feeding may considerably reduce yields (Watanabe et al., 1997). BPH can consume more than 28% of the total dry matter of rice plants infested at reproductive stage (Sogawa et al., 1994). BPH also transmits serious viral diseases, such as grassy stunt (Rivera et al., 1996) and ragged stunt virus (Ling et al., 1978). Development of resistant rice cultivars through host plant resistance is generally considered to be the most economic and effective way for controlling BPH damage. Molecular markers have demonstrated a potential to detect genetic diversity and relatedness of most crop species and to aid the management of plant genetic resources (Ford-Lloyd et al., 1997; Virk et al., 2000; Song et al., 2003; Ram et al., 2007). In contrast to morphological traits, molecular markers can reveal differences among genotypes at DNA level, providing a more direct, reliable and efficient tool for germplasm characterization, conservation and management. Among all DNA markers microsatellites (Wu and Tanksley, 1993; Yang et al., 1994) are codominant in nature; show high allelic diversity; are easily and economically assayed by PCR and can be automated. Many potential SSR markers have been identified in rice and over 25,000 have been developed as molecular markers (Temnykh et al., 2000; McCouch et al., 2002; IRGSP, 2005). These molecular markers have been effectively utilized for many purposes including genome mapping, assessment of the genetic diversity and relatedness among various cultivars and marker aided breeding (McCouch et al., 2001; Yu et al., 2003; Garris et al., 2005).

BPH populations on rice have been categorized in to four biotypes (Khush et al., 1985). The population in the east and Southeast Asia is reported as biotype 1, while biotype 2 originated in Indonesia and Vietnam as dominant biotype (Khush, 1979). Biotype 3 was produced in the laboratory at the International Rice Research Institute IRRI (Pathak and Khush, 1979) and in Japan (Ikeda and Vaughan, 1991) whereas biotype 4 is found only in South Asia. Till date, 26 BPH resistance genes have been identified in wild species *Oryza australiensis*, *Oryza officinalis*, *Oryza glaberrima*, *Oryza eichengiri*, *Oryza rufipogon*, *Oryza minuta* and Indian cultivars (Zhang, 2007; Fujita et al., 2008). The objective of the present effort was to evaluate the genetic variability among different rice genotypes based on known BPH resistant loci using closely linked simple sequence repeat (SSR) markers and by different phenotypic screening methods.

MATERIAL AND METHODS

Plant material

The experimental material consisted of 28 elite rice genotypes available at Barwale Foundation, viz; 1B, 2B, 7B, 8B, 9B, 14B, 16B, 18B, 21B, 22B, 24B, 25B, 28B, 30B, 36B, 40B, 41B, 44B, IR129, 1R150, IR157, IR168, Swarna, TN1, BPT5204, Pokkali, and PTB33,

where TN1 and PTB33 were used as susceptible and resistant check, respectively. The genotypes ranging from land race to improved lines, showed varied response to brown plant hopper stress.

Screening for BPH resistance

Initial population of BPH was collected from Maharajpet farm, Barwale Foundation and mass rearing of BPH and screening was done by following three methods, (i) standard seed box screening technique (SSST) developed at IRRI by Heinrichs et al. (1985), (ii) Honeydew test and (iii) Nymphal survival method.

Standard seed box screening technique

The experiment was conducted at a temperature of 28 to 30°C and relative humidity of 70 to 80%. The seeds were presoaked and sown in rows in 60 x 45 x 10 cm seed boxes along with resistant and susceptible checks. 25 to 30 seedlings per row were maintained per genotype. Ten (10) day old seedlings were infested with first instar nymphs at the rate of eight to 10 per seedling. Approximately one week after infestation 'hopperburn' symptom was observed. When more than 90% of susceptible check shows wilting, the plants were scored individually based on scoring system proposed by the International Rice Research Institute (IRRI, 1996) and each seedling was scored as 0 = no visible damage, 1 = partial yellowing of first leaf, 3 = first and second leaves partially yellowing, 5 = pronounced yellowing or some stunting, 7 = mostly wilted plant but still alive, 9 = the plant completely wilted or dead (Figure 1). Interpretation of results was based on standard evaluation system where the families with a mean rating of 0 to 3, 3.1 to 6.9 and 7 to 9 are designated as resistant, moderately resistant and susceptible, respectively (IRRI, 1996).

Honey dew test

The honeydew excretion is widely used to assess feeding activity and consequently a reliable index for resistance and susceptibility of a crop variety to homopteran pests (Auclair, 1959; Liu et al., 1994). Many techniques have been developed to measure the feeding response of *Nilaparvata lugens* on resistant and susceptible rice plants (Paguia et al., 1980; Pathak et al., 1982; Begum and Wilkins, 1998). The more important were the test of filter paper dipped in a solution of bromocresol green and the test of a parafilm sache. For our work, the filter paper technique was used. The plant at first tiller (40 days old) was located through two holes of the cup (up and down of the cup (5x5: HxR). The filter was placed at the base inside of the cup with a paper protecting it from humidity of the soil. For each plant to be screened, five female gravid hoppers were kept starving for 2 h 30 min. Then, the female hoppers were released on to plants to feed for 24 h, after which the filter papers were collected. Bromocresol green indicates phloem-based honeydew as blue-rimmed spots (indicate susceptible plants) and xylem-based honeydew as transparent (indicate resistant plants). The area of each spot on the bromocresol green-filter paper was measured using a digital scanner and "Image J" software (Figure 2).

Nymphal survival method

The nymphal survival test shows the difference of the survival of the nymphs on different varieties of rice plants. For this, 20 newly hatched nymphs in a pot with three rice plants (40 days old) were placed inside the mylar cages (45x5: HxR). The number of surviving nymphs was recorded every two days until they became adults (15 days). The experiment was carried in three replications and control plants were also maintained (Figure 3). Plants on which insects were released as well as control plants were cut till the base




Figure 1. Rice genotypes showing varied resistance levels for brown planthopper in standard seed box screening technique




Figure 2. Amount of honeydew excreted by brown planthopper on susceptible and resistant checks.



Figure 3. Experimental set up for screening rice genotypes by nymphal survival method.

of the stem and dried at 55°C for one week and biomass of infested plants and control plants were weighed. Number of insects surviving on individual genotype was counted.

DNA extraction, PCR amplification and PAGE

Total cellular DNA was isolated by rapid DNA extraction protocol

(Micklos and Freyer, 1990) and the quality of DNA was checked on agarose gel (0.8% w/v). Thirty four (34) SSR markers reported to be linked to 22 BPH resistant genes and quantitative trait locus (QTL) on chromosomes 2, 3, 4, 5, 6, 19, 11 and 12 were used for this study (Table 1). SSR primers were obtained from Sigma Aldrich, Bangalore. The PCR reaction was performed in 15 μ l volume using Eppendorf gradient thermocycler. The reaction mixture contained 10 ng/ μ l template DNA, 5 pM each forward and reverse primers, 10 mM each dNTPs, 2 μ l 10X PCR buffer (10 mM Tris HCl, 50 mM KCl, 0.01 mg/ml gelatin and 1.5 mM MgCl₂) and 0.5 μ l of 5 U/ μ l Taq polymerase. Thermal cycler was programmed to 1 cycle of 5 min at 94°C, 35 cycles of denaturation at 95°C for 1 min, 30 s for annealing temperature (55 to 60°C depending on the marker used) and 40 s at 72°C for primer elongation. Finally, 1 cycle of 7 min at 72°C was used for final extension. Amplified products were stored at -20°C until further use. The reproducibility of the amplification products was checked twice for each primer. SSR analysis was carried out by polyacrylamide gel electrophoresis following the protocol described by Paunaud et al. (1996) with some modifications.

Allele scoring and data analysis

All the genotypes were scored for the presence and absence of SSR bands and the data were entered in to a binary matrix as discrete variable, 1 for presence and 0 for the absence of the allele. Polymorphic information content (PIC) values were estimated using the formula: $PIC = 1 - \sum p_i^2$, where p_i is the frequency of the allele at each locus (Botstein et al., 1980). A Neighbor joining tree with bootstrap values was constructed with the help of Power marker Version 3.25 (Liu and Muse, 2005).

RESULTS

Phenotyping of rice genotypes for brown planthopper resistance

Standard seed box screening technique

The results of phenotypic response of rice genotypes to brown plant hopper screening at seedling stage (10 hoppers per seedling) indicated varied genotypic responses. Among all the 28 rice genotypes, PTB33 widely used as donor parent for BPH by rice breeders consisting of *Bph2* and *Bph3* genes (Khush, 1979) and Pokkali reported to have *Bph9* gene (Murata et al., 2001), scored as 1 and 3 respectively, and TN1 showed a score of 9 (Table 2).

Honey dew test

The amount of phloem and xylem in the honey dew excreted by the insect in the genotype was measured in mm² units (Figure 4). Among all the genotypes, TN1 has shown high rate of phloem consumption indicating that it is highly susceptible to brown planthopper/sucking pests; Whereas, insects on PTB33 showed least level of phloem consumption indicating its resistance. Genotypes like 1B, 2B, 9B, 21B, 22B, BPT5204, Swarna, showed high rate of phloem consumption. Genotypes like 7B, 36B, 41B, 44B, IR129, IR168, IR150, IR157, Pokkali and genotypes like 14B, 16B 18B, 28B, 24B, 25B, 40B, 29B, 30B and 8B

showed high to moderate levels of xylem consumption (Figure 2). The rate of phloem and xylem consumption by the genotypes was inversely proportional based on correlation of phloem consumption versus susceptibility and xylem consumption versus resistance.

Nymphal survival method

Numbers of nymphs surviving on PTB33 were less compared to other genotypes indicating its high level of resistance against brown planthopper, whereas large number of nymphs survived on TN1, indicating its susceptibility towards brown planthopper. Genotypes like 1B, 2B, 9B, 21B, 22B, BPT5204, Swarna, showed more number of insects and 7B, 36B, 41B, 44B, IR129, IR168, IR150, IR157, Pokkali and genotypes like 14B, 16B 18B, 28B, 24B, 25B, 40B, 29B, 30B and 8B showed less to moderate count of live insects (Figure 5), including number of nymphs surviving on each genotype, biomass of control plants and biomass of infested plants was recorded. All genotypes under control conditions were showing higher biomass compared to plants infested with insects. Among the infested plant, PTB33 was showing highest biomass while TN1 was showing lowest biomass after infestation among the genotypes (Figure 6).

Genotypic scoring

Thirty-four microsatellite or SSR markers reportedly linked with genes conferring BPH resistance were used to characterize and assess genetic diversity among 28 rice genotypes having varied response to brown plant hopper stress, ranging from parental lines, introgressed lines to donors. Out of the 34 markers used, four markers produced monomorphic bands while 30 markers showed polymorphism by revealing 155 alleles. The number of alleles per locus varied from 2 (RM459, RM496, RM261, RM6308, RM185) to 12 (RM8213) with an average of 4.6. The lowest amplicon size was produced by RM459 (75 bp) while highest amplicon size belonged to RM335 (315 bp). Many studies have also reported significant differences in allelic diversity among various microsatellite loci (Thompson et al., 2009). The PIC value measures the probability that two randomly chosen alleles from a population are distinguished. The PIC averaged 0.48 ranging from 0.13 (RM261) to 0.88 (RM8213). Markers (50%) were highly informative to slightly informative (pic > 0.5, informative markers). The polymorphic pattern of RM277, RM3331 and RM510 markers in 28 rice genotypes are presented in Figure 7. The gene diversity (GD) ranged from 0.15 to 0.89. The polymorphism Information Content (PIC) ranged from 0.13 to 0.88 and Allele Frequencies (AF) ranged from 0.21 to 0.89 (Table 3).

Genetic variation using SSR markers linked to BPH resistance genes

The microsatellite markers were able to distinguish

Table 1. Details of microsatellites used in the study.

| Marker | Chr | Gene/ QTL | cM | Start end | Clone | Reference(s) |
|---------|-----|-------------------|-------|------------|-----------|---|
| RM154 | 2 | Qbp3 | 1.05 | 1083810 | AP005851 | Ren et al. 2004 |
| RM3355 | 2 | Qbph2 | 95.2 | 25,728,952 | AYO20028 | Sun et al. 2007 |
| RM177 | 2 | Qbp3 | 22.35 | 22379610 | AL662937 | Ren et al. 2004 |
| RM410 | 3 | QTL | 17.6 | 17642689 | AP005676 | Sonada et al. 2003 |
| RM7 | 3 | Qbph3 | 64 | 38.6 | AF344009 | Sun et al. 2005 |
| RM6308 | 3 | Bph19(t) | 7.15 | 7160259 | AC137696 | Chen et al. 2006 |
| RM185 | 4 | Bph12(t) | 18.55 | 18564318 | AL662977 | Yang et al. 2002 |
| RM8213 | 4 | Bph17(t) | 4.4 | 4433168 | AL662959 | Sun et al. 2005 |
| RM335 | 4 | Bph12(t) | 21.5 | 679883 | AL606616 | Yang et al. 2002 |
| RM261 | 4 | Bph12(t),Bph15 | 6.55 | 6558954 | AL607008 | Yang et al. 2002 & 2004 |
| RM459 | 5 | QTL | 20.15 | 20155973 | AC130607 | Sonada et al. 2003 |
| RM589 | 6 | Bph3 | 1.35 | 1380856 | AP001168 | Jairipong Jairin et al. 2007 |
| RM510 | 6 | Qbph6 | 2.8 | 2831433 | AP006533 | Sun et al. 2007 |
| RM190 | 6 | Bph3 | 1.75 | 1764576 | AP002542 | Jairipong Jairin et al. 2007 |
| RM8072 | 6 | Bph3 | 1.4 | 1408326 | AP001168 | Jairipong Jairin et al. 2007 |
| RM217 | 6 | bph4 | 15 | 18.2 | AF344038 | Kawaguchi et al. 2001 |
| RM225 | 6 | bph4 BPH22(t) | 3.4 | 3416523 | AB023482 | Kawaguchi et al. 2001 Sai Harini et al. 2010 |
| RM588 | 6 | Bph3 | 1.6 | 1611388 | AP00391 | Jairipong Jairin et al. 2007 |
| RM19291 | 6 | Bph3 | 1.2 | 1215874 | AP003456 | Jairipong Jairin et al. 2007 |
| RM314 | 6 | Qbph6 | 33.6 | 5413067 | AF344139 | Sun et al. 2007 |
| RM496 | 10 | Qbph10 | 22.1 | 22171961 | AC087599 | Sun et al. 2005 |
| RM216 | 10 | Qbph10 | 5.5 | 5102292 | AC098566 | Jairipong Jairin et al. 2005 |
| RM209 | 11 | Bph10 | 17.75 | 17771745 | AC136150 | Jena et al. 2002 |
| RM3331 | 12 | QTL | 23.45 | 23460817 | AL845346 | Sonada et al. 2003 |
| RM277 | 12 | Qbph12 | 57.2 | 18100739 | AF344103 | Jairipong Jairin et al. 2005 |
| RM1986 | 12 | QTL | 21.2 | 21212965 | AL713932 | Sonada et al. 2003 |
| RM6869 | 12 | Bph18(t) | 22.2 | 22219502 | AL731741 | Jena et al. 2006 |
| RM463 | 12 | Bph18(t) | 22 | 22091957 | AL731742 | Jena et al. 2006 |
| RM50 | 6 | Qbph6 | 21 | 29500000 | AF343859 | Jairipong Jairin et al. 2005 |
| RM5953 | 4 | Qbph4 Bph20(t) | 9.35 | 9364068 | AP662989 | Sun et al. 2005 Rahman et al. 2009 |
| RM5479 | 12 | Bph21(t) | 57 | 24378959 | AL844880 | Rahman et al. 2009 |
| RM313 | 3 | Qbph3 | 20 | 38.8 | A02032481 | Sun et al, 2005 |
| RM484 | 10 | Qbph10 | 20.8 | 20808556 | AC073166 | Sun et al. 2005 |
| RM3134 | 3 | Bph19(t) | 7.25 | 7220514 | AC134229 | Chen et al. 2006 |

Table 2. Rice genotype ID, code and score for brown plant hopper resistance using standard seedbox screening technique.

| Genotype ID | Code | Score | Genotype ID | Code | Score |
|-------------|------|-------|----------------------|-------|-------|
| IR58025B | 1B | 8.3 | IR73793B | 30B | 6.9 |
| IR62829B | 2B | 7.3 | IR68886B | 36B | 5.8 |
| IR68888B | 7B | 5.5 | IR79156B | 40B | 5.6 |
| IR68892B | 8B | 6.1 | IR80151B | 41B | 5.8 |
| IR68897B | 9B | 7.9 | IR80156B | 44B | 5.6 |
| IR69628B | 14B | 6 | IR65482-7-216-1-2B | IR129 | 4.32 |
| IR70369B | 16B | 6.7 | IR73680-4-5-10-2-1-2 | IR150 | 4.3 |
| IR70959B | 18B | 6.6 | IR71033-121-15 | IR157 | 4.7 |
| IR72078B | 21B | 8.5 | IR73885-1-4-3-2-10 | IR168 | 5.8 |

Table 2. Contd.

| | | | | | |
|----------|-----|-----|---------------------|---------|---|
| IR72080B | 22B | 8 | MTU7029 | SWARNA | 9 |
| IR72018B | 24B | 6.6 | Thaichung native1 | TN1 | 9 |
| IR73320B | 25B | 6.8 | Samba Mahsuri | BPT5204 | 9 |
| IR73327B | 28B | 6.7 | Traditional variety | PTB 33 | 1 |
| IR73328B | 29B | 6.3 | Landrace | Pokkali | 3 |




Figure 4. Amount of phloem and xylem excreted in honeydew by brown planthopper feeding on different rice genotypes.




Figure 5. Number of nymphs surviving after 15 days on different rice genotypes.




Figure 6. Amount of biomass in control and infested plants of rice genotypes in nymphal survival method.




Figure 7. SSR banding patterns of 28 rice genotypes from RM277 (A), RM3331 (B), RM510(C).

between rice genotypes. The high degree of polymorphism of microsatellite allows rapid and efficient identification of rice genotypes. The microsatellite markers classified rice genotypes and the genetic relationships demonstrated among the genotypes in the neighbor joining tree (Figure 8) revealed some subgroups based on BPH resistant loci spread through most of the genome (chromosomes 2, 3, 4, 5, 6, 8, 10, 11, 12). The 28 genotypes

could be easily categorized into three clusters. Moving closer in terms of similarity, sub-divisions of such groups can reveal sub-sub groups. Cluster 1 included genotypes showing moderate resistance, cluster 2 is comprised of two sub clusters. Sub cluster 2.1 included all moderately resistant genotypes and sub cluster 2.2 comprised of two sub-sub groups A and B. Sub-sub group A included moderate resistance genotypes where as sub-sub cluster

Table 3. Total number of alleles, allele frequency, allele length in bp, genetic diversity, and polymorphism information content (PIC) value of 34 SSR markers assayed in 28 rice genotypes.

| Marker | Major AF | Number of alleles | Genetic diversity | PIC | AL in (bp) | AT (°C) |
|---------|----------|-------------------|-------------------|------|------------|---------|
| RM3331 | 0.48 | 6 | 0.70 | 0.69 | 120-150 | 55 |
| RM6308 | 0.78 | 3 | 0.36 | 0.35 | 120-125 | 55 |
| RM177 | 0.67 | 3 | 0.46 | 0.38 | 190-240 | 55 |
| RM261 | 0.98 | 2 | 0.15 | 0.13 | 150-160 | 55 |
| RM6869 | 0.51 | 5 | 0.63 | 0.59 | 120-140 | 55 |
| RM314 | 0.76 | 4 | 0.48 | 0.45 | 118-140 | 55 |
| RM209 | 0.53 | 4 | 0.56 | 0.36 | 145-165 | 55 |
| RM589 | 0.25 | 8 | 0.84 | 0.81 | 170-200 | 55 |
| RM19291 | 0.53 | 5 | 0.62 | 0.53 | 160-170 | 55 |
| RM185 | 0.69 | 3 | 0.44 | 0.42 | 190-205 | 55 |
| RM510 | 0.75 | 4 | 0.51 | 0.47 | 120-155 | 55 |
| RM154 | 0.3 | 8 | 0.83 | 0.81 | 170-230 | 55 |
| RM277 | 0.75 | 4 | 0.51 | 0.52 | 120-240 | 55 |
| RM588 | 0.46 | 4 | 0.64 | 0.63 | 125-145 | 55 |
| RM190 | 0.57 | 5 | 0.69 | 0.66 | 110-150 | 55 |
| RM3355 | 0.32 | 8 | 0.84 | 0.77 | 205-230 | 55 |
| RM410 | 0.25 | 8 | 0.87 | 0.86 | 180-295 | 55 |
| RM496 | 0.89 | 2 | 0.19 | 0.17 | 190-300 | 55 |
| RM217 | 0.73 | 4 | 0.52 | 0.47 | 130-145 | 55 |
| RM463 | 0.5 | 3 | 0.53 | 0.48 | 195-200 | 55 |
| RM1986 | 0.58 | 5 | 0.59 | 0.47 | 155-200 | 55 |
| RM216 | 0.32 | 7 | 0.77 | 0.77 | 145-165 | 55 |
| RM8213 | 0.21 | 12 | 0.89 | 0.88 | 155-205 | 58 |
| RM8072 | 0.82 | 3 | 0.30 | 0.30 | 120-160 | 55 |
| RM335 | 0.25 | 9 | 0.85 | 0.84 | 170-315 | 57 |
| RM459 | 0.82 | 4 | 0.31 | 0.29 | 75-150 | 55 |
| RM225 | 0.57 | 5 | 0.71 | 0.65 | 120-155 | 55 |
| RM7 | 0.58 | 4 | 0.59 | 0.40 | 120-125 | 55 |
| RM50 | 0.42 | 6 | 0.71 | 0.66 | 195-270 | 55 |
| RM5953 | 0.33 | 3 | 0.67 | 0.62 | 110-115 | 55 |
| RM5479 | 1.00 | 1 | 0.00 | 0.00 | 129-129 | 55 |
| RM3134 | 1.00 | 1 | 0.00 | 0.00 | 175-175 | 55 |
| RM313 | 1.00 | 1 | 0.00 | 0.00 | 111-111 | 55 |
| RM484 | 1.00 | 1 | 0.00 | 0.00 | 299-299 | 55 |
| Mean | 0.61 | 4.6 | 0.52 | 0.48 | | |

B comprised of resistance genotypes. All the susceptible genotypes were grouped in to cluster 3. Evidently, resistant genotypes (PTB33 and Pokkali) are undoubtedly distinct from moderately resistant genotypes (7B, 8B, 14B, 16B, 18B, 24B, 25B, 28B, 29B, 30B, 36B, 40B, 41B, 44B, IR129, IR150, IR157, IR168) and susceptible genotypes (1B, 2B, 9B, 21B, 22B, Swarna, BPT5204, TN1). The same association/relatedness is depicted by neighbor joining tree (Figure 8).

DISCUSSION

Among all the insect pests, brown planthopper, is one of the most destructive pests of rice causing severe yield

losses. Identification of donors for resistance and efficient screening techniques for evaluating breeding lines plays crucial role to transfer BPH resistance genes in to high yielding popular varieties and a high level of genetic diversity reduces the risk of wide spread epidemics of pests and diseases (Zhu et al. 2000; Newton et al. 2009).

The seedling resistance and antibiosis effects (honey dew test and nymphal survival method) of 28 rice genotypes on BPH were examined in this study. High resistance (Score 1) was shown by PTB33, followed by Pokkali (Score 3) and TN1 was found to be highly susceptible (score 9) using the three methods (SSST, honeydew test, and nymphal survival method) towards brown plant hopper infestation. Genotypes like 7B, 36B,




Figure 8. Cluster diagram based on similarity matrix calculated from 28 rice genotypes detected by 30 SSR markers.

41B, 44B, IR129, IR168, IR150, IR157 were showing upper grade of moderate resistance with score ranging between 3.1 to 5.8. Similarly, antibiosis methods revealed these genotypes to be moderately resistant towards brown plant hopper by consuming very low levels of phloem and high amounts of xylem as well as showing low amount of biomass consumption and low nymphal survival rate. Genotypes like 14B, 16B 18B, 28B, 24B, 25B, 40B, 29B, 30B and 8B were showing lower grade of moderate resistance with scores ranging between 6 to 6.9 and consuming phloem to some extent and later shifting to xylem consumption with moderate level of biomass consumption and nymphal survival. Rice genotypes like 1B, 2B, 9B, 21B, 22B, BPT5204, Swarna were highly susceptible (Score 7 to 9) feeding only on high amounts of phloem and high levels of biomass consumption and more number of nymphs.

Using 30 polymorphic SSR markers, total number of alleles (155) and the average number of alleles per locus (4.8) detected were higher than previously described for *Oryza sativa* genotypes (Vijaya Lakshmi et al., 2010; Narshimulu et al., 2010). Using 37 SSR markers, Vijaya Lakshmi et al. (2010) found 88 alleles with an average of 2.37, while Narshimulu et al. (2010) reported 96 alleles with an average of alleles per locus 2.67. The PIC values ranged from 0.13 to 0.88 which are similar (0.12 to 0.83) to those reported for genetic diversity assessment in rice genotypes (Pervaiz et al., 2010). Using high informative markers, indicated by their high PIC values maximized the probability of detecting high proportion of allelic

variation at the individual loci. The PIC values in this study were comparable to those reported in some studies (Jain et al., 2004; Saini et al., 2004; Siwach et al., 2004; Lu et al., 2005; Jayamani et al., 2007; Thomson et al., 2007) but higher than those reported by Singh et al. (2004) and Joshi and Behera (2006). It is to be noted here that other similar studies involve whole-genome markers, while in our study, high PIC values are obtained by BPH resistance loci-specific markers, otherwise expected to reveal limited polymorphism. The alleles revealed by markers showed a high degree of polymorphism, with all markers used (excluding 2, RM496 and RM459) producing 100% polymorphic bands. This amply suggests that the selected set of SSRs used in this study were extremely informative and effective for assessing the genetic diversity at specific BPH resistance loci, where a reduced level of genetic variability is generally expected.

Genetic diversity of rice genotypes has been studied based on SSR markers linked to BPH resistance genes. The tree showed a close relationship between resistance and moderately resistant genotypes based on BPH resistance loci linked marker analysis. However, phenotyping methods could clearly differentiate rice genotypes categorized as resistant and moderately resistant genotypes. The same relatedness between resistant (R) and moderately resistant (MR) types is clearly depicted by tree as well, wherein susceptible (S) types have branched separately. Cluster analysis of the 28 rice genotypes based SSR marker data divided genotypes in to three clusters

(Figure 8) with additional sub groups within each group. The parental phenotypic evaluation is in coincidence with genotypic cultivars using SSR markers.

Conclusively, evaluation of germplasm for resistance genes plays a major role in selection of parental lines and development of new breeding material. Information on target loci obtained from markers will facilitate use of germplasm efficiently. In addition to evaluation of genetic variation at specific loci associated with BPH resistance, the genotypes were taken up for phenotypic screening for BPH resistance and the score was correlated with genotype groups for resistance reaction. Our study revealed high genetic variation and clear genotypic relationship in NJ tree generated by Power Marker. The results further indicate that since the SSR markers are neutral and co-dominant, they are powerful tools to access the genetic variability of the cultivars under study. Use of marker and phenotype information together offers an efficient tool to the breeders in selecting parents for various breeding programs *viz.* mapping and marker assisted selection (MAS).

ACKNOWLEDGMENTS

The authors convey their gratitude towards Mr. Dinesh C. Joshi, Executive Director, Barwale Foundation, for valuable support and funding to carry out the work.

REFERENCES

- Auclair JL (1959). Feeding and excretion by the pea aphid *Acyrtosiphon pisum* (Harr.) (Homoptera: Aphididae) reared on different varieties of peas. *Entomologia Experimentalis et Applicata*. 2:279-286
- Begum MN, Wilkins RM (1998). A parafilm sachet technique for measuring the feeding of *Nilaparvata lugens* Stal. on rice plants with correction for evapotranspiration. *Entomologia Experimentalis et Applicata*. 88:301-304
- Botstein D, White RL, Skolnick M, Davis RW (1980). Construction of a genetic linkage map in man using restriction fragment length polymorphisms. *Am. J. Hum. Genet.* 32:314-331
- Brookes G, Barfoot P (2003). GM Rice: Will this Lead the Way for Global Acceptance of GM Crop Technology? Brief No. 28 Los Baños, Philippines, ISAAA
- Chen JW, Wang L, Pang XF, Pan H (2006). Genetic analysis and fine mapping of a rice brown planthopper (*Nilaparvata lugens* Stål) resistance gene *bph19(t)*. *Mol. Gen. Genomics* 275:321-329.
- Ford-Lloyd BV, Jackson MT, Newbury HJ (1997). Molecular markers and the management of genetic resources in seed gene banks: a case study of rice. In: Callow JA, Ford-Lloyd BV, Newbury HJ, eds. *Biotechnology and plant genetic resources: conservation and use*. Wallingford, CAB International. pp. 103-118
- Fujita D, Myint KKM, Matsumura M, Yasui H (2008). The genetics of host plant resistance to rice planthopper and leafhopper. Presented in International Conference on rice plant hopper, IRRI, Philippines, 23-25 June 2008.
- Garris J, Thomas HT, Coburn J, Kresovich S, McCouch S (2005). Genetic Structure and Diversity in *Oryza sativa* L. *Genetics Society of America* 169 (3):1631-1638
- Heinrichs EA, Medrano FG, Rapusas HR (1985). Genetic evaluation for insect resistance in rice. IRRI, Los Baños, Philippines. pp. 1-356.
- Huang Z, Shu L, Li X, Zhang Q (2001). Identification and mapping of two brown planthopper resistance genes in rice. *Theor. Appl. Genet.* 102:929-93
- IRGSP (2005). The map based sequence of rice genome. *Nature* 436:793-800
- IRRI (1996). Standard Evaluation System for Rice. IRRI, Manila, Philippines.
- Jain S, Jain RK, McCouch SR (2004). Genetic analysis of Indian aromatic and quality rice (*Oryza sativa* L.) germplasm using panels of fluorescently-labeled microsatellite markers. *Theor Appl Genet.* 109(5): 965-977
- Jairin J, Phengrat K, Teangdeerith S, Vanavichit A, Toojinda T (2007). Mapping of a broad spectrum brown planthopper resistance gene, *Bph3*, on rice chromosome 6. *Mol. Breed.* 19:35-44.
- Jairin J, Toojinda T, Tragoonrung S, Tayapat S, Vanavichit A (2005). Multiple Genes Determining Brown Planthopper (*Nilaparvata lugens* Stal) Resistance in Backcross Introgressed Lines of Thai Jasmine Rice 'KDML105'. *Science Asia* 31: 129-135
- Jayamani P, Negrao S, Martins M, Macas B. and Oliveira MM (2007). Genetic relatedness of Portuguese rice accessions from diverse origins as assessed by microsatellite markers. *Crop Sci.* 47(2):879-886.
- Jena KK, Jeung JU, Lee JH, Choi HC, Brar DS (2006). Planthopper (BPH) resistance gene, *Bph18(t)*, and marker-assisted selection for BPH resistance in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 112:288-297
- Jena KK, Pasalu IC, Rao YK, Varalaxmi Y, Krishnaiah K, Khush GS, Kochert G (2002). Molecular tagging of a gene for resistance to brown planthopper in rice (*Oryza sativa* L.). *Euphytica* 129:81-88.
- Joshi RK, Behera L (2006). Identification and differentiation of indigenous non-Basmati aromatic rice genotypes of India using microsatellite markers. *Afr. J. Biotechnol.* 6 (4): 348-354
- Kawaguchi M, Murata K, Ishii T, Takumi S, Mori N, Nakamura C (2001). Assignment of a brown planthopper (*Nilaparvata lugens* Stål) resistance gene *bph4* to the rice chromosome 6. *Breed. Sci.* 51:13-18.
- Khush GS (1979) Genetics and breeding for resistance to brown planthopper. In: IRRI (ed) *Brown planthopper: threat to rice production in Asia*. IRRI, Los Baños, Philippines. pp. 321-322
- Khush GS, Karim AR, Angeles EG (1985). Genetics of resistance of rice cultivar ARC10550 to Bangladesh brown planthopper biotype. *J. Genet.* 64(2):121-125
- Ling, KC, Tiongco ER, Aquiero VM (1978). Rice ragged stunt, a new virus disease. *Plant Dis. Rep.* 62: 701-705
- Liu G, Saxena RC, Wilkins RM (1994). Behavioral responses of the whitebacked planthopper, *Sogatella furcifera* (Homoptera: Delphacidae) on rice plants whose odors have been masked. *J. Insect Behav.* 7: 343-353.
- Liu K, Muse SV (2005). Power Marker: integrated analysis environment for genetic marker data. *Bioinformatics* 21: 2128-2129
- Lu Hong, Redus, Mare A, Coburn, Jason Rutger, Neil J, McCouch Susan R, Tai, Thomas H (2005). Population structure and breeding patterns of 145 US rice cultivars based on SSR marker analysis. *Crop Sci.* 45(1): 66-76.
- McCouch S, Teytelman L, Y Xu, Lobos K, Clare K, Walton M, Fu B, Maghirang R, Li Z, Xing Y, Zhang Q, Kono I, Yano M, Fjellstrom R, DeClerck G, Schneider D, Cartinhour S, Ware D, Lincoln S (2002). Development and mapping of 2240 new SSR markers for rice (*Oryza sativa* L.). *DNA Research* 9: 199-207
- McCouch SR, Temnykh S, Lukashova A, Coburn J, Declerck G, Cartinhour S (2001). Microsatellite markers in rice: Abundance, diversity and applications. In: *Rice genetics IV*. IRRI, Manila, Philippines, p 117-135.
- Micklos DA, Freyer GA (1990). *DNA Science: A First Course in Recombinant DNA Technology*. Carolina Biological Supply Company and Cold Spring Harbour Laboratory Press, North Carolina, USA. pp. 266.
- Murata K, Fujiwara M, Murai H, Takumi S, Mori N, Nakamura C (2001). Mapping of a brown planthopper (*Nilaparvata lugens* Stål) resistance gene *Bph9* on the long arm of chromosome 12. *Cereal Res. Commun.* 29:245-250
- Narasimhulu G, Jamloddin M, Vemireddy LR, Anuradha G, Siddiq EA (2010). Potentiality of evenly distributed hypervariable microsatellite markers in marker assisted breeding of rice. *Plant Breed.* 130(3): 314-320

- Newton AC, Begg GS, Swanston JS (2009). Deployment of diversity for enhanced crop function. *Ann. Appl. Biol.* 154:309-322
- Paguia P, Pathak MD, Heinrichs EA (1980). Honeydew Excretion Measurement Techniques for Determining Differential Feeding Activity of Biotype of *Nilaparvata lugens* on Rice Varieties. *J. Econ. Entomol.* 73:35-40.
- Park DS, Song MY, Park SK, Lee SK, Lee JH (2008). Molecular tagging of the Bph locus for resistance to brown planthopper (*Nilaparvata lugens* Stal) through representational divergence analysis. *Mol. Genet. Genomics* 280:163-172.
- Pathak MD, Khush GS (1979). Studies of varietal resistance in rice to the brown planthopper at the International Rice Research Institute. In: IRRI (ed) *Brown planthopper: threat to rice production in Asia*. IRRI, Los Banos, Philippines. pp 285-301.
- Pathak PK, Saxena RC, Heinrichs EA (1982). Parafilm Sachet for Measuring Honeydew Excretion by *Nilaparvata lugens* on Rice. *J. Econ. Entomol.* 75: 194-195
- Rahman ML, Jiang W, Chu SH, Qiao Y, Ham TH, Woo MK, Lee J, Khanam MS, Chin JH, Jeung JU, Brar DS, Jena KK, Koh HJ (2009). High-resolution mapping of two brown planthopper resistance genes, *Bph20(t)* and *Bph21(t)*, originating from *Oryza minuta*. *Theor. Appl. Genet.* 119:1237-1246
- Ram, Venkatesan Thiruvengadam, Kunnummal Krungari Vinod (2007). Genetic Diversity among cultivars, landraces and wild relatives of rice as revealed by microsatellite markers. *J. Appl. Genet.* 48(4):337-345.
- Ren X, Wang X, Yuan H, Weng Q, Zhu L, He G (2004). Mapping quantitative trait loci and expressed sequence tags related to brown planthopper resistance in rice. *Plant Breed.* 123:342-348
- Rivera CT, Ou SH, and Lida TT (1966). Grassy stunt disease of rice and its transmission by *Nilaparvata lugens* (Stal). *Plant Dis. Rep.* 50: 453-456.
- Sai Harini A, Santhana lakshmi S, Sai kumar S, SivaramaKrishnan S, Kadirvel P (2010). Validation and fine-mapping of genetic locus associated with resistance to Brown planthopper, *Nilaparvata lugens* (Stal.) in rice, *Oryza sativa* L. *Asian J. Biosci.* 5: 32-37.
- Saini N, Jain Neelu, Jain S, Jain RK (2004). Assessment of genetic diversity within and among Basmati and non-Basmati rice varieties using AFLP, ISSR and SSR markers. *Euphytica.* 140 (3):133-146
- Singh RK, Sharma RK, Singh AK, Singh VP, Singh NK, Tiwari SP, Mohopatra T (2004). Suitability of mapped sequence tagged microsatellite site markers for establishing distinctness, uniformity and stability in aromatic rice. *Euphytica.* 135(2):135-143
- Siwach P, Jain S, Saini N, Chowdhury VK, Jain RK (2004). Allelic diversity among Basmati and non-Basmati long-grain indica rice varieties using microsatellite markers. *J. Plant Biochem. Biotechnol.* 13(1) 25-32.
- Sogawa K, Gallagher KD, Kenmore PPE (1994). Pest management in tropical paddy ecosystems. *Japanese J. Trop. Agric.* 38:361-368. (in Japanese with English abstract)
- Sonada TA, Yoshimura, Yasui H (2003). Detection of QTLs for antibiosis to brown plant hopper. *Rice Genet. Newsl.* 20:83-85
- Song ZP, Xu X, Wang B, Chen JK, Lu BR (2003). Genetic diversity in the northernmost *Oryza rufipogon* populations estimated by SSR markers. *Theor Appl Genet.* 107: 1492-1499
- Sun L, Liu Y, Jiang L, Su C, Wang G, Zhai H, Wan J (2007). Identification of quantitative trait loci associated with resistance to brown planthopper in the indica rice cultivar Col.5 Thailand. *Hereditas* 144(2):48-52
- Sun L, Su C, Wang C, Zai H, Wan J (2005). Mapping of a major resistance gene to brown planthopper in the rice cultivar Rathu Heenati. *Breed Sci.* 55:391-396
- Temnykh S, Park WD, Ayres N, Cartinhour S, Hauck N, Lipovtiesich L (2000). Mapping and genome organization of microsatellite sequences in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 100:697-712.
- Thomson MJ, Septiningsih EM, Suwardjo Fatimah, Santoso Tri J, Silitonga, Tiur S, McCouch SR (2007). Genetic diversity analysis of traditional and improved Indonesian rice (*Oryza sativa* L.) germplasm using microsatellite markers. *Theor. Appl. Genet.* 114(3): 559-568
- Vijaya Lakshmi B, Srinivas B, Vinaykumar M, Vijay D (2010). Assessment of Rice genotypes For Brown Planthopper using Microsatellite Markers. *Curr. Trends Biotechnol. Pharm.* 4: 526-534
- Virk PS, Newbury JH, Bryan GJ, Jackson MT, Ford-Lloyd BV (2000). Are mapped or anonymous markers more useful for assessing genetic diversity? *Theor. Appl. Genet.* 100: 607-613.
- Watanabe T, Fabellar LT, Almazan LP, Rubia EG, Heong KL, Sogawa K (1997). Quantitative evaluation of growth and yield of rice plants infested with rice planthoppers. In M.J. Kropff et al. [eds.] *Applications of Systems Approaches at the Field Level*. Kluwer Academic Publishers, Dordrecht. pp. 365 - 382.
- Wu KS, Tanksley SD (1993). Genetic and physical mapping of telomeres and microsatellites of rice. *Plant Mol. Biol.* 22: 861-872.
- Yang GP, Maroof MAS, Xu CG, Zhang Q, Biyashev RM (1994). comparative analysis of microsatellite DNA polymorphism in landraces and cultivars of rice. *Mol. Gen. Genet.* 245: 187-194
- Yang H, You A, Yang Z, Zhang F, He R, Zu L, He G (2004). High resolution genetic mapping at the *Bph15* locus for brown planthopper resistance in rice (*Oryza sativa* L.). *Theor. Appl. Genet.* 110:182-191
- Yang HY, Ren X, Weng QM, Zhu LL, He GC (2002). Molecular mapping and genetic analysis of a rice brown planthopper (*Nilaparvata lugens* Stål) resistance gene. *Hereditas* 136:39-43.
- Yu SB, Xu WJ, Vijaykumar CHM, Ali J, Fu BY, Xu JL (2003). Molecular diversity and multilocus organization of the parental lines used in the International Rice Molecular Breeding program. *Theor. Appl. Genet.* 108:131-40
- Pervaiz ZH, Rabbani AM, Ishtiaq Khaliq, Pearce SR, Malik SA (2010). Genetic diversity associated with agronomic traits using microsatellite markers in Pakistani rice landraces. *Electronic J. Biotechnol.* 13 (3). doi: 10.2225
- Zhang Q (2007). Strategies for developing Green Super Rice. *Proc. Natl. Acad. Sci. USA* 104:16402-16409.
- Zhu Y, H.Chen, J Fan, Wang Y, Li Y (2000). Genetic diversity and disease control in rice. *Nature* 406: 718-72