

Short Communication

Antimicrobial activity of *Mucuna pruriens* on selected bacteria

Salau, A. O.* and Odeleye, O. M.

¹Department of Pharmaceutics, Faculty of Pharmacy, Obafemi Awolowo University Ife, Osun State, Nigeria.

²Department of Pharmacognosy, Faculty of Pharmacy, Obafemi Awolowo University Ife, Osun State, Nigeria.

Accepted 20 February, 2007

The investigation is aimed to carry out the antimicrobial activities of the methanol extract of *Mucuna pruriens* leaves. The antimicrobial activity of the extract was determined by the agar well diffusion method against various gram positive, gram negative and spore forming microorganisms and fungi. The methanol extract of *M. pruriens* showed broad-spectrum antimicrobial activity against all the tested microorganisms except *Candida albicans*. The results obtained in the study shows that *M. pruriens* extract can be a potential source of natural antimicrobial agent.

Key words: Agar diffusion, antimicrobial activity, *Mucuna pruriens*.

INTRODUCTION

Plant based antimicrobials represent a vast untapped source for medicines and further exploration of plant antimicrobials needs to occur. Antimicrobials of plant origin have enormous therapeutic potential. They are effective in the treatment of infectious diseases while simultaneously mitigating many of the side effects that are often associated with synthetic antimicrobials (Iwu et al., 1999).

Mucuna pruriens belongs to the family Fabaceae. Some of the common names include: Cow itch, Common Cow-itch, Konch, velvet bean, mucuna, nescafé, pó de mico, fava- coceira, cabeça-de-frade, cowage, cowhage, bengal bean, Mauritius bean, itchy bean, krame, picapica, chiporro, and buffalo bean. The roots are bitter, sweet thermogenic emollient, stimulant, purgative, aphrodisiac and diuretic. The leaves are aphrodisiac. The seeds are astringent, laxative, anthelmintic, alexipharmic and tonic (Taylor, 2005). A clinical study confirmed the efficacy of the seeds in the management of Parkinson's disease by virtue of their L-DOPA content (Manyam et al., 1995; Bell et al., 1971). *M. pruriens* has been shown to increase testosterone levels (Amin et al., 1996), leading to deposition of protein in the muscles and increased muscle mass and strength (Bhasin et al., 1996).

Much work has not been done on the leaf, hence the main reason of this investigation.

M. pruriens itch-producing property is attributed to the trichomes (hair) present on the pods. It has been established that this unique property is accounted by the presence of 5-hydroxy tryptamine (5-HT) in the hair (Armstrong et al., 1953). Some reports show that anti-histaminics afford protection against the itch (Broadbent, 1953). In this work, we tested the antimicrobial activity (against gram positive, gram negative and spore forming bacteria and also fungi) of the methanol extract of the leaf of *M. pruriens*

MATERIALS AND METHOD

The leaf of *M. pruriens* was collected from bushes around the Obafemi Awolowo University Ile-Ife Osun State, Nigeria. The leaf was identified in the herbarium of the Faculty of Pharmacy of the Obafemi Awolowo University Ile-Ife Osun State, Nigeria. The leaf was extracted using methanol.

Microorganisms (*Staphylococcus aureus* NCTC 6571, *Escherichia coli* NCTC 10418, *Bacillus subtilis* NCTC 8263, *Pseudomonas aeruginosa* ATCC 10145) were obtained from the stock culture of Department of Pharmaceutics, Obafemi Awolowo University Ile-Ife Osun State, Nigeria.

Antimicrobial activity was determined by agar-well diffusion method (Kavanagh, 1963) and minimum inhibitory concentrations were determined (Russell and Furr, 1977).

*Corresponding author. E-mail: dtwoabiola@yahoo.com. Tel: +2348034725754.

Table 1. Antimicrobial activity of *Mucuna pruriens* leaf extract

Zones of inhibition (mm) ³								
Microorganism	<i>Mucuna pruriens</i> (mg/ml)						MIC (mg/m)	Streptomycin (1 mg/ml)
	240	160	80	40	20	5		
<i>Staph. aureus</i> NCTC	45	15	12	0	0	0	80	21
<i>E. coli</i> NCTC 10418	28	13	0	0	0	0	160	24
<i>B. subtilis</i> NCTC 8263	38	14	12	11	0	0	40	26
<i>P.aeruginosa</i> ATCC 10145	30	15	11	0	0	0	80	0
<i>Proteus mirabilis</i> NCIB 67	34	18	17	14	0	0	40	22
<i>Candida albicans</i>	0	0	0	0	0	0	-	0

RESULTS AND DISCUSSION

Table 1 above indicates that the extract inhibited the growth of all the tested microorganisms with the exception of *Candida albicans* to various degrees. The extract showed strong antibacterial activity against *S. aureus* NCTC 6571, *E. coli* NCTC 10418, *B. subtilis* NCTC 8263, *Proteus mirabilis* NCIB 67, *P. aeruginosa* ATCC 10145 (which is most times resistant to most antimicrobial agents). Their activity at 240 mg/ml is comparable to that of the control used (streptomycin at a concentration of mg/ml). *Candida albicans* was not sensitive to the extract.

On the basis of the result obtained in this present investigation, we conclude that the methanol extract of *M. pruriens* leaves had significant *in vitro* antimicrobial activity. The obtained results may provide a support to some uses of the plant in traditional medicine. Further studies are recommended to isolate the active components responsible for the antimicrobial activity.

REFERENCES

- Amin KMY, Khan MN, Zillur-Rehman S (1996). Sexual function improving effect of *Mucuna pruriens* in sexually normal male rats. J. Study Med. Plant. Fitoterapia, 67(nr.1): 53-58.
- Armstrong D, Arcy RML, Keela CA, Maikhana M (1953). Observations on chemical excitants of cutaneous pain in man. J. Physiol. 120: 326-351.
- Bell EA, Nulu JR, Cone C (1971). Swaminathan Research Foundation of Chennai. Phytochemistry. 10: 2191-2194.
- Bhasin S, Storer TW, Berman N, Callegari C, Clevenger B, Phillips J, Bunnell TJ, Tricker R, Shirazi A, Casaburi R (1996). The Effects of Supraphysiologic Doses of Testosterone on Muscle Size and Strength in Normal Men. N. Engl. J. Med., 335: 1-7.

Broadbent JL (1953). Observation on itching produced by Cowhage and on the part played by histamine as a mediator of the itch sensation Brit. J. Pharmacol. Chemother. 8: 263-270.

Iwu MW, Duncan AR, Okunji CO (1999). New antimicrobials of plant origin. p. 457-462. In: J. Janick (ed.), Perspectives on new crops and new uses. ASHS Press, Alexandria, VA.

Kavanagh F (1963). Analytical microbiology. New York: Academic Press. p. 290.

Manyam BV, Dhanasekaran M, Hare TA (1995). An Alternative Medicine Treatment for Parkinson's disease: Results of a Multicenter Clinical Trial, J. Altern. Complement Med., 1(3): 249-255.

Russell AD, Furr JR (1977). Antibacterial activity of a new chloroxylenol preparation containing ethylenediamine tetra acetic acid. J. Appl. Bacteriol. 243: 253.

Taylor L (2005). The Healing Power of Rainforest Herbs, p. 444.