

Full Length Research Paper

Construction and analysis of a cDNA library from yellow-fruit ginseng (*Panax ginseng* C.A.Meyer.) leaf tissue

Chengjun Yang^{1,2}, Jun Wang^{2,3*}, Chuanping Yang² and Guifeng Liu²

¹Postdoctoral Research Centres of Biology, Northeast Forestry University, Harbin 150040, P. R. China.

²Key Laboratory of Forest Tree Genetic Improvement and Biotechnology, Ministry of Education, College of forestry, Northeast Forestry University, Harbin 150040, P. R. China.

³College of Food Science and Nutritional Engineering, China Agricultural University, Beijing 100083, P. R. China.

Accepted 9 July, 2009

The total RNA was isolated from yellow-fruit ginseng (*Panax ginseng* C.A. Meyer) leaf tissue. A cDNA library of panax ginseng leaves was constructed by using pDNR-LIB vector according to the SMART cDNA library construction kit protocol. We obtained 378 high quality sequences (GenBank accession number: ES672876-ES673253). ESTs were annotated, analyzed by BlastX and functional classified based on gene ontology, the results showed that 221 ESTs showed significant similarities to gene sequences in Nr database and were known genes, 21 ESTs were non-significance and unknown function genes, and 136 ESTs were considered novel genes. Most of the ESTs appeared to be related to physiological and cellular processes.

Key words: cDNA library, expressed sequence tags (EST), *Panax ginseng*.

INTRODUCTION

Ginseng (*Panax ginseng* C. A. Meyer), a perennial herb from the Araliaceae family, is one of the most commonly utilized medicinal plants. Ginseng is considered to be one of the most potent medicinal plants that have been used to bolster immunity, provide nutrition, ameliorate fatigue and enhance resistance to stress, disease and exhaustion. Ginsenosides, which are triterpene glycosides (saponins), are believed to be the main active compounds in ginseng tissues (Choi et al., 2005; Kim et al., 2006). More than 30 different ginsenosides have been isolated from ginseng plants (Sun et al., 2001; Zheng et al., 2001). Despite the considerable commercial interest in ginsenosides, little is known about the genes and biochemical pathways of ginsenoside biosynthesis. 'Jilin yellow-fruit ginseng' is identified as a homozygous recessive mutant from ordinary panax ginseng and contains high levels of ginsenosides, rich proteins, amino acids

and other nutrients (Zhao et al., 1998).

cDNA library is one of the basic means to study functional genomics. Expressed sequence tags (ESTs) are partial sequences of randomly selected complementary DNA (cDNA) clones; automated sequencing techniques make it possible to generate large numbers of EST at one time. Expressed sequence tags (EST) analysis is an effective method to discover novel genes and investigate gene expression in different organs and tissues (Wang et al., 2006). The generation and analysis of expressed sequence tags provides useful information on development, metabolism and signaling in various organisms. Expressed sequence tags have applications in the discovery of new genes, mapping of the genome and identification of coding regions in genomic sequences (Miyahara et al., 2000).

In the study, we constructed a cDNA library of yellow-fruit ginseng leaf tissue and obtained 378 high quality EST sequences. Through EST analysis and gene ontology, this will help us to understand the gene expression pattern, discover novel genes and study the biochemical pathways of ginsenoside biosynthesis.

*Corresponding author. E-mail: junwang1966@yahoo.com.cn.
Tel: 86-045-82191829(O). Fax: 86-451-82190607-11.



Figure 1. Agarose gel electrophoresis of total RNA from yellow-fruit ginseng leaf.




Figure 2. Agarose gel electrophoresis of double-stranded cDNA. Lane 1: double-strand cDNA; lane M: DL2000 marker.

MATERIALS AND METHODS

Plant materials

Actively growing 2 year old yellow-fruit ginseng (*Panax ginseng* C. A. Meyer) leaves were obtained from ZuoJia, Jilin province, China, on June 20, 2006. The harvested leaves were immediately frozen in liquid nitrogen and then stored at -70°C until RNA isolation.

Isolation and quantification of RNA

The total RNA was isolated from ginseng leaf tissue using SDS methods, as described by Yang et al. (2008). Total RNA was quantified by measuring the optical density of a dilute RNA solution. The integrity of the RNA was analyzed using 1.1% agarose/EtBr gel electrophoresis. The purity of the RNA was checked by the ratio of $\text{OD}_{260}/\text{OD}_{280}$.

cDNA synthesis and library construction

In accordance with the creator^{im} SMARTTM cDNA library construction kit user manual provided by the manufacturers (Clontech), total RNA (1.0 μg) as starting material was reverse transcribed to synthesize first-strand cDNA, and double-strand cDNA was synthesis by LD-PCR. 5 μg of the double-stranded cDNA were taken for analysis by electrophoresis on a 1.1% agarose/EtBr gel. Then the amplified double-strand cDNA was digested with proteinase K and Sfi I. After digestion, cDNA size fractionation was performed using chroma spin-400 columns to collect cDNA large than 400 bp and checked the profile of fractions on a 1.1% agarose/EtBr gel. The cDNA was ligated to the Sfi I-digested dephosphorylated pDNR-LIB vector provided with the kit and electroporated into DH5 α *Escherichia coli* bacteria to develop the cDNA library of ginseng leaf tissue. To make a large, stable quantity of a high-titer stock of the library, we amplified the primary cDNA library.

Identification of the cDNA library

According to the library titrating protocol, the unamplified and amplified cDNA library were tittered. To identify the cDNA inserts of

the recombinants and determined the percentage of recombinant clones, 16 plaques were randomly picked from plate. Then PCR was performed with M13 primers provided by the advantage 2 PCR kit. The PCR products checked on 1.2% TAE/agarose gel with DNA size markers.

Library sequencing and analysis

The cDNA library clones were plated into LB agar plate containing 30 $\mu\text{g}/\text{ml}$ of chloramphenicol, white clones were picked in to 96 well plates randomly, plasmids DNA of each clone were prepared by standard alkaline lysis preparation protocol and then sequenced by Beijing genomics institute. Sequencing was performed in a Mega BACE1000 DNA capillary sequence machine. The raw expressed sequence tags (EST) were edited to remove vector and poor quality sequences. The remaining sequences were subjected to blast analysis against the non-redundant database on the GeneBank (<http://www.ncbi.nlm.nih.gov/blast>) for similarity. The confirmed sequences were submitted to the dbEST database of GeneBank. Identified genes were classified according to gene ontology. The network information of the gene ontology database is categorized into 3 groups: cellular component, molecular function and biological process.

RESULTS AND DISCUSSION

High-quality total RNA was isolated from the leaf tissue of ginseng. Electrophoresis of the total RNA on 1.1% agarose/EtBr showed distinct 28S and 18S rRNA bands and the ratio of intensities of 28S and 18S was about 2:1 (Figure 1), the ratio of $\text{OD}_{260}/\text{OD}_{280}$ to the total RNA was 1.90. The total RNA isolated was integrated and suitable for constructing the cDNA library. The double-strand cDNA synthesized using LD-PCR was analyzed on a 1.1% agarose/EtBr gel and the product showed a smear from 0.2 to 2.5 kb (Figure 2). The double-strand cDNA fractionated using CHROMA SPIN-400 columns was




Figure 3. Double-stranded cDNA fractionated on a 1.1% agarose/EtBr gel 1-18: double-stranded cDNA; M: DL2000 marker.




Figure 4. 16 clones in the cDNA library were selected randomly to evaluate their insert sizes. Lanes 1-16: Products of inserts; lane M: DL2000 marker.

collected and pipette 3 μ l of cDNA to run on a 1.1% agarose/EtBr gel at 150 V for 10 min. Electrophoresis results showed the cDNA in lanes 7 - 11 was larger than 400 bp (Figure 3) and was ready to be ligated to the Sfi I-digested, dephosphorylated pDNR-LIB vector.

The titer of the un-amplified constructed cDNA library was approximately 1.01×10^6 pfu/ml, the titer of the amplified cDNA library was 2.16×10^9 pfu/ml. 16 independent clones were selected randomly from unamplified cDNA library and were amplified by PCR to check the percentage of recombinants and the insert size of the recombinants. The results showed the percentage of recombinants was 93.75%, the cDNA inserts ranged from 0.3 to 2 kb (Figure 4), with an average size of 700 bp. The results indicated that the quality of cDNA library should be sufficient to identify the expressed genes in yellow-fruit ginseng.

A total of 400 randomly selected clones were sequenced from the library. Of these, 378 high quality sequences

were obtained after deletion of the vector sequences and sequences with short base pairs. The obtained 378 EST sequences were submitted to GeneBank (accession No: ES672876-ES673253) and dbEST (ID: 46881386 46881763). Through BlastX analysis, 221 ESTs showed significant similarities to gene sequences in Nr database and were known genes, 21 ESTs were non-significance and unknown function genes, and 136 ESTs, which have no matched were considered novel genes in *P. ginseng*. The results of BLASTX search and annotation are shown in Table 1. The generated EST were categorized using gene ontology terms as shown in Table 2, which provide a structured vocabulary to describe a sequence according to its cellular component status, molecular function and biological process. Most EST appeared to be related to physiological and cellular processes.

In conclusion, we described the construction of a cDNA library from yellow-fruit ginseng leaf tissue, EST analysis and gene ontology. The results of analysis of 378 cDNA

Table 1. List of BlastX searched and annotated expressed sequence tags from yellow-fruit ginseng leaf cDNA library.

| Accession oo. | Identity (%) | Score (bit) | E-value | Annotation | No of EST |
|---------------|--------------|-------------|----------|---|-----------|
| ES672914 | 92 | 310 | 1.00E-83 | (S)-2-hydroxy-acid oxidase (EC 1.1.3.15) - cucurbit | 1 |
| ES673235 | 87 | 61.2 | 2.00E-10 | 19S proteasome regulatory complex subunit S6A | 1 |
| ES673125 | 81 | 66.6 | 3.00E-10 | 1-aminocyclopropane-1-carboxylate oxidase | 1 |
| ES673214 | 80 | 279 | 3.00E-74 | 1-deoxyxylulose 5-phosphate synthase | 1 |
| ES672956 | 86 | 303 | 4.00E-81 | 60S ribosomal protein L13 | 1 |
| ES673025 | 97 | 186 | 3.00E-46 | 60S ribosomal protein L37a | 1 |
| ES672934 | 79 | 183 | 3.00E-45 | GTP binding | 1 |
| ES673044 | 84 | 183 | 2.00E-45 | AT-HF | 1 |
| ES672970 | 100 | 253 | 1.00E-66 | ATP synthase beta subunit | 1 |
| ES673152 | 93 | 63.2 | 8.00E-12 | ATP synthase CF-0 subunit I | 1 |
| ES672877 | 100 | 174 | 6.00E-43 | ATPase epsilon subunit | 3 |
| ES672903 | 73 | 227 | 7.00E-59 | putative AtRer1A protein | 1 |
| ES672922 | 83 | 150 | 1.00E-35 | auxin response factor 3 | 1 |
| ES672960 | 68 | 184 | 1.00E-45 | blight-associated protein p12 precursor | 2 |
| ES673234 | 78 | 241 | 5.00E-63 | carbonate dehydratase/ zinc ion binding | 1 |
| ES673212 | 94 | 133 | 1.00E-30 | CAB-like protein [[pomoea nil] | 1 |
| ES673022 | 71 | 163 | 1.00E-39 | calcium ion binding | 2 |
| ES673029 | 60 | 156 | 4.00E-37 | calcium-binding protein | 1 |
| ES672945 | 48 | 98.2 | 7.00E-20 | Ribonuclease Mc1 | 1 |
| ES672901 | 67 | 151 | 1.00E-35 | chitinase-like protein | 1 |
| ES672917 | 99 | 275 | 8.00E-73 | chlorophyll a/b binding protein | 3 |
| ES673014 | 99 | 295 | 4.00E-79 | chlorophyll a/b binding protein of LHCII type I precursor | 5 |
| ES673007 | 94 | 193 | 2.00E-48 | Chlorophyll a-b binding protein 13, chloroplast precursor | 1 |
| ES673114 | 92 | 112 | 9.00E-49 | Chlorophyll a-b binding protein, chloroplast precursor | 1 |
| ES672969 | 90 | 134 | 9.00E-31 | chloroplast hypothetical protein | 2 |
| ES672953 | 64 | 217 | 2.00E-55 | chloroplast oxygen-evolving enhancer protein | 2 |
| ES673194 | 86 | 304 | 1.00E-81 | chloroplast pigment-binding protein CP24 | 1 |
| ES673148 | 56 | 75.1 | 7.00E-13 | conserved hypothetical protein | 1 |
| ES672983 | 70 | 213 | 2.00E-54 | CONSTANS-like protein | 1 |
| ES673055 | 78 | 132 | 5.00E-30 | copper chaperone | 1 |
| ES672886 | 70 | 270 | 7.00E-72 | CPD photolyase | 1 |
| ES673171 | 89 | 296 | 3.00E-79 | cyclophilin | 1 |
| ES673041 | 100 | 303 | 1.00E-81 | cytoplasmic ribosomal protein S13 | 1 |
| ES672961 | 64 | 132 | 5.00E-30 | dehydrin 4 | 1 |
| ES673219 | 76 | 85.1 | 7.00E-16 | delta 12 oleic acid desaturase FAD2 | 1 |
| ES673113 | 67 | 199 | 3.00E-50 | Desiccation protectant protein Lea14 homolog | 2 |
| ES673180 | 95 | 316 | 3.00E-85 | DSK2 | 1 |
| ES673046 | 77 | 195 | 7.00E-49 | electron transporter/ thiol-disulfide exchange intermediate | 1 |
| ES672999 | 77 | 164 | 1.00E-39 | elongation factor 1-beta | 1 |
| ES672981 | 71 | 115 | 5.00E-25 | enoyl-CoA-hydratase | 1 |
| ES673040 | 71 | 54.3 | 1.00E-06 | expressed protein | 1 |
| ES672915 | 82 | 253 | 9.00E-67 | ferritin | 1 |
| ES673057 | 75 | 253 | 2.00E-66 | galactinol synthase, isoform GoIS-1 | 1 |
| ES673002 | 100 | 95.5 | 5.00E-19 | GBR3 | 3 |
| ES673076 | 39 | 84.7 | 1.00E-15 | GBR5 | 5 |
| ES673205 | 73 | 192 | 3.00E-48 | GDSL-lipase protein | 1 |
| ES673079 | 94 | 105 | 3.00E-22 | geranylgeranyl reductase | 1 |
| ES673091 | 83 | 60.1 | 3.00E-08 | glossy1 homolog | 1 |

Table 1. contd.

| | | | | | |
|----------|-----|------|----------|---|----|
| ES672967 | 88 | 215 | 7.00E-55 | glutathione peroxidase | 1 |
| ES673220 | 84 | 66.6 | 2.00E-10 | glyceraldehyde 3-phosphate dehydrogenase | 1 |
| ES673215 | 78 | 220 | 2.00E-56 | glyceraldehyde 3-phosphate dehydrogenase B subunit | 1 |
| ES673075 | 84 | 278 | 5.00E-74 | glyceraldehyde-3-phosphate dehydrogenase | 1 |
| ES672997 | 52 | 109 | 5.00E-23 | Harpin-induced 1 | 1 |
| ES673176 | 81 | 171 | 1.00E-41 | heat-shock protein 80 | 1 |
| ES673042 | 46 | 145 | 9.00E-34 | histone acetyltransferase complex component | 1 |
| ES673020 | 75 | 124 | 1.00E-27 | hypothetical protein | 1 |
| ES672963 | 67 | 164 | 2.00E-39 | hypothetical protein | 1 |
| ES673016 | 47 | 65.9 | 4.00E-10 | hypothetical protein Afu4g09870 | 1 |
| ES673132 | 85 | 52 | 1.00E-07 | hypothetical protein CIMG_06034 | 1 |
| ES673048 | 92 | 123 | 4.00E-27 | hypothetical protein MtrDRAFT_AC151668g11v1 | 17 |
| ES673230 | 92 | 106 | 2.00E-37 | hypothetical protein MtrDRAFT_AC151668g27v1 | 1 |
| ES673043 | 88 | 131 | 8.00E-30 | hypothetical protein OeelhCp020 | 2 |
| ES673035 | 81 | 91.3 | 1.00E-17 | hypothetical protein OeelhCp021 | 1 |
| ES673010 | 97 | 169 | 4.00E-41 | hypothetical protein PhapfoPp090 | 2 |
| ES673039 | 63 | 121 | 1.00E-26 | hypothetical protein SNOG_04123 | 1 |
| ES673162 | 73 | 77 | 2.00E-13 | late embryogenesis abundant protein 5 | 1 |
| ES673151 | 99 | 340 | 3.00E-92 | light harvesting chlorophyll a /b binding protein | 7 |
| ES673197 | 88 | 167 | 2.00E-40 | Single-stranded DNA-binding protein | 1 |
| ES672906 | 58 | 57.4 | 1.00E-07 | low temperature and salt responsive protein | 1 |
| ES673232 | 78 | 206 | 1.00E-52 | ly200 protein | 1 |
| ES673061 | 56 | 151 | 9.00E-36 | Major sperm protein | 1 |
| ES673185 | 74 | 243 | 2.00E-63 | malate dehydrogenase | 1 |
| ES673017 | 43 | 62 | 1.00E-08 | metal ion binding | 1 |
| ES672955 | 75 | 86.7 | 4.00E-16 | metallothionein-1 like protein | 1 |
| ES672883 | 82 | 82.4 | 6.00E-15 | mRNA-binding protein precursor | 1 |
| ES673068 | 61 | 77.4 | 1.00E-13 | NAK-type protein kinase | 1 |
| ES673078 | 64 | 136 | 7.00E-31 | nonspecific lipid transfer protein 1 | 2 |
| ES672938 | 74 | 143 | 2.00E-33 | O-GlcNAc-transferase-like protein | 1 |
| ES673139 | 93 | 61.2 | 1.00E-08 | putative GMPase | 1 |
| ES673119 | 86 | 193 | 2.00E-54 | ALM beta-like | 1 |
| ES673000 | 50 | 158 | 8.00E-38 | putative adapter-related protein complex 4 epsilon 1 subunit | 1 |
| ES673030 | 78 | 53.1 | 3.00E-06 | putative lipase | 1 |
| ES673155 | 61 | 147 | 2.00E-34 | early flowering 4 | 1 |
| ES673242 | 50 | 62.4 | 6.00E-16 | hypothetical protein | 1 |
| ES673141 | 76 | 75.1 | 9.00E-13 | oxidoreductase | 1 |
| ES673071 | 94 | 296 | 1.00E-79 | oxygen evolving complex 33 kDa photosystem II protein | 1 |
| ES673240 | 39 | 59.7 | 3.00E-08 | 16 kDa protein of the photosynthetic oxygen- evolving protein | 1 |
| ES673161 | 89 | 54.7 | 1.00E-06 | PAP fibrillin | 1 |
| ES673051 | 62 | 75.5 | 6.00E-13 | peptidase/ threonine endopeptidase | 1 |
| ES673237 | 88 | 270 | 1.00E-71 | permease | 1 |
| ES673033 | 90 | 117 | 1.00E-25 | peroxiredoxin Q | 1 |
| ES673054 | 35 | 58.9 | 7.00E-08 | phloem protein 2-1 | 4 |
| ES672921 | 39 | 66.2 | 5.00E-10 | phloem protein 2-2 | 9 |
| ES673208 | 89 | 238 | 7.00E-62 | Phosphoribulokinase | 1 |
| ES673053 | 100 | 183 | 1.00E-45 | photosynthetic electron transfer-like protein | 1 |
| ES673207 | 80 | 196 | 2.00E-49 | photosystem I subunit XI | 1 |
| ES673104 | 45 | 75.1 | 1.00E-12 | photosystem II | 1 |
| ES672994 | 71 | 168 | 2.00E-41 | photosystem II 10 kDa protein | 3 |

Table 1. contd.

| | | | | | |
|----------|-----|------|----------|---|---|
| ES673116 | 41 | 72.8 | 5.00E-12 | Photosystem II 5 kDa protein, chloroplast precursor (PSII-T) | 1 |
| ES673227 | 100 | 56.6 | 3.00E-07 | photosystem II CP47 protein | 1 |
| ES673177 | 100 | 39.7 | 3.00E-06 | photosystem II M protein | 2 |
| ES673239 | 87 | 200 | 1.00E-50 | photosystem II protein D1 | 2 |
| ES673252 | 94 | 105 | 7.00E-22 | photosystem II protein K | 1 |
| ES673146 | 63 | 125 | 4.00E-28 | plastidic aldolase NPALDP1 | 1 |
| ES672927 | 52 | 95.1 | 1.00E-18 | plastoquinol-plastocyanin reductase | 2 |
| ES673221 | 92 | 140 | 2.00E-32 | poly(A)-binding protein | 2 |
| ES672984 | 84 | 79 | 5.00E-14 | Polygalacturonase-1 non-catalytic subunit beta precursor (AroGP1) | 1 |
| ES673015 | 81 | 184 | 2.00E-45 | protein binding / ubiquitin-protein ligase/ zinc ion binding | 1 |
| ES672980 | 84 | 144 | 2.00E-33 | protein disulfide isomerase | 1 |
| ES673203 | 86 | 59.7 | 4.00E-08 | PsbC | 1 |
| ES673005 | 98 | 178 | 4.00E-44 | PSI 9 kDa protein | 1 |
| ES673027 | 68 | 53.5 | 2.00E-06 | PSI-H precursor | 1 |
| ES672992 | 100 | 75.5 | 7.00E-13 | PSII 44 kDa protein | 1 |
| ES672925 | 100 | 96.3 | 3.00E-19 | PSII K protein | 1 |
| ES672896 | 52 | 112 | 8.00E-24 | structural constituent of ribosome | 1 |
| ES672978 | 72 | 221 | 2.00E-56 | putative 60S ribosomal protein L7-like protein | 1 |
| ES673059 | 82 | 232 | 6.00E-60 | Putative auxin efflux carrier component 8 (AtPIN8) | 1 |
| ES673135 | 91 | 161 | 9.00E-39 | putative chlorophyll A-B binding protein of LHCl type II precursor | 1 |
| ES673200 | 98 | 115 | 4.00E-25 | putative chloroplast chlorophyll A-B binding protein type I | 1 |
| ES672923 | 94 | 77.4 | 1.00E-13 | putative chloroplast thiazole biosynthetic protein | 1 |
| ES672946 | 76 | 157 | 3.00E-37 | putative E2, ubiquitin-conjugating enzyme UBC7 | 1 |
| ES672943 | 41 | 93.6 | 2.00E-18 | putative F-box and leucine-rich repeat protein | 1 |
| ES673154 | 76 | 213 | 2.00E-54 | putative L24 ribosomal protein | 1 |
| ES673236 | 61 | 140 | 2.00E-32 | putative phosphatidylcholine-sterol acyltransferase | 1 |
| ES672880 | 80 | 169 | 4.00E-41 | putative phosphatidylinositol- phosphatidylcholine transfer protein SEC14 | 1 |
| ES673217 | 74 | 187 | 7.00E-47 | putative photosystem I reaction centre PSI-D subunit precursor | 2 |
| ES673011 | 76 | 172 | 5.00E-42 | putative rubisco subunit binding-protein alpha subunit | 2 |
| ES673112 | 54 | 164 | 1.00E-39 | receptor-like protein kinase homolog RK20-1 | 1 |
| ES673244 | 81 | 202 | 5.00E-51 | Ribonuclease 1 | 3 |
| ES672907 | 100 | 181 | 7.00E-45 | ribosomal protein L16 | 1 |
| ES672876 | 83 | 106 | 3.00E-22 | Ribosomal protein S30 | 1 |
| ES673003 | 78 | 114 | 1.00E-24 | Ribosomal protein S5, bacterial and organelle form | 1 |
| ES672940 | 94 | 99.8 | 4.00E-20 | ribosomal protein small subunit 28 | 2 |
| ES673123 | 94 | 298 | 5.00E-83 | ribulose-1,5-bisphosphate carboxylase/oxygenase activase | 2 |
| ES673006 | 98 | 342 | 3.00E-93 | ribulose-1,5-bisphosphate carboxylase/oxygenase small subunit | 3 |
| ES673175 | 45 | 135 | 5.00E-31 | RNA-binding protein-like | 1 |
| ES673201 | 90 | 80.1 | 3.00E-14 | rRNA intron-encoded homing endonuclease | 1 |
| ES673065 | 93 | 67.8 | 9.00E-16 | S-adenosylmethionine synthetase | 1 |
| ES673173 | 93 | 69.3 | 1.00E-15 | selenium binding | 1 |
| ES672919 | 88 | 209 | 3.00E-53 | small GTP-binding protein | 1 |
| ES672924 | 75 | 88.6 | 8.00E-17 | sorbitol related enzyme | 1 |
| ES673074 | 72 | 102 | 5.00E-21 | structural constituent of ribosome | 2 |
| ES673143 | 60 | 94.7 | 1.00E-18 | T-complex protein 1 epsilon subunit | 1 |
| ES673064 | 46 | 103 | 3.00E-25 | TGF-beta receptor, type I/II extracellular region | 1 |
| ES672993 | 49 | 118 | 1.00E-25 | thioredoxin H | 2 |
| ES673225 | 74 | 150 | 1.00E-35 | Thioredoxin H-type (TRX-H) emb CAA94534.1 thioredoxin | 1 |
| ES672979 | 60 | 128 | 6.00E-29 | TIR-NBS disease resistance-like protein [Populus trichocarpa] | 1 |

Table 1. contd.

| | | | | | |
|---------------|----|------|----------|--|-----|
| ES672944 | 80 | 224 | 1.00E-57 | tonoplast intrinsic protein | 1 |
| ES673110 | 63 | 156 | 4.00E-37 | vacuolar ATPase subunit E-like protein | 1 |
| ES673164 | 76 | 80.9 | 2.00E-14 | type 2 metallothionein | 1 |
| unknown | | | | | 21 |
| no matched | | | | | 136 |
| Total | | | | | 378 |

Table 2. Gene ontology of expressed sequence tags from yellow-fruit ginseng leaf cDNA library.

| Gene ontology term | No. of genes |
|----------------------------------|--------------|
| Cell component | |
| Cell | 46 |
| Cell part | 46 |
| Organelle | 20 |
| Organelle part | 8 |
| Protein complex | 26 |
| Molecular function | |
| Antioxidant activity | 1 |
| Binding | 30 |
| Catalytic activity | 25 |
| Enzyme regulator activity | 2 |
| Molecular function unknown | 2 |
| Obsolete molecular function | 5 |
| Signal transducer activity | 1 |
| Structural molecule activity | 13 |
| Translation regulator activity | 1 |
| Transporter activity | 8 |
| Biological process | |
| Cellular process | 65 |
| Physiological process | 75 |
| Regulation of biological process | 2 |
| Response to stimulus | 4 |

demonstrated EST sequencing and data analysis as a useful and efficient approach to identifying novel genes and for functional genes expression, which would help us understand the mechanisms of ginseng plants during development stages and enrich our knowledge of functional genomic research in ginseng.

ACKNOWLEDGEMENT

This work was supported by a grant from State Forestry Administration, P.R. China (010-413255).

REFERENCES

- Choi DW, Jung JD, Ha YI, Park HW, In DS, Chung HJ, Liu JR (2005). Analysis of transcripts in methyl jasmonate-treated ginseng hairy roots to identify genes involved in the biosynthesis of ginsenosides and other secondary metabolites *Plant Cell Rep.* 23: 557-566
- Kim MK, Lee BS, In JG, Sun H, Yoon JH, Yang DK (2006). Comparative analysis of expressed sequence tags (ESTs) of ginseng leaf. *Plant Cell Rep.* 25: 599-606
- Miyahara T, Hirono I, Aoki T (2000). Analysis of expressed sequence tags from a Japanese eel *Anguilla japonica* spleen cDNA library. *Fish. Sci.* 66: 257-260
- Sun YL, Xue WZ (2001). Study on the main chemical constituents in ginseng. *Chin. J. Health Lab. Technol.* 5: 555-556. (in chinese.)
- Wang YC, Yang CP, Liu GF, Jiang J, Wu JH (2006). Generation and analysis of expressed sequence tags from a cDNA library of *Tamarix androssowii*. *Plant Sci.* 170: 28-36
- Yang CJ, Wang J, Zhou L, Liu GJ (2008). Extraction of total RNA from *Panax ginseng* C.A. Meyer. cv. Hongguo leaves. *Biotechnol. Bull.* 1: 136-139. (in chinese.)
- Zhao SJ, Liu YZ, Zhao YH, Li FY, Huang ZH, Wu LJ, Guo J, Liu JY (1998). Comprehensive evaluation on characters of Jilin yellow-fruit ginseng. *Special wild economic animal plant res.* 4: 1-6. (in chinese.)
- Zheng YL, Zhang CX, Li XG (2001). Indicators and methods of quality evaluation to Jilin ginseng. *Ginseng res.* 2: 12-14. (in chinese.)