

Full Length Research Paper

Phylogenetic analysis of human Tp53 gene using computational approach

Mohammad Haroon Khan¹, Hamid Rashid¹ and Asif Mir²

¹Department of Bioinformatics, Mohammad Ali Jinnah University, Islamabad, Pakistan.

²Department of Bioinformatics and Biotechnology, International Islamic University, Islamabad. Pakistan.

Accepted 4 November, 2010

The TP53 gene encoding p53 protein is involved in regulating a series of pathways. New discoveries about the function and control of p53 are still in progress and it is hoped to develop better therapeutics and diagnostics by exploiting this system. Evolutionary studies are of prime importance in the field of biological research since very long as provide the basis for comparative genomics. The sequence of Homo sapiens human TP53, transcript variant-1 mRNA sequence was retrieved from the NCBI in FASTA format and was studied for its relationships and percent similarity within human and others species. Genetic variation among TP53 found in human beings and other organisms were studied in detail. Multiple sequence alignment and phylogenetic analysis of the human TP53, transcript variant-1 mRNA sequence through UPGMA was performed which showed its relationship and pattern of variations among different organisms. This study will help in modern research strategies through the manipulation of p53 as its pathways are emerging rapidly and one can predict its extensive clinical use in the near future for the human benefit worldwide.

Key words: P53, tumour, cancer, phylogeny, sequence alignment.

INTRODUCTION

The *TP53* gene encoding p53 protein is involved in regulating a series of pathways including apoptosis, DNA repair, transcription, cell cycle control and genomic stability (Hussain and Harris, 2006). It has also been shown to act directly upon mitochondria through a transcription-independent pathway (Mihara et al., 2003; Park et al., 2005). The qualitative and quantitative activity of it depends on its integrity, amount and posttranslational modifications induced by different stress induced signaling pathways (Lacroix et al., 2009). Cancer research has reached an exciting phase of its evolution and investigation of the p53 tumour suppressor pathway, in particular, has become a key focus of current cancer research (Woods and Lane, 2003). New discoveries about the function and control of p53 are still in progress and it is hoped to develop better therapeutics and diagnostics by exploiting this system (Hupp et al., 2000; Lane and Hupp, 2003).

Evolutionary studies are of prime importance in the field

of biological research since very long (Pavlopoulos et al., 2010). It has become an essential element combining diverse range of research fields investigating the patterns and mechanism of evolution (Brooks et al., 2007). Phylogeny reconstruction is not only important for the organisms that house genes but is also to the evolutionary history of the genes themselves (Soltis and Soltis, 2003). Phylogenetics also provides the basis for comparative genomics (Pryer et al., 2002; Doyle and Luckow, 2003) and thus more and more molecular biologists are using phylogenetic trees to guide their sampling of taxa for comparative research (Soltis and Soltis, 2003).

Evolution of transcription network is an important genetic component in diversification between species (Shapiro et al., 2004). The evolution pattern of the p53 and related family members was studied previously which provided insights into the phylogeny of these genes and their relations (Saccone et al., 2002; Waddell et al., 2001). Modern research strategies through manipulating the p53 pathway are emerging rapidly and one can predict its extensive clinical use in the near future for the human benefit worldwide. This study was focused to explore the distribution pattern of genetic variation in p53

*Corresponding author: E-mail: drhamid@jinnah.edu.pk.

found in different organisms including man.

MATERIALS AND METHODS

Sequence retrieval

The sequence of *Homo sapiens* tumor protein p53 (TP53), transcript variant 1 mRNA was retrieved from the NCBI (<http://www.ncbi.nlm.nih.gov>) in FASTA format.

Local sequence alignment

BLAST (Altschul et al., 1990) was performed for the human TP53, transcript variant-1 mRNA sequence retrieved from NCBI to identify its relatives in different organisms including man using the online Geneious 4.8.3 software (<http://www.geneious.com>). This software takes the data in FASTA format and produces the BLAST table.

Phylogenetic analysis

Phylogenetic analysis of *H. sapiens* tumor protein p53 (TP53), transcript variant-1 mRNA sequence through UPGMA was carried out using Geneious software. Phylogenetic tree was constructed by the software showing the ancestral relationship among the sequences. The tree gives different clusters showing their relationship with each other. The sequences which lie in the same cluster are closely related.

RESULTS AND DISCUSSION

Sequence retrieval

H. sapiens tumor protein p53 (TP53), transcript variant-1 mRNA was retrieved from the NCBI in FASTA format. The sequence of the transcript variant-1 mRNA (NM_000546.4) is as the following:

>gi|187830767|ref|NM_000546.4| *Homo sapiens* tumor protein p53 (TP53), transcript variant 1, mRNA

GATTGGGGTTTTCCCTCCCATGTGCTCAAGACTGG
 CGCTAAAAGTTTTGAGCTTCTCAAAGTCTAGAGCCA
 CCGTCCAGGGAGCAGGTAGCTGCTGGGCTCCGGGG
 ACACTTTGCGTTCGGGCTGGGAGCGTGCTTTCCACG
 ACGGTGACACGCTTCCCTGGATTGGCAGCCAGACTG
 CCTTCCGGGTCACTGCCATGGAGGAGCCGCAGTCA
 GATCCTAGCGTCGAGCCCCCTCTGAGTCAGGAAACA
 TTTTCAGACCTATGGAACTACTTCTGAAAACAACG
 TTCTGTCCCCCTTGCCGTCCAAGCAATGGATGATTT
 GATGCTGTCCCCGGACGATATTGAACAATGGTTCAC
 TGAAGACCAGGTCCAGATGAAGCTCCCAGAATGCC
 AGAGGCTGCTCCCCCGTGGCCCCCTGCACCAGCAG
 CTCCTACACGGCGGCCCTGCACCAGCCCCCTCCT
 GGCCCCGTGCATCTTCTGTCCCTTCCCAGAAAACCTA
 CCAGGGCAGCTACGGTTTCCGTCTGGGCTTCTTGCA
 TTCTGGGACAGCCAAGTCTGTGACTTGCACGTACTC
 CCCTGCCCTCAACAAGATGTTTTGCCAACTGGCCAA
 GACCTGCCCTGTGCAGCTGTGGGTTGATTCCACACC
 CCCGCCGGCACCCGCGTCCGCGCCATGGCCATCT

ACAAGCAGTCACAGCACATGACGGAGGTTGTGAGGC
 GCTGCCCCCACCATGAGCGCTGCTCAGATAGCGATG
 GTCTGGCCCCCTCCTCAGCATCTTATCCGAGTGGAAG
 GAAATTTGCGTGTGGAGTATTTGGATGACAGAAACAC
 TTTTCGACATAGTGTGGTGGTGCCCTATGAGCCGCC
 TGAGGTTGGCTCTGACTGTACCACCATCCACTACAA
 CTACATGTGTAACAGTTCCTGCATGGGCGGCATGAA
 CCGGAGGCCATCCTCACCATCATCACACTGGAAGA
 CTCCAGTGGTAATCTACTGGGACGGAACAGCTTTGA
 GGTGCGTGTGTTGTGCCTGTCTGGGAGAGACCGGC
 GCACAGAGGAAGAGAATCTCCGCAAGAAAGGGGAG
 CCTCACCACGAGCTGCCCCAGGGAGCACTAAGCG
 AGCACTGCCCAACAACACCAGCTCCTCTCCCAGCC
 AAAGAAGAAACCCTGGATGGAGAATATTTACCCTT
 CAGATCCGTGGGCGTGAGCGCTTCGAGATGTTCCGA
 GAGCTGAATGAGGCCTTGGAACTCAAGGATGCCAG
 GCTGGGAAGGAGCCAGGGGGGAGCAGGGGCTCACTC
 CAGCCACCTGAAGTCCAAAAGGGTCACTACCTC
 CCGCCATAAAAACTCATGTTCAAGACAGAAGGGCC
 TGACTCAGACTGACATTCTCCACTTCTTGTCCCCAC
 TGACAGCCTCCCACCCCTCTCTCCCTCCCCTGCC
 ATTTTGGGTTTTGGGTCTTTGAACCCTTGCTTGCAAT
 AGGTGTGCGTCAGAAGCACCCAGGACTTCCATTTGC
 TTTGTCCCAGGGCTCCACTGAACAAGTTGGCCTGCA
 CTGGTGTGTTTGTGTTGGGGAGGAGGATGGGGAGTAG
 GACATACCAGCTTAGATTTTTAAGTTTTTACTGTGAG
 GGATGTTTGGGAGATGTAAGAAATGTTCTTGCAAGTA
 AGGGTTAGTTTACAATCAGCCACATTCTAGGTAGGG
 GCCCACTTACCCTACTAACCAGGGAAGCTGTCCCT
 CACTGTTGAATTTTCTCTAATTCAAGGCCCATATCT
 GTGAAATGCTGGCATTGTCACCTACCTCACAGAGTG
 CATTGTGAGGGTTAATGAAATAATGTACATCTGGCCT
 TGAAACCACCTTTTATTACATGGGGTCTAGAATTGA
 CCCCTTGAGGGTGCTTGTCCCTCTCCCTGTTGGT
 CGGTGGGTTGGTAGTTTCTACAGTTGGGCAGCTGGT
 TAGGTAGAGGGAGTTGTCAAGTCTCTGCTGGCCAG
 CCAAACCCTGTCTGACAACCTCTTGGTGAACCTTAGT
 ACCTAAAAGGAAATCTCACCCCATCCCACACCCTGG
 AGGATTTTCTCTTGTATATGATGATCTGGATCCAC
 CAAGACTTGTTTTATGCTCAGGGTCAATTTCTTTTTTC
 TTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTTT
 TCGCTTTGTTGCCAGGCTGGAGTGGAGTGGCGTGA
 TCTTGGCTTACTGCAGCCTTTGCCTCCCCGGCTCGA
 GCAGTCTGCCTCAGCCTCCGGAGTAGTCTGGGACC
 ACAGGTTTATGCCACCATGGCCAGCCAACTTTTGCA
 TTTTTGTAGAGATGGGGTCTCACAGTGTGCCCAG
 GCTGGTCTCAAACCTCTGGGCTCAGGCGATCCACCT
 GTCTCAGCCTCCCAGAGTGCTGGGATTACAATTGTG
 AGCCACCACGTCCAGCTGGAAGGGTCAACATCTTTT
 ACATTCTGCAAGCACATCTGCATTTTACCCACCCCT
 TCCCCTCCTTCTCCCTTTTTATATCCCATTTTTATATC
 GATCTCTTATTTTACAATAAACTTTGCTGCCACCTGT
 GTGTCTGAGGGGTG

Local sequence alignment

H. sapiens tumor protein p53 (TP53), transcript variant-1

Table 1. BLAST table of Human p53.

S/N	AN	E-Value	Organism	Description	Sequence Length	% Pairwise Identity
1	NM_000546	0	Homo sapiens	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 1, mRNA	2586	100.00
2	DQ191317	0	Homo sapiens	<i>Homo sapiens</i> p53 protein (TP53) mRNA, complete cds, alternatively spliced	2589	99.10
3	DQ286964	0	Homo sapiens	<i>Homo sapiens</i> p53 protein (TP53) mRNA, complete cds, alternatively spliced	2534	99.10
4	FJ207420	0	Homo sapiens	<i>Homo sapiens</i> mutant p53 mRNA, complete cds	2517	99.10
5	AB082923	0	Homo sapiens	<i>Homo sapiens</i> mRNA for P53, complete cds	2436	99.70
6	X01405	0	Human mRNA	Human mRNA fragment for phosphoprotein p53	2073	99.40
7	NM_001126115	0	Homo sapiens	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 5, mRNA	2015	100.00
8	XM_001172061	0	PREDICTED:Pan troglodytes	PREDICTED: Pan troglodytes tumor protein p53, transcript variant 2 (TP53), mRNA	2016	99.40
9	XM_001172050	0	PREDICTED:Pan troglodytes	PREDICTED: Pan troglodytes tumor protein p53, transcript variant 1 (TP53), mRNA	1997	99.30
10	K03199	0	Human p53	Human p53 cellular tumor antigen mRNA, complete cds	1684	99.90
11	AK223026	0	Homo sapiens	<i>Homo sapiens</i> mRNA for tumor protein p53 variant, clone: JTH07296	1665	99.90
12	AK225838	0	Homo sapiens	<i>Homo sapiens</i> mRNA for Cellular tumor antigen p53 variant, clone: FCC127A11	1618	99.90
13	NM_001126113	0	Homo sapiens	<i>Homo sapiens</i> tumor protein p53 (TP53), transcript variant 4, mRNA	1396	100.00
14	AK297462	0	Homo sapiens	<i>Homo sapiens</i> cDNA FLJ54297 complete cds, highly similar to Cellular tumor antigen p53	1362	99.80
15	DQ186649	0	Homo sapiens	<i>Homo sapiens</i> p53 gamma isoform (TP53) mRNA, complete cds	1406	98.60
16	DQ186648	0	Homo sapiens	<i>Homo sapiens</i> p53 beta isoform (TP53) mRNA, complete cds	1406	98.60
17	DQ186652	0	Homo sapiens	<i>Homo sapiens</i> del133 p53 gamma isoform (TP53) mRNA, complete cds	1408	98.50
18	DQ186651	0	Homo sapiens	<i>Homo sapiens</i> del133 p53 beta isoform (TP53) mRNA, complete cds	1408	98.50
19	M14694	0	Human p53	Human p53 cellular tumor antigen mRNA, complete cds	1308	99.80
20	M14695	0	Human p53	Human p53 cellular tumor antigen mRNA, complete cds	1305	99.80
21	AC007421	0	Homo sapiens	<i>Homo sapiens</i> chromosome 17, clone RP5-1030O14, complete sequence	1299	99.80
22	AY838896	0	Homo sapiens	<i>Homo sapiens</i> tumor protein p53 (Li-Fraumeni syndrome) (TP53) gene, complete cds	1299	99.80
23	BT019622	0	Homo sapiens	<i>Homo sapiens</i> tumor protein p53 (Li-Fraumeni syndrome) mRNA, complete cds	1180	99.80
24	AY888314	0	Synthetic construct	Synthetic construct <i>Homo sapiens</i> clone FLH008485.01X tumor protein p53 (TP53) mRNA, complete cds	1180	99.80
25	DQ895704	0	Synthetic construct	Synthetic construct <i>Homo sapiens</i> clone IMAGE:100010164; FLH186805.01L; RZPDo839A0962D tumor protein p53 (Li-Fraumeni syndrome) (TP53) gene, encodes complete protein	1182	99.70
26	X60011	0	Human mRNA	Human mRNA for mutated p53 transformation suppressor gene	1179	99.80
27	X60020	0	Human mRNA	Human mRNA for mutated p53 transformation suppressor gene	1180	99.70
28	AY429684	0	Homo sapiens	<i>Homo sapiens</i> tumor suppressor p53 (TP53) mRNA, partial cds	1137	99.60
29	AF456343	0	Macaca fascicularis	Macaca fascicularis p53 mRNA, complete cds	1182	96.40

Table 1. Contd.

30	U48956	0	Macaca mulatta	Macaca mulatta p53 gene, complete cds	1182	96.40
31	U48957	0	Macaca fascicularis	Macaca fascicularis p53 gene, complete cds	1182	96.10
32	EF101868	0	Homo sapiens	<i>Homo sapiens</i> cell-line L1236 nonfunctional tumor suppressor p53 (TP53) mRNA, complete cds	994	99.90
33	AF175893	0	Tupaia belangeri	Tupaia belangeri chinensis p53 tumor suppressor protein (p53) mRNA, complete cds	1206	91.10
34	EF101867	0	Homo sapiens	<i>Homo sapiens</i> cell-line L428 nonfunctional tumor suppressor p53 (TP53) mRNA, complete cds	818	100.00
35	AM076971	0	Homo sapiens	<i>Homo sapiens</i> partial mRNA for tumor protein p53 mutant form (TP53 gene), classical Hodgkin Lymphoma cell line HDLM-2	782	100.00
36	X60010	0	Human mRNA	Human mRNA for mutated p53 transformation suppressor gene	736	99.70
37	NM_001082404	0	Oryctolagus cuniculus	Oryctolagus cuniculus tumor protein p53 (TP53), mRNA >gi 1532043 emb X90592.1 O.cuniculus mRNA for p53 protein	1464	83.50
38	AF475081	0	Delphinapterus leucas	Delphinapterus leucas P53 (p53) mRNA, complete cds	1198	86.60
39	NM_213824	0	Sus scrofa	Sus scrofa p53 protein (LOC396767), mRNA >gi 115550411 dbj AK238676.1 Sus scrofa mRNA, clone:THY010016F09, expressed in thymus	1344	83.60
40	NM_214145	0	Sus scrofa	Sus scrofa tumor suppressor p53 (P53), mRNA >gi 6165622 gb AF098067.1 AF098067 Sus scrofa tumor suppressor p53 (p53) mRNA, complete cds	1204	85.00
41	NM_001009294	0	Felis catus	Felis catus tumor protein p53 (TP53), mRNA >gi 538224 dbj D26608.1 CATP53 Felis catus mRNA for p53, complete cds	1321	83.10
42	NM_174201	0	Bos taurus	<i>Bos taurus</i> tumor protein p53 (TP53), mRNA >gi 73587286 gb BC102440.1 <i>Bos taurus</i> tumor protein p53, mRNA (cDNA clone MGC:127557 IMAGE:7953339), complete cds	1346	82.50
43	AJ009673	0	Cavia porcellus	Cavia porcellus mRNA for p53 protein	1721	79.30
44	FJ855223	0	Ovis aries	Ovis aries p53 mRNA, complete cds	946	87.60
45	AF124298	0	Sus scrofa	Sus scrofa p53 protein mRNA, complete cds	941	87.60
46	D49825	0	Bos primigenius	Bos primigenius p53 mRNA, partial cds	974	86.90
47	X81704	0	B.taurus p53	B.taurus p53 mRNA	972	86.70
48	DQ656490	0	Bos taurus	<i>Bos taurus</i> cell-line TaA288 p53 mRNA, complete cds	973	86.60
49	U07182	0	Mesocricetus auratus	Mesocricetus auratus tumor suppressor p53 (p53) mRNA, complete cds	1428	80.70
50	DQ656491	0	Bos taurus	<i>Bos taurus</i> cell-line TpM803 p53 mRNA, complete cds	1173	83.60

AN = Accession number; EV = E-value; SL = Sequence length; PPI = Percent pair wise Identity.

mRNA sequence was studied for its similarity patterns and BLAST was therefore performed by feeding the FASTA format of sequence into the

online Geneious 4.8.3 software. After performing BLAST, the software produced BLAST table showing the accession numbers, percent similarity,

e-value etc (Table 1). The sequences having lowest e-value are more closely related while the difference in e-value shows the dissimilarity among




Figure 1. Phylogenetic analysis of human p53.

them.

It is clear from the results that *H. sapiens* tumor protein p53 (TP53), transcript variant 4 mRNA (NM_001126113), Expression vector Ad5CMV-p53 tumor suppressor protein p53 (p53) expression cassette, complete sequence (AF192534), *H.*

sapiens cell-line L428 nonfunctional tumor suppressor p53 (TP53) mRNA, complete cds (EF101867), *H. sapiens* partial mRNA for tumor protein p53 mutant form (TP53 gene), classical Hodgkin Lymphoma cell line HDLM-2 (AM076971) and full-length cDNA clone CS0DC002YP20 of

Neuroblastoma Cot 25-normalized of *H. sapiens* (CR608294) are 100% identical with *H. sapiens* tumor protein p53 (TP53), transcript variant 1(NM_000546.4), while *Bos taurus* tumor protein p53 (TP53), mRNA (cDNA clone MGC:127557 IMAGE:7953339), and complete cds (NM_174201)

are the most dissimilar sequences with 82% identity (Table) (Pintusa et al., 2006; Mills, 2005; Osnat et al., 2004; Kathryn et al., 2002; Shengkan et al., 2000).

Phylogenetic analysis

Phylogenetic analysis (Koref et al., 2003) often includes the search for evidence of directional selection in molecular evolution (Hsu et al., 2005; Hofmann et al., 2003; Yang and Bielawski, 2000). Evolution of the TP53 was studied in different organisms and adaptive changes were in the sequences. Phylogenetic analysis of *H. sapiens* tumor protein p53 (TP53), transcript variant-1 mRNA sequence was performed through the Geneious software using Tamura-Nei Algorithm (Tamura and Nei, 1993). The UPGMA rooted tree diagram of *H. sapiens* tumor protein p53 (TP53), transcript variant-1 mRNA sequence showed different clusters formation. Organism that originated from same ancestors having same e-value and 100% pair wise identity, are placed in same clusters whereas those which are distant from each other are placed in separate clusters.

Majority of human p53 sequences are lying in the same clusters. The genes lie in four distinct clusters while *Cavia porcellus* mRNA for p53 protein (AJ009673) is the most distinct one with 79.5% pair wise identity with the *H. sapiens* tumor protein p53. The phylogeny of the TP53 gene was also studied previously by different workers (Pintusa et al, 2006; Waddell et al, 2001; Ford, 2000). It can be concluded from the results that Human p53 may be evolved from the *C. porcellus* or both the species have evolved it from the same ancestor (Table). These results are confirmatory with the results of Villiard et al., 2007. Phylogenetic analysis was performed for the Pheromone-binding proteins (PBP) in the genus *Choristoneura* (Lepidoptera: Tortricidae) to determine either selection has acted on these genes or not (Willett, 2000). Five fish TPI sequences obtained from GenBank were used for phylogenetic analyses to describe the TPI evolution in gnathostome vertebrates (Merritt and Quattro, 2001). (Fig. 1)

REFERENCES

- Brooks DR, Jaret B, Charmaine C, David CE, Kaila EF, Jörg F, Dominik H, Stephanie H, Deborah AM, Michelle M, Linda AT, Jessica LW, Niklas W, David Z, David Z (2007). Quantitative Phylogenetic Analysis in the 21st Century. *Revista Mexicana de Biodiversidad*. 78: 225-252.
- Doyle JJ, Luckow MA (2003). The rest of the iceberg. *Legume diversity and evolution in a phylogenetic context*. *Plant Physiol*. 131: 900-910.
- Ford MJ (2000). Effects of natural selection on patterns of DNA sequence variation at the transferrin, somatolactin, and p53 genes within and among chinook salmon (*Oncorhynchus tshawytscha*) populations. *Mol. Ecol*. 9: 843-855.
- Hofmann J, Renz M, Meyer S (2003). Phylogenetic analysis of rubella virus including new genotype I isolates. *Virus Res*. 96: 123-128.
- Hsu HW, Hung YS, Huang PH, Lee LH, Liu HJ (2005). Sequence and Phylogenetic Analysis of P10- and P17-Encoding Genes of Avian Reovirus. *Avian Dis*. 49(1): 36-42.
- Hupp TR, Lane DP, Ball KL (2000). Strategies for manipulating the p53 pathway in the treatment of human cancer. *Biochem. J*. 352(1): 1-17.
- Hussain SP, Harris CC (2006). p53 biological network: at the crossroads of the cellular-stress response pathway and molecular carcinogenesis. *J. Nippon Med. Sch*. 73: 54-64.
- Kathryn JE, Jill AK, Gail SB, Marjorie ES, Charles WW, Raymond ES, Carol LR (2002). A New Invertebrate Member of the p53 Gene Family Is Developmentally Expressed and Responds to Polychlorinated Biphenyls. *Environ. Health Pers*. 110(4): 377-385.
- Koref MFS, Gangeswaran R, Santibanez KIP, Shanahan N, Hancock JM (2003). A Phylogenetic Approach to Assessing the Significance of Missense Mutations in Disease Genes. *Hum. Mutat*. 22: 51-58.
- Lacroix M, Toillon RA, Leclercq G (2009). p53 and breast cancer, an update. *Endo. Rel. Cancer*, 13(2): 293-325.
- Lane DP, Hupp TR (2003). Drug discovery and p53. *Drug Discov. Today*, 8: 347-355.
- Merritt TJS, Quattro JM (2001). Evidence for a Period of Directional Selection Following Gene Duplication in a Neurally Expressed Locus of Triosephosphate Isomerase. *Genetics*, Vol. 159: 689-697.
- Mihara M, Erster S, Zaika A, Petrenko O, Chittenden T, Pancoska P, Moll UM (2003). p53 has a direct apoptogenic role at the mitochondria. *Mol. Cell*. 11: 577-590
- Mills AA (2005). p53: link to the past, bridge to the future. *Genes Dev*. 19: 2091-2099.
- Osnat AF, Aaron A, Luba T, Konstantin A, Meytal C, Gadi K, Alma J, Ninette A, Eviatar N, Gideon R (2004). Evolution of p53 in hypoxia-stressed *Spalax* mimics human tumor mutation. *PNAS*, 101(33): 12236-12241.
- Park BS, Song YS, Yee SB, Lee BG, Seo SY, Park YC, Kim JM (2005). Phospho-ser 15-p53 translocates into mitochondria and interacts with Bcl-2 and Bcl-xL in eugenol-induced apoptosis. *Apoptosis*, 10: 193-200.
- Pavlopoulos GA, Theodoros GS, Adriano BS, Reinhard S (2010). A reference guide for tree analysis and visualization. *BioData Min*. 3(1): p. 1.
- Pintusa SS, Fomin ES, Ivanisenko VA, Kolchanov NA (2006). Phylogenetic Analysis of the p53 Family. *Biophysics*, 51(4): 571-580.
- Pryer KM, Schneider H, Zimmer EA, Banks JA (2002) Deciding among green plants for whole genome studies. *Trends Plant Sci* 7: 550-554.
- Saccone C, Barome PO, D'Erchia AM, D'Errico I, Pesole G, Sbisà E, Tullo A (2002). Molecular strategies in Metazoan genomic evolution. *Gene*. 300: 195-201.
- Shapiro MD, Marks ME, Peichel CL, Blackman BK, Nereng KS, Jonsson B, Schluter D, Kingsley DM (2004). Genetic and developmental basis of evolutionary pelvic reduction in threespine sticklebacks. *Nature*, 428: 717-723.
- Shengkan J, Sebastian M, Woo SJ, Jennifer RW, Nebojsa M, Andrej S, Mark DY, Nikola PP, Michael WY, Arnold JL (2000). Identification and characterization of a p53 homologue in *Drosophila melanogaster*. *PNAS*, 97(13): 7301-7306.
- Soltis DE, Soltis PS (2003). The Role of Phylogenetics in Comparative Genetics. *Plant Physiol*. 132: 1790-1800.
- Tamura K, Nei M (1993). Estimation of the number of nucleotide substitutions in the control region of mitochondrial DNA in humans and chimpanzees. *Mol. Biol. Evol*. 10: 512-526.
- Villiard E, Brinkmann H, Moiseeva O, Mallette FA, Ferbeyre G, Roy S (2007). Urodele p53 tolerates amino acid changes found in p53 variants linked to human cancer. *BMC Evol. Biol*. 7: p. 180.
- Waddell PJ, Kishino H, Ota R (2001). A phylogenetic foundation for comparative mammalian genomics. *Genome Inform*. 12: 141-154.
- Willett CS (2000). Evidence for Directional Selection Acting on Pheromone-Binding Proteins in the Genus *Choristoneura*. *Mol. Biol. Evol*. 17: 553-562.
- Woods YL, Lane DP (2003). Exploiting the p53 pathway for cancer diagnosis and therapy. *Hematol. J*. 4(4): 233-247.
- Yang Z, Bielawski JP (2000). Statistical methods for detecting molecular adaptation. *Trends Ecol. Evol*. 15(12): 496-503.