

Full Length Research Paper

# The use of microsatellite markers for genetic diversity assessment of genus *Hordeum* L. in Kerman province (Iran)

M. Adeli Sardou<sup>2\*</sup>, A. Baghizadeh<sup>2</sup>, A. Tavasoli<sup>1</sup> and S. Babaei<sup>3</sup>

<sup>1</sup>Biology Department, Faculty of Science, Alzahra University, Tehran, Iran.

<sup>2</sup>Biotechnology Department, International Center for Science, High Technology and Environmental Sciences, Kerman, Iran.

<sup>3</sup>Kerman Medical University of Kerman, Kerman, Iran.

Accepted 20 January, 2011

In this study, gene diversity and genetic relationships among 30 genotypes of genus *Hordeum* from Kerman province (Iran) were assessed using 10 simple sequence repeat (SSR) primers. Seven of these markers were highly polymorphic. A total of 96 alleles were detected. The number of alleles per microsatellite marker varied from 2 to 18. Except for Bmag0603, HVM36 and HvWAXY4, all the other markers showed on average, a high value of gene diversity ranging from 0.68 to 0.89. Primer GMS003 and GMS056 had the highest polymorphism information content (PIC) and so were able to distinguish different *Hordeum* genotypes. Cluster analysis based on unweighted pair group method with arithmetic averages (UPGMA) was accomplished by using the NTSYS-pc version 2.02 software for 4 major clusters which were finally extracted. Cluster 1 includes *Hordeum distichon*, *Hordeum vulgare* and *Hordeum spontaneum*, cluster 2 comprises *Hordeum leporinum* and *Hordeum glaucum*, cluster 3 has *Hordeum bulbosum* and cluster 4 includes *Hordeum brevisubulatum* ssp. *iranicum*. Data obtained from analysis are in complete agreement with taxonomic and gene pool classifications proposed previously. This study reports the first application of the SSR technique in characterization of genus *Hordeum* from Kerman province (Iran).

**Key words:** Microsatellite markers, *Hordeum*, gene diversity, genetic relationships.

## INTRODUCTION

The genus *Hordeum*, belonging to the *Triticeae* tribe, has a wide distribution through the world and comprises 32 species and 45 taxa. Thirteen species are reported from Iran, of which 7 are found in Kerman province. It is one of the 4th world important crops and has been subjected to considerable genetics studies. Genus *Hordeum* is characterized by a spike with 3 single-flowered spikelets at each rachis node (Love, 1984). On the basis of mor-

phological traits proposed by Von Bothmer and Jacobsen (1985), *Hordeum* has been divided into 4 sections, *Hordeum*, *Anisolepis*, *Stenostachys* and *Critesion* (Von Bothmer et al., 1991). Based on cross ability, the genus *Hordeum* can be divided into 3 genetic pools. The primary gene pool comprises *Hordeum vulgare*, *Hordeum spontaneum* and *Hordeum distichon*. The secondary gene pool is represented solely by *Hordeum bulbosum*. The tertiary gene pool comprises all the other *Hordeum* species. Cytological and molecular tools have been used extensively to study the relationships of the *Hordeum* species. Cytogenetic studies based on meiotic pairing among different diploid species of *Hordeum* showed four basic genomes: genome I common to *H. vulgare* and *H. bulbosum* (Kasha and Sadasivaiah, 1971; von Bothmer et al., 1983), genome X of the *Hordeum marinum* group (von Bothmer et al., 1986a, b), genome Y from *H.*

\*Corresponding author. E-mail: [m\\_adeli622002@yahoo.com](mailto:m_adeli622002@yahoo.com).  
Tel: 0098- 9133995437.

**Abbreviations:** PCR, Polymerase chain reaction; SSR, simple sequence repeat; GD, genetic distances; PIC, polymorphism information content; RFLP, restriction fragment length polymorphism; STS, sequence-tagged site.

**Table 1.** List of *Hordeum* genotypes examined for genetic relatedness using SSR marker system.

No.	Code	Genotype	Location	No.	Code	Genotype	Location
1	H. vul1	<i>H. vulgare</i>	Joopar	16	H. gla3	<i>H. glaucum</i>	Kerman
2	H. vul2	<i>H. vulgare</i>	Dehbakri	17	H. gla4	<i>H. glaucum</i>	Estakhroie
3	H. vul3	<i>H. vulgare</i>	Jiroft	18	H. gla5	<i>H. glaucum</i>	Kohpayeh
4	H. vul4	<i>H. vulgare</i>	Mahan	19	H. gla6	<i>H. glaucum</i>	Baft
5	H. dis1	<i>H. distichon</i>	Baft	20	H. gla7	<i>H. glaucum</i>	Baghin
6	H. dis2	<i>H. distichon</i>	Baft	21	H. bre1	<i>H. brevisubulatum</i> ssp. <i>iranicum</i>	Baft
7	H. dis3	<i>H. distichon</i>	Kerman	22	H. bre2	<i>H. brevisubulatum</i> ssp. <i>iranicum</i>	Lalezar
8	H. spo	<i>H. spontaneum</i>	Sarasiab	23	H. bre3	<i>H. brevisubulatum</i> ssp. <i>iranicum</i>	Lalezar
9	H. spo	<i>H. spontaneum</i>	Estakhroie	24	H. bre3	<i>H. brevisubulatum</i> ssp. <i>iranicum</i>	Lalezar
10	H. spo	<i>H. spontaneum</i>	Sirjan	25	H. lep1	<i>H. leporinum</i>	Baghin
11	H. bul1	<i>H. bulbosum</i>	Dehbakri	26	H. lep2	<i>H. leporinum</i>	Estakhroie
12	H. bul2	<i>H. bulbosum</i>	Dokol	27	H. lep3	<i>H. leporinum</i>	Estakhroie
13	H. bul3	<i>H. bulbosum</i>	Maskoon	28	H. lep4	<i>H. leporinum</i>	Kerman
14	H. gla1	<i>H. glaucum</i>	Sarasiab	29	H. lep5	<i>H. leporinum</i>	Kabootarkahn
15	H. gla2	<i>H. glaucum</i>	Kerman	30	H. lep6	<i>H. leporinum</i>	Rafsanjan

*murinum* and the ge-nome H, shared by Asiatic, North and South American diploid barley (Dewey, 1984; Love, 1984; von Bothmer et al., 1986a). The genome designation was changed and Xa corresponds to X and Xu corresponds to Y according to von Bothmer et al. (1986a). Since the mid 1980s, ge-nome identification has progressed rapidly with the help of polymerase chain reaction (PCR) technology. Mole-cular polymorphism has been developed and used in genetic and breeding studies in many species of plants (Hu and Quiros, 1991; Svitashv et al. 1994; Bell and Ecker, 1994; Matsumoto and Fukui, 1996; Guilford et al., 1997; Haddad, 2008). Microsatellites as a marker have a high potential in using genome and genetic analysis of self-pollinating crops such as wheat, because of their high degree of polymorphism (Roder et al., 1995). Seve-ral studies on estimation of genetic diversity among *Hordeum* species have been carried out, for example, Graner et al. (1994), Saghai Maroof et al. (1994), Selbach and Cavalli-Molina (2000) and Struss and Plieske, (1998).

The objectives of this study are to define genetic diversity and relationship among species of the genus *Hordeum* using SSR markers and to determine the efficiency of the markers for further genome analysis.

## MATERIALS AND METHODS

### Plant material and DNA extraction

Leaf samples of 30 *Hordeum* genotypes (wild and cultivated forms) from Kerman (Table 1) were evaluated using 10 simple sequence repeat primers (Table 2). Genomic DNA was extracted by mini prep\_ isolation method (Dellaporta and Hicks, 1983) with minor modifications. Approximately 0.2 g of young leaves from 8 mature plants of each accession was pooled for the extraction of genomic DNA. The leaves were frozen in liquid N<sub>2</sub>, mixed with 400 µl of extraction buffer (50 mM Trisbase pH 8, 300 mM NaCl, 25 mM

EDTA pH 8 and 1% SDS) and incubated at 65 °C for 30 min. 200 µl sodium acetate 5 mM was added to each tube and placed about 10 min on ice. 500 µl chloroform/ isoamyl alcohol (24:1) was added and mixed well. The mix was centrifuged at 12000 g for 15 min. The supernatant was precipitated with an equal volume of ice-cold isopropanol and centrifuged at 5000 g for 15 min. At this stage, DNA was recovered by centrifuging. The pellet was hooked out by sterile pipettes, washed in 70% ethanol and air dried and suspended in 300 µl of 1x TBE buffer. DNA quantity and quality were estimated by both using an UV spectrophotometer (Carry 50) by measuring absorbencies at A260 and A280 nm and 1% agarose gel electrophoresis by comparing band intensity with DNA ladder of known concentrations. DNA samples were diluted to 50 ng/µl for simple sequence repeat (SSR) reactions.

### SSR reaction

PCR reactions were performed in 25 µl reaction volumes containing 2.5 µl PCR buffer (1x), 0.75 µl MgCl<sub>2</sub> (1.5 mM), 0.5 µl of each dNTPs (200 µM), 2 µl of each primer (10 pmol) , 0.2 µl of Taq polymerase (1 Unit) (Fermentas), 17.05 µl of deionized water and 2 µl of genomic DNA (50 ng/µl). Amplifications were performed in a gradient thermal cycler (Eppendorf, Hamburg) and the program included 1 cycle of 4 min at 94°C, followed by 35 cycle of 1 min at 94°C, 30 s at 46 to 58°C (depending on primer) and 45 s at 72°C, followed by a final extension for 5 min at 72°C.

The PCR products were sepatated on sequencing gel containing 8% acrylamide (AA : BIS = 19:1) in 1x TBE buffer. For this 5 µl PCR product plus 2 µl loading dye (19 ml formamide (95%), 1 ml NaOH (10 mM), 0.05 g bromophenol blue (0.05%) (w/v) and 0.05 g xylene cyanol (0.05%) (w/v), were mixed and the gel was run for approximately 4 h at 100 W. After running, glass plate covering the gel was silver stained according to Bassam et al. (1991).

### Data analysis

The amplified bands were scored manually as 0 (absent) or 1 (present). Genetic distances (GD) between samples were calculated using the DICE (equivalent to Nei and Li (1979) algorithm) based on the probability that the amplified fragment from one

**Table 2.** Primers sequences and annealing temperature of 10 *Hordeum* microsatellite markers.

Marker	Primer sequences (5'→3')	Annealing temperature (°C)
GMS003	F: TTTCAGCATCACACGAAAGC R: TTGCATGCATGCATACCC	5
GMS027	F: CTTTTTCTTTGACGATGCACC R: TGAGTTTGTGAGAACTGGATGG	51
GMS056	F: GAGAAACGCAGCTGTGGC R: GTCACCGAGGCCTTCCTC	50
Bmag13	F: AAGGGGAATCAAATGGGAG R: TCGAATAGGTCTCCGAAGAAA	60
Bmag0603	F: ATACCATGATACATCACATCG R: GGGGTATGTACGACTAATA	60
Bmac40	F: AGCCCGATCAGATTTACG R: TTCTCCCTTTGGTCCTTG	56
Bmac192	F: GGGTGGTGTGCTTAAAC R: TCAACATTCATCTACCACCA	56
Bmac306	F: CCTTGTGTGAGTGTGTGTGT R: ACATGCACATGAACTAATCAA	59
HVM36	F: TCCAGCCGACAATTTCTTG R: AGTACTCCGACACCACGTCC	57
HvWAXY4	F: AGTATCGCAGACGCTCAC R: GTTATGTA CTGCTCGCTC	59

genotype will be present in another genotype,  $GD = 1 - 2a/2a+b+c$ , (a: number of shared fragments, b: number of fragments in line A, and c: number of fragments in line B). Dendrogram was constructed using Cluster method of the Unweighted Pair Group method with Arithmetic Averages (UPGMA) by NTSYS\_pc 2.02i (Rohlf 1998). Cut line was designed using molecular analysis of variance (MANOVA) method. Gene diversity (heterozygosity) was calculated according to Weir (1990):

$$\text{Gene diversity} = 1 - \sum P_{ij}^2$$

Where,  $P_{ij}$  is the frequency of the  $j$ th pattern for microsatellite marker and is summed across in patterns.

Anderson et al. (1993) suggest that gene diversity is the same as the polymorphism information content (PIC). Diversity can also be calculated by Shannon's Information Index:  $H' = -\sum P_i \ln p_i$ ,  $P_i$  is the allele frequency. These analyses were performed by POPGENE 32 software.

## RESULTS AND DISCUSSION



The results of molecular assay in studying 30 *Hordeum* genotypes by use of 10 microsatellite markers are divided into 2 parts: 1) Determining the efficiency of markers and 2) genetic relationships among genotypes. All primers were also tested for their efficiency to reveal polymorphism in different *Hordeum* species by calculating allele length, number of alleles, PIC, heterozygosity and Shannon's information index (Table 3). The amplified fragments length was from 102 to 379 bp. The number of alleles per marker varied from 2 to 18. PIC ranged from


0.29 to 0.89. Seven of the markers were highly polymorphic in this study. Except for HVM36, Bmag0603 and HvWAXY4, all other markers showed a high value of gene diversity (Table 3). GMS003 (Figure 1A) and GMS056 (Figure 1B) exhibited the highest gene diversity (0.89 and 0.80) among the markers, so these markers could separate the genotypes better and can be useful candidates for further studies. The number of alleles detected by microsatellite markers correlated with the gene diversity, for example the marker Bmag0603 showed an average gene diversity of 0.42, detecting 3 alleles within *Hordeum* genotypes, whereas GMS003 detected 18 alleles and showed a gene diversity value of 0.89. This finding was in contrast with Struss and Plieske, (1998) who used 15 microsatellite markers to estimate the genetic diversity among 163 barley genotypes. Hadad et al. (2008) working with grape also found correlation between number of alleles and gene diversity. Graner et al. (1990) and Selbach and Cavalli-Molina (2000) using restriction fragment length polymorphism (RFLP) and randomly amplified polymorphic DNA (RAPD) marker, respectively, observed a relatively low diversity within barley germplasm.

Genetic relationships among the accessions were further studied by cluster analysis (Figure 2). The dendrogram discriminated between 30 barley genotypes. The pattern of cluster analysis based on DICE similarity coefficient and UPGMA algorithm placed the genotypes into 4 major clusters. The cut line was at point 0.05.

**Table 3.** Allele length, number of alleles, gene diversity and Shannon's information index of 10 microsatellite markers in 30 *Hordeum* genotypes.

Marker	Allele length in bp	Number of alleles	PIC (heterozygosity)	Shannon's information index
GMS003	103-379	18	0.89	2.9201
GMS027	129-197	11	0.79	2.7013
GMS56	124-184	13	0.80	2.8468
Bmag13	150-186	7	0.68	2.0083
Bmag0603	127-188	3	0.42	1.2346
Bmac40	202-283	8	0.71	2.4592
Bmac192	133-187	10	0.73	2.8025
Bmac306	112-158	8	0.70	2.4602
HVM36	102-158	4	0.36	1.1129
HvWaxy4	127-183	2	0.29	1.0002

**A****B****Figure 1.** PCR product of microsatellite marker GMS056 showing 13 (A) and 18 (B) polymorphic alleles among 1 to 15 different *Hordeum* genotypes. The allele length is in bp



**Figure 2.** UPGMA dendrogram based on Dice similarity coefficient illustrating the relationships among 30 *Hordeum* genotypes.

Cluster 1 was represented by *H. distichon*, *H. vulgare* and *H. spontaneum*. Cluster 2 comprises *H. leporinum* and *H. glaucum*. Cluster 3 includes *H. bulbosum* as its sole representative. Cluster 4 comprises by *H. brevisubulatum* ssp. *iranicum*. Our result based on SSR markers was consistent with taxonomic classification based on numerous morphological (Bor, 1970), cytological and cross ability characters, delineated by and sequence-tagged site (STS) markers (Terzi et al., 2001). All *Hordeum* species with the I genome clustered together.

Our results grouped *Hordeum leporinum* and *Hordeum glaucum*, which all share the Xu genome according to

von Bothmer et al. (1986a). Position of *H. bulbosum* confirms the findings of Svitashv et al. (1994), who used six cloned barely repetitive DNA sequences, regarding the clear distinction between *H. vulgare* and *H. bulbosum*. This was despite the attribution of the I genome to both species. In the dendrogram, all *Hordeum brevisubulatum* ssp. *iranicum* genotypes were grouped together, which was according to cross ability and cytological studies. Cluster analysis could clearly identify the genetic relationships of the *Hordeum* genotypes examined in each set and demonstrates the potential and ability of microsatellite markers for genome analysis in the genus *Hordeum*. Totally, Our results using SSR markers are

in agreement with taxonomical classification based on cross ability and morphological studies (Sahebi, 2002), and cytogenetic studies based on meiotic pairing except for the *H. bulbosum* and molecular studies of Svitashv et al. (1994) and Struss and Plieske, (1998). In this study, microsatellite markers were able to distinguish between different *Hordeum* genotypes. The high gene diversity value of markers (with an average of 0.75) makes them ideal markers for differentiation between *Hordeum* genotypes. Similarly, high values of gene diversity were reported for other plant species (Saghai et al., 1994; Rongwen et al., 1995; Szewc-McFadden et al., 1996; Guilford et al., 1997; Hernandez et al., 2001; Najafi et al., 2006; Hadad et al., 2008). The high degree of polymorphisms in our microsatellite markers is not only because of the germplasm tested but is also based on the variable character of microsatellites. The use of microsatellite markers employed in this study demonstrates the usefulness of these markers for identification of *Hordeum* genotypes. They provide an optimal system which can increase the efficiency of germplasm evaluation in the *Hordeum* germplasm collection of Kerman province (Iran).

## REFERENCES

- Anderson JA, Churchill GA, Autrique JE, Tanksley SD, Sorrells ME (1993). Optimizing parental selection for genetic linkage maps. *Genome*, 36: 181-186.
- Bassam BJ, Caetano-Anolles G, Gressho PM (1991). Fast and sensitive silver staining of DNA in polyacrylamide gels. *Appl. Biochem. Biotechnol.* 42: 181-188.
- Bell CJ, Ecker JH (1994). Assignment of 30 microsatellite loci to the linkage map of *Arabidopsis*. *Genomics*, 19: 137-144.
- Bor NL (1970). Graminea. In: Flora Iranica. Vol. 70 In: Rechinger KH ed. Graz.
- von Bothmer R, Flink J, Jacobsen N, Kotimaki M, Landstrom T (1983). Interspecific hybridization with cultivated barley (*Hordeum vulgare* L.). *Hereditas*, 99: 219-244.
- von Bothmer R, Flink J, Landstrom T (1986a). Meiosis in interspecific *Hordeum* hybrids. I. Diploid combinations. *Can. J. Genet. Cytol.* 28: 525-535.
- von Bothmer R, Giles BR, Jacobsen N (1986b). Crosses and genome relationships in the *Hordeum patagonicum* group. *Genetica*, 71: 75-80.
- von Bothmer R, Jacobsen N, Baden C, Linde-Laursen I, Jorgensen RB (1991). An ecogeographical study of the genus *Hordeum*. Systematic and ecogeographic studies on crop gene pools 7. International Board for Plant Genetics Resources, Rome, 2nd ed.
- Dellaporta SL, Hicks JB (1983). A plant DNA miniprep: version II. *Plant Mol. Biol. Rep.* 1(14): 19-21.
- Dewey DR (1984). The genome system of classification as a guide to hybridization with the perennial *Triticeae*. In: Gustafson JP (ed.). *Gene Manipulation in Plant Improvement*. Plenum Publ. Corp. New York. pp. 209-279.
- Graner A, Siedler H, Jahoor A, Herrmann RG, Wenzel G (1990). Assessment of the degree and the type of restriction fragment length polymorphism in barley (*Hordeum vulgare*). *Theor. Appl. Genet.* 80: 826-832.
- Graner A, Ludwig WF, Melchinger AE (1994). Relationship among European barley germplasm. II. Comparison of RFLP and pedigree data. *Crop Sci.* 34: 1199-1205.
- Guilford P, Prakash S, Zhu JM, Rikkerink E, Gardiner S, Bassett H, Forster R (1997). Microsatellites in *Malus domestica* (apple): abundance, polymorphism and cultivar identification. *Theor. Appl. Genet.* 94: 245-249.
- Hadad R, Ramezani A, Mardi M (2008). Genetic diversity of some Iranian grapes using microsatellite markers. *Iran. J. Genet.* 4(2): 31-38.
- Hernandez P, Dorado G, Laurie DA, Martln A, Snape JW (2001). Microsatellites and RFLP probes from maize are efficient sources of molecular markers for the biomass energy crop *Miscanthus*. *Theor. Appl. Genet.* 102: 616-622.
- Hu J, Quiros CF (1991). Identification of broccoli and cauliflower cultivars with RAPD markers. *Plant Cell Rep.* 10(10): 505-511.
- Kasha K, Sadasivaiah RS (1971). Genome relationships between *Hordeum vulgare* L. and *H. bulbosum*. *Chromosoma*, 35: 264-287.
- Love A (1984). *Conspectus of the Triticeae*. Feddes Rep. 95: 425-521.
- Matsumoto S, Fukui H (1996). Identification of rose cultivars and clonal plants by random amplified polymorphic DNA. *Scientia Horticulturae*, 67(1-2): 49-54.
- Nei M, Li WH (1979). Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. USA.* 76: 5269-5273.
- Najafi J, Alipanah L, Ghareyazie B, Mohammadi SA, Hagh-Nazari A (2006). Genetic diversity of Iranian and some European grapes revealed by microsatellite markers. *Iran. J. Biotechnol.* 4(1): 36-44.
- Roder MS, Plaschke J, Konig SU, Borner A, Sorrells E, Tanksley SD, Ganal M (1995). Abundance, variability and chromosomal location of microsatellites in wheat. *Mol. Genet.* 246: 327-333.
- Rohlf FJ (1998). *Ntsys-pc: Numerical taxonomy: the principles and practice of numerical classification*. WH. freeman and company, San Francisco.
- Rongwen J, Akkaya MS, Bhagwat AA, Lavi U, Gregan PB (1995). The use of microsatellite DNA markers for soybean genotype identification. *Theor. Appl. Genet.* 90: 43-48.
- Saghai Maroof MA, Biyashev RM, Yang GP, Zhang Q, Allard RW (1994). Extraordinarily polymorphic microsatellite DNA in barley: species diversity, chromosomal locations and population dynamics. *Proc. Natl. Acad. Sci. USA.* 91: 5466-5470.
- Sahebi SJ (2002). Systematic and biosystematic study of genus *Hordeum* L. from Iran. Phd thesis, Tehran university, Iran.
- Selbach A, Cavalli-Molina S (2000). RAPD characterization of Brazilian barley (*Hordeum vulgare* ssp. *vulgare*) varieties. *Euphytica*, 111: 127-135.
- Struss D, Plieske J (1998). The use of microsatellite markers for detection of genetic diversity in barley populations. *Theor. Appl. Genet.* 97: 308-315.
- Svitashv S, Bryngelsson T, Vershinin A, Pedersen C, Sall T, von Bothmer R (1994). Phylogenetic analysis of the genus *Hordeum* using repetitive DNA sequences. *Theor. Appl. Genet.* 89: 801-810.
- Szewc-McFadden AK, Kresovich S, Blied SM, Mitchell SE, McFerson JR (1996). Identification of polymorphic, conserved simple sequence repeats (SSRs) in cultivated *Brassica* species. *Theor. Appl. Genet.* 93: 534-538.
- Terzi V, Pecchioni N, Faccioli P, Kucera L, Stanca M (2001). Phyletic relationships within the genus *Hordeum* using PCR-based Markers. *Genet. Resour. Crop Evol.* pp. 1-10.
- Weir BS (1990). *Genetic data analysis methods for discrete genetic data*. Sinauer assoc. Inc. Sunderland, Massachusetts. USA.