

*Full Length Research Paper*

# Comparison of two different endurance training methods on glutathione s-transferase level

Murat Tas

Physical Education and Sports School, Agri Ibrahim Cecen University, 04100, Agri-TR, Turkey. E-mail: murattas25@gmail.com. Tel: +904722159248. Fax: +904722162036.

Accepted 24 March, 2011

This study compared the effects of different training methods (continuous and interval) carried out in a hot environment on glutathione s-transferase level (GST). The study group comprised 30 male volunteers who do not exercise regularly. The participants were categorized into 2 groups, continuous running (CRG, n: 15) and interval running (IRG, n: 15). The participants followed a training program on 3 days per week for 8 weeks. Comparison of the pre- and post-training samples showed a statistically significant increase in GST value ( $P < 0.01$ ) within the interval running group (IRG) and a significant decrease in GST value ( $P < 0.01$ ) within the continuous running group (CRG). GST values were compared on the basis of the pre-test and post-test results of various training methods carried out in a hot environment. In this scope, statistically significant differences were observed between the groups in the pre-tests ( $p < 0.05$ ) and the post-tests ( $p < 0.01$ ). As a result, it can be said that the CRG in a hot environment had a greater effect on oxidative stress by increasing lipid peroxidation, and the IRG prevented the formation of free radicals by producing a positive increase in the glutathione s-transferase level.

**Key words:** Hot environment, endurance training, glutathione s-transferase, oxidative stress.

## INTRODUCTION

Oxygen consumption inherent in cell growth leads to the generation of a series of reactive oxygen species (ROS) (Gülçin et al., 2003a, 2004a and b; Oktay et al., 2003). They are continuously produced by the body's normal use of oxygen such as respiration and some cell-mediated immune functions. ROS include free radicals such as superoxide anion radicals ( $O_2^{\cdot-}$ ), hydroxyl radicals ( $OH\cdot$ ) and non-free radical species such as hydrogen peroxide ( $H_2O_2$ ) and singlet oxygen ( $^1O_2$ ) (Gülçin, 2006a, and b, 2007). ROS are continuously produced during normal physiological events and can easily initiate the peroxidation of membrane lipids, leading to the accumulation of lipid peroxides (Gülçin; 2010; Gülçin and Dastan, 2007; Balaydin et al., 2010;

Şerbetçi and Gülçin, 2010; Gülçin et al., 2010a; 2010b). ROS are also capable of damaging crucial biomolecules such as nucleic acids, lipids, proteins and carbohydrates and may cause DNA damage that can lead to mutations. If ROS are not effectively scavenged by cellular constituents, they lead to disease conditions. ROS have been implicated in more than 100 diseases (Halliwell and Gutteridge, 1990; Gülçin et al., 2004c, 2005a and b, 2006a and b).

All aerobic organisms have antioxidant defences, including antioxidant enzymes and antioxidant food constituents, to remove or repair the damaged molecules. Antioxidant compounds can scavenge free radicals and oxidative damage (Ak and Gülçin, 2008; Gülçin et al., 2007a, 2009a, 2010c; Talaz et al., 2009). The level of oxidative damage which can occur during physical training is not only determined by the formation of free radicals, but also by the defence capacity of antioxidants. While it is expressed that especially acute, intensive training may cause oxidative stress, it is suggested that regular endurance training can reduce oxidative stress and muscle damage after the exercise,

---

**Abbreviations:** ROS, Reactive oxygen species; **ETDA**, ethylenediaminetetraacetic acid; **ELISA**, enzyme-linked immunosorbent assay; **MDA**, malondialdehyde; **CRG**, continuous running group; **IRG**, interval running group; **GST**, glutathione s-transferase.

and develop antioxidant defence capacity (Finaud et al., 2006; Galassetti et al., 2006; Elosua et al., 2003).

Physical exercises require new arrangements in the organism. However, physiological adaptations becomes more important if the organism, which is in a more intensive arrangement compared with the resting position, is exposed to stress resulting from a different sources. Thus, the adaptations of organisms to altitude, underwater, in hot and cold environments, the physiological stresses and the precautions are discussed (Goto et al., 2003; Durstine et al. 2001; Stein and Colditz 2004).

The effects of climate change are evident in every aspect of our lives. Increased temperatures associated with climate change may exacerbate the negative effects of outdoor sporting activities. The pre-season preparation period of many sports generally coincides with the summer months.

Depending on the climatic conditions of this season, training carried out in hot weather increases the importance of the issue. These training programs specifically include exercises that improve endurance. These endurance exercises use a range of different training models, which often use interval and continuous running methods. Also, the effects of the interval and continuous running methods on the antioxidants and oxidative stress, two of the physiological changes occurring in our body, in hot weather are subjects of debate.

In the light of the earlier information, the purpose of the present study is to compare the effects of continuous running and interval running models, two training methods aimed at improving endurance, which are often carried in a hot environment during pre-season preparation, on glutathione s-transferase level and to assess the results within the framework of the literature.

## MATERIALS AND METHODS

### Subject selection

Thirty male students from Physical Education and Sports School, Atatürk University, participated in this study. The subjects were categorized into 2 groups, termed continuous running (CRG, n: 15) and interval running (IRG, n: 15). The subjects trained for three days per week for 8 weeks in a hyperthermic environment in which the average temperature was 29 to 34°C. The training was carried out on the athletic field of Physical Education and Sports School, Atatürk University.

The body weights of the subjects were measured (barefooted, wearing t-shirt and tights) using a bascule with 0.01 kg sensitivity. Weather temperature and humidity were taken from official data from the Erzurum Provincial Directorate of State Meteorology.

### Exercise program

The target number of heart beats of the subjects in the continuous running group was determined by means of the Karvonen method (Özer, 2006) and the subjects followed 25 to 60 min duration running exercises with 50 to 70% intensity on three days per week

for 8 weeks. The maximal running times over distances of 250, 400, 650 and 900 m were determined for each subject, and a common interval training program was applied at 250, 400, 650 and 900 m; subjects were required to run using pyramidal loading method at an intensity of 60 to 80% (250, 400, 650, 900, 650, 400 and 250 m). The interval-training group also exercised for 3 days per week for 8 weeks. The exercise was applied until the heart rate reduced to 120 to 130 between the loadings. In order to make the subjects adapt to the training, the interval training program was applied as 1 set for the first two weeks, 2 sets from the third week to the seventh week and 3 sets in the last two weeks. Both groups completed warm up exercises for 5 to 10 min before starting the training and 5 to 10 min of cooling down exercises after the training.

### Blood sample

Blood uptake and haemolyzate preparation were performed as in previous studies (Beydemir et al., 2003; Gülçin et al., 2004d; Beydemir and Gülçin, 2004; ArasHisar et al., 2004; Hisar et al., 2005a and b). Blood samples were taken 2 days before and 2 days after the two different training programs, each of which lasted for 8 weeks, for comparison. The samples were taken into normal biochemistry and ethylenediaminetetraacetic acid (ETDA) tubes. The samples taken into the ETDA tubes were inverted 3 to 5 times. After the samples in the biochemistry tubes were left at room temperature for 20 min, they were stored at -80°C prior to analysis. For the analysis, the samples were centrifuged at 3500 rpm for 5 min to precipitate the shaped particles (Gülçin et al., 2005b, 2008a, 2009; Çoban et al., 2008, 2009; Beydemir et al., 2005; Şentürk et al., 2008; 2009; Öztürk Sarıkaya et al. 2010).

### Glutathione s-transferase level (GST) activity assay

GST assay reactive was adjusted at 36.20 ml 0.1 M K-phosphate buffer and pH was taken as 6.5 on the day of measurements. At that time, the microplate which was sample pipetted was covered up and kept at 25°C, the assay reactive was mixed slightly after being incubated at 37°C for 3 min and its kinetic reading was carried out in an enzyme-linked immunosorbent assay (ELISA) Reader at 340 nm. The results were printed out as GST (U/L) after being calculated from the standard graph (Donald et al., 1967).

### Statistical analysis

The data analysis was carried out using the Statistical Package for the Social Sciences (SPSS) (version 15.0) statistical analysis program. The arithmetic averages and the standard deviations of the data were calculated and given as descriptive statistics. The Mann-Whitney U test, which is a nonparametric test, was used to examine the differences between independent groups, and the Wilcoxon test, which is a nonparametric test, was used to examine the differences between the dependent groups. Values of  $p < 0.01$  and  $p < 0.05$  were taken as the significance level

## RESULTS

This study compared the effects of two different endurance-training programs (interval and continuous running), conducted in hot conditions, on glutathione s-transferase enzyme. The study was carried out in an average temperature of  $29.40 \pm 1.49^\circ\text{C}$  and average humidity level was  $50.71 \pm 8.46\%$ . The average and

**Table 1.** Average and standard deviation of continuous and interval running groups before and after exercises.

Variables	CRG <sup>§</sup>	IRG <sup>§</sup>
Age (year)	22.73 ± 3.51	24.27 ± 2.71
Height (cm)	1.73 ± 0.06	1.75 ± 0.06
Weight (kg)	Pre-test	71.99 ± 10.70
	Post-test	70.30 ± 12.57**
		75.35 ± 7.14
		73.46 ± 6.89*

\*p < 0.05, \*\*p < 0.01, <sup>§</sup>: Mean ± SD, IRG: Interval running group, CRG: Continuous running group.

**Table 2.** Comparison of pre-test and post-test GST level of the interval running group.

Variable	Interval running group (IRG)				Z
	Pre-test		Post-test		
	X	SS	X	SS	
<b>GST (U/L)</b>	10.80	1.08	12.94	0.48	-3.181**

GST: Glutathione S-transferase; \*\*p < 0.01.

**Table 3.** Comparison of pre-test and post-test GST level of the continuous running group.

Variables	Continuous running group (CRG)				Z
	Pre-test		Post-test		
	X	SS	X	SS	
<b>GST (U/L)</b>	12.38	2.03	10.13	2.05	-3.408**

GST: Glutathione S-transferase; \*\*p < 0.01.

standard deviation of continuous and interval running groups before and after exercises is given in Table 1. The results in Table 2 shows that after training was carried in a hot environment, there was a significant increase in the level of GST of the IRG (p < 0.01). The results in Table 3 show that, after training in a hot environment, there was a significant decrease in GST value of the CRG group (p < 0.01).

The pre-test and post-test GST values were compared according to the different training methods used in a hot environment. Statistically significant differences between the groups were seen in the pre-tests (p < 0.05) in comparison with the post-test results (p < 0.01)

Tissue damage can occur during sports activities, depending on muscle damage, thermal temperature and ischemia reperfusion. The sports scientists depend on the ROS production during sports activities and some elements such as mitochondrial electron transfer chain, xanthine oxidase system, the metal catalyzed reactions and activated neutrophils (Peake and Suzuki, 2004). It is thought that malondialdehyde (MDA) causes oxidative stress in direct proportion to the intensity and the duration of the exercises and increases lipid peroxidation reactions. In situations in which oxygen use is low, the superoxide radical and its derivatives are deactivated by the antioxidant defence. However, in an

exercise in which oxygen consumption increased considerably, these defence mechanisms may not keep up with the formation of free radicals and this may result in cell damage (Cheeseman and Slater, 1993). If the level of free radicals exceeds the antioxidant capacity, the fats, proteins and other parts of the cells will be oxidized (Clarkson and Thompson, 2000; Smith and Miles, 2000). Endurance exercises were used in most of the experimental studies examining the oxidative stress which results from exercise and its exact effect on plasma antioxidants or fat peroxidation (Sanchez et al., 1995).

Instances of heat exhaustion occurring during and after exercise cause the increase of reactive oxygen species (Salo et al., 1991). In a study by Osorio et al. (2003), it was suggested that exercise under thermal stress increased the effectiveness of antioxidants; also, although the metabolic responses continued, a mother was exposed daily to extreme heat stress, which had a dangerous effect on the baby and it was necessary, when swimming, to avoid hot water, since it could damage foetal development

In the study carried out in a hot environment, it was observed that while there was a significant increase in the GST value of the IRG before and after the training (p < 0.01 level, Table 2), there was a significant decrease in

**Table 4.** Comparison of pre-test and post-test GST values between groups.

Variable		Group comparison <sup>§</sup>		Z
GST(U/L)	Pre-test	11.59	1.79	-2.178 *
	Post-test	11.54	2.19	-3.505**

GST: Glutathione S-transferase; \*p < 0.05; \*\*p < 0.01; §: Mean ± SD.

the values of the CRG before and after the training ( $p < 0.01$ , Table 3). When the values were compared according to the pre-test and post-test results of the different training methods, it was seen that there are statistically significant differences between the groups at  $p < 0.05$  and  $< 0.01$  levels (Table 4).

Many previous studies stated that a regular training program clearly increased the GST levels (Melikoğlu et al., 2008). In this study, it is seen that the interval training method shows parallel results to the findings of previous studies. Gül et al. (2006) examined the effects of endurance training and acute exhaustive exercises on the antioxidant defence mechanisms in rats. The training group exercised on a treadmill for 1.5 h per day, 5 days a week for 8 weeks. Both the training and the control groups then carried out exhaustive exercises. It was found that the GST level in the trained and untrained rats was not affected by the acute exhaustive exercise.

Kaldırımçı (2010) conducted a study that involved basketball training at 60 and 75% max  $\text{VO}_2$  on 3 days per week for 12 weeks. There was a significant increase in the GST values of the basketball players and, as a result, this increase raised the antioxidant activities of the physical exercise, which continued for long periods and caused adaptations in the antioxidant defense. In another study, it was observed that although acute exercise caused a decrease in the GST activities of the liver tissue of untrained rats, this decrease was also seen in trained rats. It was stated that the GST activity in the livers of the rats doing endurance exercise was not affected. Moreover, acute exercise-induced GST-reduction is prevented in trained rats and ROS production is probably reduced. These are partially responsible for the positive effects of the endurance training. Also, it was determined that there was a decrease in the fat tissue around the testicles of rats doing endurance training and this decrease might affect the body composition, depending on the use of fats rather than carbohydrates as the source of energy during the endurance training (Öztaşan et al., 2007).

In the present study carried out in a hot environment, it was seen that there was a statistically significant increase in the interval running group, a statistically significant decrease in the continuous running group and statistically significant differences in the comparison of the pre-test and post-test values between the groups.

Consequently, it can be said that the continuous running training in a hot environment affected the oxidative stress much more by increasing lipid peroxidation, while the interval running prevented the formation of free radicals by providing a positive increase in the glutathione S-transferase level.

In conclusion, it can be suggested that athletes and their trainers should generally apply the interval exercise program in order to reduce oxidative stress to a simpler state while doing endurance training in a hot environment.

## ACKNOWLEDGEMENTS

The author would like to avail this opportunity to thank Dr. Akar Karakoç and Dr. İlhami Gülçin, instructors at the Department of Chemistry, Faculty of Sciences, Atatürk University, for assisting in the biochemical analyses of the study.

## REFERENCES

- Ak T, Gülçin İ (2008). Antioxidant and radical scavenging properties of curcumin. *Chem. Biol. Interact.* 174: 27-37.
- ArasHisar Ş, Hisar O, Beydemir Ş, Gülçin İ, Yanık T (2004). Effect of vitamin E on carbonic anhydrase enzyme activity in rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vitro and in vivo. *Acta Vet. Hung.* 52: 413-422.
- Balaydın HT, Gülçin İ, Menzek A, Göksu S, Şahin E (2010). Synthesis and antioxidant properties of diphenylmethane derivative bromophenols including a natural product. *J. Enzym. Inhib. Med. Chem.* 25: 685-695.
- Beydemir Ş, Gülçin İ, Küfrevioğlu Öİ, Çiftçi M (2003). Glucose 6-phosphate dehydrogenase: In vitro and In vivo effects of dantrolene sodium. *Pol. J. Pharmacol.* 55: 787-792.
- Beydemir Ş, Gülçin İ (2004). Effect of melatonin on carbonic anhydrase from human erythrocyte in vitro and from rat erythrocyte in vivo. *J. Enzym. Inhib. Med. Chem.* 19: 193-197.
- Beydemir Ş, Gülçin İ, Hisar O, Küfrevioğlu Öİ, Yanık T (2005). Effect of melatonin on glucose-6-phosphate dehydrogenase from rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vitro and in vivo. *J. Appl. Anim. Res.* 28: 65-68.
- Clarkson PM, Thompson HS (2000). Antioxidants: What role do they play in physical exercise and health. *Am. J. Clin. Nutr.* 72: 637-645.
- Cheeseman KH, Slater TF (1993). An introduction to free radical biochemistry. *Brit Med. Bull.* 49: 481-493.
- Çoban TA, Beydemir Ş, Gülçin İ, Ekinci D (2008). The inhibitory effect of ethanol on carbonic anhydrase isoenzymes: in vivo and in vitro studies. *J. Enzym. Inhib. Med. Chem.* 23: 266-270.
- Çoban TA, Beydemir S, Gülçin İ, Ekinci D, Innocenti A, Vullo D, Supuran CT (2009). Sildenafil is a strong activator of mammalian carbonic anhydrase isoforms I-XIV. *Bioorg. Med. Chem.* 17: 5791-5795.
- Donald EP, William NV (1967). Studies on the quantitative characterization of erythrocyte glutathione peroxidase. *J. Lab. Clin. Med.* 70: 158-169.
- Durstine JL, Grandjean PW, Davis PG, Ferguson MA, Alderson NL, Dubose KD (2001). Blood lipid and lipoprotein adaptations to exercise: a quantitative analysis. *Sports Med.* 31: 1033-1062.
- Elosua R, Molina L, Fito M, Arquer A, Sanchez-Quesada JL, Covas MI, Ordóñez-Llanos J, Marrugat J (2003). Response of oxidative stress biomarkers to a 16-week aerobic physical activity program and to acute physical activity in healthy young men and women. *Atherosclerosis.* 167: 327-334.
- Finaud J, Scislawski V, Lac G, Durand D, Vidalin H, Robert A, Filair E

- (2006). Antioxidant status and oxidative stress in professional rugby players: evolution throughout a season. *Int. J. Sports Med.* 27: 87-93.
- Galassetti PR, Nemet D, Pescatello A, Rose-Gottron C, Larson J, Cooper DM (2006). Exercise, caloric restriction and systemic oxidative stress. *J. Investig. Med.* 54: 67-75.
- Goto C, Higashi Y, Kimura M, Noma K, Hara K, Nakagawa K, Kawamura M, Chayama K, Yoshizumi M, Nara I (2003). Effect Of Different intensities of exercise on endothelium dependent vasodilation on humans: Role of endothelium-dependent nitric oxide and oxidative stress. *Circulation.* 108: 530-535.
- Gül M, Demircan B, Taysi S, Öztasan N, Gümüştekin K, Siktar E, Polat MF, Akar S, Akcay F, Dane S (2006). Effects of endurance training and acute exhaustive exercise on antioxidant defense mechanisms in rat heart. *Comp. Biochem. Physiol. A. Mol. Integr. Physiol.* 143: 239-245.
- Gülçin İ, Oktay M, Kireççi E, Küfrevioğlu Öİ (2003a). Screening of antioxidant and antimicrobial activities of anise (*Pimpinella anisum* L.) seed extracts. *Food Chem.* 83: 371-382.
- Gülçin İ, Beydemir Ş, Alici HA, Elmastaş M, Büyükkuroğlu ME (2004b). In vitro antioxidant properties of morphine. *Pharmacol. Res.* 49: 59-66.
- Gülçin İ, Beydemir Ş, Büyükkuroğlu ME (2004d). In vitro and in vivo effects of dantrolene on carbonic anhydrase enzyme activities. *Biol. Pharm. Bull.* 27: 613-616.
- Gülçin İ, Mshvildadze V, Gepdiremen A, Elias R (2004a). Antioxidant activity of saponins isolated from ivy:  $\alpha$ -Hederin, hederasaponin-C, hederacolchiside-E and hederacolchiside F. *Planta Med.* 70: 561-563.
- Gülçin İ, Küfrevioğlu Öİ, Oktay M, Büyükkuroğlu ME (2004c). Antioxidant, antimicrobial, antiulcer and analgesic activities of nettle (*Urtica dioica* L.). *J. Ethnopharmacol.* 90: 205-215.
- Gülçin İ, Beydemir Ş, Şat İG, Küfrevioğlu Öİ (2005a). Evaluation of antioxidant activity of cornelian cherry (*Cornus mas* L.). *Acta Aliment. Hung.* 34: 193-202.
- Gülçin İ, Berashvili D, Gepdiremen A (2005b). Antiradical and antioxidant activity of total anthocyanins from *Perilla pankinensis* decne. *J. Ethnopharmacol.* 101: 287-293.
- Gülçin İ, Beydemir Ş, Hisar O (2005b). The effect of  $\alpha$ -tocopherol on the antioxidant enzymes activities and lipid peroxidation of rainbow trout (*Oncorhynchus mykiss*). *Acta Vet. Hung.* 53: 425-433.
- Gülçin İ, Mshvildadze V, Gepdiremen A, Elias R (2006a). Screening of antioxidant and antiradical activity of monodesmosides and crude extract from *Leontice smirnowii* Tuber. *Phytomedicine.* 13: 343-351.
- Gülçin İ, Elias R, Gepdiremen A, Boyer L (2006b). Antioxidant activity of lignans from fringe tree (*Chionanthus virginicus* L.). *Eur. Food Res. Technol.* 223: 759-767.
- Gülçin İ (2006a). Antioxidant activity of caffeic acid (3,4-dihydroxycinnamic acid). *Toxicology.* 217: 213-220.
- Gülçin İ (2006b). Antioxidant and antiradical activities of L-Carnitine. *Life Sci.* 78: 803-811.
- Gülçin İ (2007). Comparison of in vitro antioxidant and antiradical activities of L-tyrosine and L-Dopa. *Amino Acids.* 32: 431-438.
- Gülçin İ, Elmastas M, Aboul-Enein HY (2007a). Determination of antioxidant and radical scavenging activity of basil (*Ocimum basilicum*) assayed by different methodologies. *Phytother. Res.* 21: 354-361.
- Gülçin İ, Daştan A (2007). Synthesis of dimeric phenol derivatives and determination of in vitro antioxidant and radical scavenging activities. *J. Enzym. Inhib. Med. Chem.* 22: 685-695.
- Gülçin İ, Beydemir Ş, Çoban TA, Ekinci D (2008a). The inhibitory effect of dantrolene sodium and propofol on 6-phosphogluconate dehydrogenase from rat erythrocyte. *Fresen. Environ. Bull.* 17(9A): 1283-1287.
- Gülçin İ, Beydemir Ş, Hisar O, Köksal E, Reiter RJ (2009a). Melatonin administration increase antioxidant enzymes activities and reduce lipid peroxidation in the rainbow trout (*Oncorhynchus mykiss*, Walbaum) erythrocytes. *Turk. J. Vet. Anim. Sci.* 33: 241-245.
- Gülçin İ, Elias R, Gepdiremen A, Taoubi K, Köksal E (2009a). Antioxidant secoiridoids from fringe tree (*Chionanthus virginicus* L.). *Wood Sci. Technol.* 43: 195-212.
- Gülçin İ, Elias R, Gepdiremen A, Chea A, Topal F (2010c). Antioxidant activity of bisbenzylisoquinoline alkaloids from *Stephania rotunda*: Cepharanthine and fangchinoline. *J. Enzym. Inhib. Med. Chem.* 25: 44-53.
- Gülçin İ (2010). Antioxidant properties of resveratrol: A structure-activity insight. *Innov. Food Sci. Emerg.* 11: 210-218.
- Gülçin İ, Kirecci E, Akkemik E, Topal F, Hisar O (2010a). Antioxidant and antimicrobial activities of an aquatic plant: Duckweed (*Lemna minor* L.). *Turk. J. Biol.* 34:175-188.
- Gülçin İ, Bursal E, Şehitoğlu HM, Bilsel M, Gören AC (2010b). Polyphenol contents and antioxidant activity of lyophilized aqueous extract of propolis from Erzurum, Turkey. *Food Chem. Toxicol.* 48: 2227-2238.
- Halliwell B, Gutteridge JMC (1990). Role of free radicals and catalyticmetal ions in human disease: an overview. *Method Enzymol.* 186: 1-85.
- Hisar O, Beydemir Ş, Gülçin İ, ArasHisar Ş, Yanık T, Küfrevioğlu Öİ (2005a). The effect of melatonin hormone on carbonic anhydrase enzyme activity in rainbow trout (*Oncorhynchus mykiss*) erythrocytes in vitro and in vivo. *Turk. J. Vet. Anim. Sci.* 29: 841-845.
- Hisar O, Beydemir Ş, Gülçin İ, Küfrevioğlu Öİ, Supuran CT (2005b). Effect of low molecular weight plasma inhibitors of rainbow trout (*Oncorhynchus mykiss*) on human erythrocytes carbonic anhydrase-II isozyme activity in vitro and rat erythrocytes in vivo. *J. Enzym. Inhib. Med. Chem.* 20: 35-39.
- Kaldırımçı M (2010). The effect of training on G6FD and antioxidant activities. *Turk. J. Rheumatol.* 25: 34-36.
- Oktay M, Gülçin İ, Küfrevioğlu Öİ (2003). Determination of in vitro antioxidant activity of fennel (*Foeniculum vulgare*) seed extracts. *Lebensm Wissen Technol.* 36: 263-271.
- Osorio RAL, Christofani JS, Almeida VD, Picarro IC (2003). Swimming of pregnant rats at different water temperatures. *Comp. Biochem. Phys. Part A.* 135: 605-611.
- Özer MK (2006). Physical fitness (In Turkish). Second Edition. Ankara: Nobel Press.
- Öztaşan N, Timur H, Siktar E, Gümüştekin K, Akar S, Dane S, Gül M (2007). Effects of endurance training on gonadal fat pad and vertebral mass in rat. *Biol. Sports.* 24: 265-273.
- Öztürk Sarıkaya SB, Gülçin İ, Supurana CT (2010). Carbonic anhydrase inhibitors. Inhibition of human erythrocyte isozymes I and II with a series of phenolic acids. *Chem. Biol. Drug Des.* 75: 515-520.
- Peake J, Suzuki K (2004). Neutrophil activation, antioxidant supplements and exercise-induced oxidative stress exercise. *Exerc. Immunol. Rev.* 10: 129-141.
- Salo DC, Donovan CM, Davies KJA (1991). HSP70 and other possible heat shock or oxidative stressprotein are inducedin skeletal muscle, heart and liver during exercise. *Free Radical Biol. Med.* 11: 239-246.
- Sanchez-Quesada JL, Holms-Serradesanferm R, Serrat J, Grima JR, Gonzalez-Sastre J, Ordonez LJ (1995). Increase of ldl susceptibility to oxidation occurring after intense, long duration aerobic exercise. *Atherosclerosis.* 118: 297-305.
- Smith LL, Miles MP (2000). Exercise induced muscle injury and inflammation. Philadelphia. 401-411.
- Stein CJ, Colditz GA (2004). Modifiable risk factors for cancer. *Brit. J. Cancer.* 90: 299-303.
- Şentürk M, Gülçin İ, Çiftçi M, Küfrevioğlu Öİ (2008). Dantrolene inhibits human erythrocyte glutathione reductase. *Biol. Pharm. Bull.* 31: 2036-2039.
- Şentürk M, Gülçin İ, Daştan A, Küfrevioğlu Öİ, Supuran CT (2009). Carbonic anhydrase inhibitors. Inhibition of human erythrocyte isozymes I and II with a series of antioxidant phenols. *Bioorg. Med. Chem.* 17: 3207-3211.
- Şerbetçi Tohma H, Gülçin İ (2010). Antioxidant and radical scavenging activity of aerial parts and roots of Turkish liquorice (*Glycyrrhiza glabra* L.). *Int. J. Food Propert.* 13: 657-671.
- Talaz O, Gülçin İ, Göksu S, Saracoglu N (2009). Antioxidant activity of 5,10-dihydroindeno[1,2-b]indoles containing substituents on dihydroindeno part. *Bioorg. Med. Chem.* 17: 6583-6589.