

*Full Length Research Paper*

# Generation and analysis of cDNA library from lipopolysaccharide-stimulated gastropod abalone (*Haliotis diversicolor supertexta*) hemocytes

Liuji Wu<sup>1,3</sup>, Li Wang<sup>2</sup> and Xinzhong Wu<sup>3\*</sup>

<sup>1</sup>College of Agronomy, Henan Agricultural University, 63 Nongye Road, Zhengzhou 450002, P. R. China.

<sup>2</sup>Key Laboratory of Animal Genetics and Breeding of State Ethnic Affairs Commission and Ministry of Education, College of Life Science and Technology, Southwest University for Nationalities, Chengdu 610041, China.

<sup>3</sup>Laboratory of Marine Life Science and Technology, College of Animal Sciences, Zhejiang University, 268 Kaixuan Road, Hangzhou 310029, P. R. China.

Accepted 13 June, 2011

**A cDNA Library of lipopolysaccharide-stimulated gastropod abalone hemocytes was constructed using switching mechanism at 5' end of RNA transcript (SMART) approach. 300 random clones were selected and sequenced. 180 ESTs was gained. After assembling, we obtained 24 contigs and 119 singlets finally, some of which were immune-related genes. These immune-related genes include cytidine deaminase, ferritin, nonmuscle myosin essential light chain, cytochrome c oxidase subunit I, CD63 antigen-like protein and lysosomal-associated transmembrane protein. This study may contribute to the understanding of the immune mechanism of gastropod abalone *Haliotis diversicolor supertexta*.**

**Key words:** *Haliotis diversicolor supertexta*, cDNA Library, gene expression, immune mechanism.

## INTRODUCTION

Abalone is an important mollusc species for commercial production in the world including Australia, China, Japan, Korea, Mexico, South Africa, and the United States (Gordon and Cook, 2001). Since the late 1990s, the global industry of abalone has been gradually decreased due to diseases and environmental pollution (Gardner et al., 1995; Moore et al., 2000). Therefore, more investigation is required to aid in understanding the innate immune system of abalone for aquaculture development.

A critical step in any immune response is the recognition of invading organisms. It is mediated by many proteins, including pattern recognition proteins (PRPs), which recognize and bind to molecules present on the surface of microorganisms (Janeway, 1989). The binding of PRPs to lipopolysaccharide (LPS) triggers a series of


responses which lead to the activation of the host defence system (Lee and Soderhall, 2002). LPS is an essential cell wall component of Gram-negative bacteria.

LPS stimulated polyclonal proliferation of salmonid lymphocytes (Warr and Simon, 1983), respiratory burst and phagocytic activity of macrophages (Solem et al., 1995; Dalmo and Seljelid, 1995), and elicited secretion of IL-1 like compounds in channel catfish (Clem et al., 1985). LPS injected into red sea bream, *Pagrus major*, enhanced macrophage phagocytic activity (Salati et al., 1987). Similarly, LPS stimulated the production of macrophage activating factor in goldfish lymphocytes (Neumann et al., 1995).

Some reports showed that the effect of LPS has been studied in some aquatic animals, such as: Red sea bream ((Salati et al., 1987), trout (MacKenzie et al., 2003) and yellow grouper (Wang and Wu, 2007). However, there is still little information about the molecular response of abalone to LPS. In this study, a cDNA library was constructed to identify expressed genes in LPS-stimulated gastropod abalone hemocytes in an attempt to understand the molecular processes involved in bacteria

\*Corresponding author. E-mail: [wuxz@zju.edu.cn](mailto:wuxz@zju.edu.cn). Tel: 86-571-86971960. Fax: 86-571-86971960.

**Abbreviations:** PRPs, Pattern recognition proteins; LPS, lipopolysaccharide.



**Figure 1.** 1.2% agarose gel electrophoresis of the total RNA extracted from LPS-stimulated *H. diversicolor supertexta* hemocytes.

disease primary prevention.

## MATERIALS AND METHODS

### Animals, immune challenge and tissue collection

Healthy abalones (*Haliotis diversicolor supertexta*), 3 years of age, were collected from an abalone farm in Xiamen (Fujian, China) and kept in artificial seawater with a cycling system at 23°C (Jiang and Wu, 2007). Six abalones were challenged by injecting 50  $\mu$ L *Escherichia coli* lipopolysaccharide (LPS) (1 mg/mL diluted in sterile 0.9% sodium chloride). Hemocytes were collected at 12 h post injection by centrifugation (700 g, 10 min, 4°C).

### Total RNA isolation

Total RNA was isolated from LPS-stimulated abalone hemocytes using TRIzol reagent (Invitrogen, USA) according to the manufacturer's instructions. RNA integrity was checked using 1% formaldehyde agarose gel. The quantity and quality of RNA were determined by absorbance at A260 and at A260/280 ratios, respectively, using a Perkin-Elmer Lambda 3B spectrophotometer.

### Construction of cDNA library

Abalone cDNA library was constructed by a SMART<sup>TM</sup> cDNA library construction kit (Clontech, USA) according to the manufacturer's instruction with a few modifications. Briefly, the isolated RNA was reverse transcribed to synthesize first strand cDNA at 42°C for 1 h. Then first strand cDNA was mixed with PCR primer and other PCR reagents at 95°C for 30 s followed by 20 cycles of 95°C for 10 s and 68°C for 6 min to synthesize double-strand DNA. After digestion and purification, double-strand DNA was ligated to TriplEx2 vectors. Then the resulting ligation reaction was packed using the Lambda DNA Packaging System (Promega, USA) following the manufacturer's specifications. Screening of the abalone cDNA library was based on polymerase chain reaction (PCR) with the pTriplEx2 sequencing primers. The forward primer is 5'-CTCCGAGATCTGGACGAGC-3', and the reverse primer is 5'-

TAATACGACTCACTATAGGGC-3'. Positive plasmid clones were grown in liquid cultures and induced to a high copy number for direct sequencing using an ABI PRISM 3730 automated sequencer (Applied Biosystems).

### Sequence analysis

DNA traces were assembled with PHRED/PHRAP/CONSED package ([www.phrap.org](http://www.phrap.org)). Vector contamination sequences were removed with the program Cross\_match. Low-quality bases (quality score <20) were trimmed from both ends of sequences using a PHRED. EST sequences corresponding to the same transcript were then assembled with the program PHRAP. Sequence similarity analysis was performed by using the BLASTX program (Altschul et al., 1997) with threshold E-value  $\leq 1e-5$ , against the non-redundant database provided by National Center for Biotechnology Information (<http://www.ncbi.nlm.nih.gov/>). Classification of annotated sequence was searched by Gene Ontology (<http://www.godatabase.org/>).

## RESULTS


### Construction of abalone cDNA library

In order to analyze gene expressed in LPS-stimulated abalone hemocytes and to facilitate functional analysis of cDNA clones, a SMART cDNA library was constructed. Total RNA with high quality is shown in Figure 1. The primary titer of the library was  $2 \times 10^8$  pfu/ml, and the recombinant phage in the library was 90%. The quality of cDNA library was examined by sequencing of the inserted cDNA clones (Julio and Miguel, 2002; Wakimoto, 2000). The insert size was distributed in a range from 200 to 2000 bp which reflected the size distribution of the first-strand cDNA was convergence (Figure 2).

### Sequence analysis

From cloned PCR products, 300 randomly-selected clones containing inserts were sequenced. After re-moving the vector sequence and the poor-quality sequence (with trim cutoff 0.05), 180 qualified ESTs were grouped into 24 contigs, and another 119 singlets with an average length of approximately 400 bp. They were submitted to GenBank database with accession numbers HS040369-HS040662. Tentative annotations were performed by using BLASTX and results were manually validated. These sequences were used to analyze gene expression profile from the LPS-stimulated abalone hemocytes.

Fifty seven (57) sequences were assigned to defined annotations, whereas 32 sequences showed similarity with unclassified genes and 54 sequences remained unknown. These sequences according to the functions were involved in immune response, protein biosynthesis, metabolism, biological process and cellular component, while the remainder belonged to none of the categories mentioned above (Table 1).



**Figure 2.** PCR identification of randomly selected clones from cDNA library. Lane1-35: randomly selected clones; M: marker DL2000.

## DISCUSSION

To date, there were several methods used for constructing cDNA library, for example; Oligo-Capping (Maruyama and Sugano, 1994), SMART (Chenchik et al., 1996), Cap-Select (Schmidt et al., 1999) and CAP-jumping (Fimove et al., 2001). Among these methods, the SMART<sup>TM</sup> cDNA Library Construction Kit provides a way for producing high-quality, full-length cDNA libraries from nanograms of total or poly A+ RNA. All commonly used cDNA synthesis methods rely on the ability of reverse transcriptase to transcribe mRNA into single-stranded (ss) DNA in the first strand reaction. In some cases, RT terminates before transcribing the complete mRNA sequence. This is particularly true for long mRNAs, especially if the first strand synthesis is primed with oligo (dT) primers only or if the mRNA contains abundant secondary structures. In addition, conventional cDNA cloning procedures use the T4 DNA polymerase to generate blunt cDNA ends after second strand synthesis. As a result, under-represented 5' ends of genes in cDNA populations tend to be 5–30 nucleotides shorter than the original mRNA.

The SMART protocols are designed to preferentially

enrich for full-length cDNAs, while eliminating T4 DNA polymerase and adaptor ligation. SMART libraries are proven to contain a higher percentage of full-length clones than libraries constructed by conventional methods or other full-length cDNA synthesis protocols (Sughura et al., 2001; Kato et al., 1994). Thus, clones isolated from SMART cDNA libraries contain sequences corresponding to the complete 5' untranslated region of the mRNA. SMART cDNA libraries can facilitate preliminary mapping of transcription start sites due to the high percentage of full length clones. However, SMART cDNA libraries may not be suitable for immunoscreening for certain proteins.

The major characteristic of cDNA construction by the SMART method could improve the ratio of full length cDNA sequences. It is important to identify a new gene by constructing a cDNA library and EST sequencing to get full-length cDNA from a special tissue, even if that EST is a fragment of one gene. The SMART method has been used in aquatic organism, such as *Tegillarca granosa* (Bao and Lin, 2010) and *Miichthys miiuy* (Xu et al., 2010). Using this approach, we identified 24 contigs and 119 singlets finally. All of the genes are reported for the first time from gastropod abalone, which are involved

**Table 1.** Key unigenes shared significant similarities with known genes of other species.

Length	Score	E value	Identity	Subject Name	Annotation
149	303	3.00E-80	100	gb ABW90692.1	cytidine deaminase
151	187	2.00E-45	68	ref XP_002066344.1	GK18243
231	149	6.00E-68	81	gb ABY87393.1	ATP synthase F0 subunit 6
142	154	2.00E-35	52	gb ABY87349.1	profilin
128	161	2.00E-37	69	ref NP_001080724.1	NHP2 non-histone chromosome protein 2-like 1
239	169	5.00E-40	41	ref XP_002588730.1	hypothetical protein BRAFLDRAFT_100190
206	281	9.00E-74	72	gb ABY87388.1	calmodulin-dependent protein kinase
171	344	1.00E-92	100	gb ABY87353.1	ferritin
1040	221	4.00E-56	44	ref XP_002719999.1	connector enhancer of kinase suppressor of Ras 2 isoform 1
147	208	1.00E-51	68	ref XP_002407055.1	nonmuscle myosin essential light chain
189	304	1.00E-80	85	gb ABO26639.1	transgelin
142	290	3.00E-76	97	gb ABY87349.1	profilin
205	441	1.00E-121	100	gb ABY87355.1	stanniocalcin-like protein
76	77.8	3.00E-12	56	emb CAX70431.1	LIM, zinc-binding, domain-containing protein
257	491	1.00E-137	99	gb ABY87386.1	60S acidic ribosomal protein P0
246	388	1.00E-106	74	ref XP_002736568.1	PREDICTED: exosome component 4-like
535	208	7.00E-52	65	ref NP_001122125.1	cortactin
106	116	4.00E-24	52	ref XP_001969325.1	GG10043 [ <i>Drosophila erecta</i> ]
72	87.4	3.00E-15	100	gb ABY87379.1	omega class glutathione S-transferase
326	70.9	2.00E-22	100	ref ZP_06718036.1	hypothetical protein CUS_0263
127	135	6.00E-39	100	gb ABV25015.1	beta-galactosidase a-peptide
103	171	2.00E-40	86	ref ZP_04858619.1	ribosomal protein S10
126	120	3.00E-25	53	ref XP_784178.1	PREDICTED: hypothetical protein isoform
580	478	1.00E-133	80	gb ACB73226.1	NADH dehydrogenase subunit 5
140	114	2.00E-23	45	ref XP_970004.1	similar to mitochondrial NADH
600	103	6.00E-20	25	ref YP_519561.1	hypothetical protein DSY3328
426	277	2.00E-72	83	ref ZP_05633094.1	adenylosuccinate synthetase
147	296	2.00E-78	91	ref XP_001120139.1	similar to Ubiquitin-conjugating enzyme E2-17
513	354	1.00E-95	93	gb ACL99801.1	cytochrome c oxidase subunit I
533	79	4.00E-13	56	ref XP_002734661.1	PREDICTED: sialin-like
174	172	9.00E-41	53	ref XP_002424313.1	Nucleoside diphosphate kinase, putative
174	172	9.00E-41	53	ref XP_002424313.1	Nucleoside diphosphate kinase, putative
192	350	2.00E-94	91	gb ABY87354.1	calmodulin 2
449	212	8.00E-53	83	gb AAT44354.1	isocitrate dehydrogenase
483	204	2.00E-50	62	ref XP_001517928.1	PREDICTED: similar to PLC alpha, partial
122	190	3.00E-51	88	gb ABY87409.1	CD63 antigen-like protein

Table 1. continues

176	226	4.00E-57	70	ref NP_001090206.1	ribosomal protein L18a
832	48.5	4.00E-15	43	ref XP_002933957.1	PREDICTED: hypothetical protein
107	49.3	3.00E-10	56	ref ZP_02207458.1	hypothetical protein COPEUT_02273
140	234	1.00E-59	98	gb ABV44723.1	60S ribosomal protein L23-like protein
314	500	1.00E-139	84	gb ACL99797.1	NADH dehydrogenase subunit 1
486	99.4	8.00E-19	50	gb AAF91388.1	AF263243_1 SocE
520	511	1.00E-143	79	ref XP_001603746.1	similar to nucleolar KKE/D repeat protein
1222	70.1	5.00E-10	38	gb ABN58714.1	pol-like protein
176	311	9.00E-83	83	emb CAD91425.1	actin related protein 2/3 complex,
257	397	1.00E-108	77	gb ABW23163.1	ribosomal protein rpl7a
192	57.4	3.00E-06	100	gb ABY87354.1	calmodulin 2
786	144	2.00E-32	40	ref XP_002733949.1	PREDICTED: CG1104-like
307	145	7.00E-33	39	ref XP_002931362.1	PREDICTED: GTPase IMAP family member 4-like
341	81.3	2.00E-13	79	ref XP_002639536.1	<i>C. briggsae</i> CBR-FBP-1 protein
326	69.3	8.00E-22	100	ref ZP_06718036.1	hypothetical protein CUS_0263
245	165	1.00E-38	38	ref NP_001133826.1	Lysosomal-associated transmembrane protein 4A
235	464	1.00E-128	99	gb ABO26645.1	proteasome alpha type 2
341	80.5	4.00E-13	34	ref XP_002738783.1	PREDICTED: transcription factor AP-4
113	218	8.00E-55	85	gb EFA01469.1	hypothetical protein TcasGA2_TC007016
340	145	1.00E-32	42	gb AAG61258.1	CCAAT/enhancer binding protein
220	389	1.00E-106	88	gb ABO26631.1	Ran-1-prov protein
163	106	5.00E-21	50	ref XP_002734957.1	PREDICTED: <i>C. briggsae</i> CBR-GLRX-10 protein-like

in immune reaction, biological process, metabolism, protein binding and so on. Of the 57 known genes, 6 immune-relevant genes were identified and isolated: cytidine deaminase, ferritin, nonmuscle myosin essential light chain, cytochrome c oxidase subunit I, CD63 antigen-like protein, and lysosomal-associated transmembrane protein. These immune-relevant genes should be useful for analyzing gene function during disease defense and for developing molecular markers related to disease-resistance. Possible reasons for the low number of immune-

related genes uncovered in this study were as follows: the need to obtain more transcripts from fully differentiated abalone hemocytes treated with different dose of LPS and sampled from other times, and the need for additional sequencing of the library.

In summary, a cDNA Library was constructed to isolate and identify expressed genes in LPS-treated gastropod abalone hemocytes. 24 contigs and 119 singlets were obtained finally; 6 of which appeared to be important immune-relevant genes. Further investigations will be required to deter-

mine expression mechanisms and functions of these genes and processes involved in the defense mechanism of the gastropod abalone - *H. diversicolor supertexta*.

#### ACKNOWLEDGEMENTS

This work was supported by the National Natural Science Foundation of China (NFSC 30671619). We thank Dr. Haifeng Pan for his helpful suggestions.

## REFERENCES

- Altschul SF, Madden TL, Schaffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ (1997). Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucleic Acid Res.* 25(17): 3389-3402.
- Bao YB, Lin ZH (2010). Generation, annotation, and analysis of ESTs from hemocyte of the bloody clam, *Tegillarca granosa*. *Fish Shellfish Immunol.* 29(5): 740-746.
- Chenchik A, Moqadam F, Siebert P (1996). *RNA: Isolation, Analysis and Synthesis*. New York: Wiley-Liss, pp. 273-321.
- Clem LW, Sizemore RC, Ellsaesser CF, Miller NW (1985). Monocytes as accessory cells in fish immune responses. *Dev. Comp. Immunol.* 9(4): 803-809.
- Dalmo RA, Seljelid R (1995). The immunomodulatory effect of LPS, laminaran and sulfated laminaran [ $\beta(1, 3)$ -D-glucan] on Atlantic salmon, *Salmo salar* L, macrophages *in vitro*. *J. Fish. Dis.*, 18(2): 175-185.
- Fimove VA, Chakhmakhcheva OG, Archdeacon J, Fernande JM, Fedorkin ON, Dorokhov YL, Atabekov JG (2001). Detection of the 5'-cap structure of messenger RNAs with the use of the CAP-jumping approach. *Nucleic Acids Res.* 29(22): 4751-4759.
- Gardner GR, Harshbarger JC, Lake JL, Sawyer TK, Price KL, Stephenson MD, Haaker PL, Togstad HA (1995). Association of prokaryotes with symptomatic appearance of withering syndrome in black abalone *Haliotis cracherodii*. *J. Invertebr. Pathol.* 66(2): 111-115.
- Gordon HR, Cook PA (2001). World abalone supply, markets and pricing: historical, current and future. *J. Shellfish. Res.* 20: 567-570.
- Janeway CA (1989). Approaching the asymptote, evolution and revolution in immunology. *Cold Spring Harb. Symp. Quant. Biol.* 54: 1-13.
- Jiang YS, Wu XZ (2007). Characterization of a Rel/NF- $\kappa$ B homologue in a gastropod abalone, *Haliotis diversicolor*. *Dev. Comp. Immunol.* 31(2): 121-131.
- Julio E, Miguel CP (2002). Bioinformatics and reanalysis of subtracted expressed sequence tags from the human ciliary body: identification of novel biological functions. *Mol. Vision*, 8: 315-332.
- Kato S, Sekine S, Oh SW, Kim NS, Umezawa Y, Abe N, Yokoyama-Kobayashi M, Aoki T (1994). Construction of a human full-length cDNA bank. *Gene*, 150(2): 243-250.
- Lee SY, Soderhall K (2002). Early events in crustacean innate immunity. *Fish Shellfish Immunol.* 12(5): 421-437.
- MacKenzie S, Planas J, Goetz F (2003). LPS-stimulated expression of a tumor necrosis factor-alpha mRNA in primary trout monocytes and *in vitro* differentiated macrophages. *Dev. Comp. Immunol.* 27(5): 393-400.
- Maruyama K, Sugano S (1994). Oligo-capping: a simple method to replace the cap structure of eukaryotic mRNAs with oligoribonucleotides. *Gene*, 138(1-2): 171-174.
- Moore JD, Robbins TT, Friedman CS (2000). Withering syndrome in farmed red abalone, *Haliotis rufescens*. *J. Aquat. Anim. Health*, 12(1): 26-34.
- Neumann NF, Fagan D, Belosevic M (1995). Macrophage activating factor s secreted by mitogen stimulated goldfish kidney leucocytes synergies with bacterial lipopolysaccharide to induce nitric oxide production in teleost macrophages. *Dev. Comp. Immunol.* 19(6): 475-482.
- Salati F, Hamaguchi M, Kusuda R (1987). Immune response of red sea bream to *Edwardsiella tarda* antigens. *Fish Pathol.* 22(2): 291-304.
- Schmidt WM, Mueller MW (1999). CapSelect: A highly sensitive method for 5-CAP-dependent enrichment of full-length cDNA in PCR-mediated analysis of mRNAs. *Nucleic Acids Res.* 27(21): 31-31.
- Solem ST, Jørgensen JB, Robertsen B (1995). Stimulation of respiratory burst and phagocytic activity in Atlantic salmon (*Salmo salar* L.) macrophages by lipopolysaccharide. *Fish Shellfish Immunol.* 5(7): 475-491.
- Sughura Y, Carninci P, Iton M, Shibata K, Konno H, Endo T, Muramatsu M, Hayashizaki Y (2001). Comparative evaluation of 5-end-sequence quality of clones in CAP trapper and other full length cDNA libraries. *Gene*, 263(2): 93-102.
- Wakimoto BT (2000). Doubling the rewards: testis ESTs for *Drosophila* gene discovery and spermatogenesis expression profile analysis. *Genome Res.* 10(12): 1841-1842.
- Wang L, Wu XZ (2007). Identification of differentially expressed genes in lipopolysaccharide-stimulated yellow grouper *Epinephelus awoara* spleen. *Fish Shellfish Immunol.* 23(2): 354-363.
- Warr GW, Simon RC (1983). The mitogen response potential of lymphocytes from the rainbow trout (*Salmo gairdneri*) re-examined. *Dev. Comp. Immunol.* 7(2): 379-384.
- Xu TG, Meng FX, Sun YN, Shi G, Wang RX (2010). Identification of immune genes of the miiuy croaker (*Miichthys miiuy*) by sequencing and bioinformatic analysis of ESTs. *Fish Shellfish Immunol.* 29(6): 1099-1105.