

Full Length Research Paper

Screening of autochthonous *Lactobacillus* species from Algerian raw goats' milk for the production of bacteriocin-like compounds against *Staphylococcus aureus*

Mami Anas*, Boumehira Ali Zinedine, Hamedi Amine Rizk, Henni Jamal Eddine and Kihal Mebrouk

Laboratory of Applied Microbiology, Department of Biology, Faculty of Science, Oran University, BP 16, Es-senia, 31100, Oran, Algeria.

Accepted 27 January, 2012

Lactic acid bacteria play a key role in maintaining the balance of normal gastro-intestinal microflora. Food contamination by *Staphylococcus aureus* is a major problem for consumer's health in Algeria, especially during the summer period. The use of bacterial interactions is a new way to limit the pathogenic germs growth. Detection of antimicrobial substances produced by lactic acid bacteria against the undesirable germs is the objective of this work. Microbiological and biochemical methods were used to identify lactic acid bacteria having an antimicrobial activity. The 02 isolates of lactic acid bacteria obtained from raw goats' milk in western Algeria areas were identified as: *Lactobacillus rhamnosus* and *Lactobacillus plantarum*. The interactions study revealed that two lactobacilli species: *L. plantarum* (58) and *L. plantarum* (68) are able to inhibit *S. aureus*' growth. In mixed culture after 24 h, *L. plantarum* reduces the growth of *S. aureus* by 1.6 log and this latter bacteria was not found after 72 h. The various tests used revealed the proteinic nature of the substance, which was responsible of the growth inhibition of *S. aureus*. The ecological adaptation and growth characteristics of cultures of *L. plantarum* in food products will determine their effectiveness as biocontrol agent in dairy foods.

Key words: Raw goat's milk, lactic acid bacteria, *Lactobacillus plantarum*, interaction, bacteriocin, *Staphylococcus*, mixed culture.

INTRODUCTION

Control of both pathogenic and spoilage microbe in a variety of foods is important to guarantee food quality and safety. Recently, biopreservation has become a topic of interest (Kabuki, et al., 2007; Keihei et al., 2011). This technique is used as an alternative to chemical additives for increasing self-life storage and enhancing safety of food by using natural microflora and their antimicrobial products (Stiles, 1996). Lactic acid bacteria are believed to be safe because they have been long established as the normal flora in fermented food; thus, they have great potential for use in biopreservation. The preserving

effects of lactic acid bacteria are due to the production of antimicrobial agents such as organic acids, hydrogen peroxide and bacteriocin or related substances (Desmazeaud and Cogan, 1996; Cocolin et al., 2007). Bacteriocins are proteinaceous compounds that mainly inhibit closely related species (Klaenhammer, 1993). Some bacteriocins have been shown to possess the ability to inhibit the actions of unrelated genera such as Clostridia, Listeria, enteropathogenic bacteria and Gram-negative bacteria. For these reasons, bacteriocins are promising candidates for biopreservation of food (Cleveland, 2001). Several *Lactobacillus* strains are important dairy culture starter and are used for the manufacturing of fermented food (Fitzsimmons et al., 1999; Badis et al., 2004; Pringsulaka et al., 2011b).

The discovery of bacteriocins gave a new way for food

*Corresponding author. E-mail: anas.mami@yahoo.fr. Tel: 00213664260403. Fax: 0021341582540

development in better hygienic quality (Fitzsimmons et al., 1999; Badis et al., 2004; Pringsulaka et al., 2011b). In recent years, there have been many reports on bacteriocins that are produced by lactic acid bacteria. However, most reports deal with bacteriocins that are produced by various Lactococci, Pediococcus, Leuconostoc, Enterococci and Lactobacilli (Ennahar et al., 2000; Mc Auliffe et al., 2001; Moll et al., 1999). The search for new strains of lactic acid bacteria that produce antimicrobial substances is a universal objective for the creation of new cultures starter with a high biosafety for fermented food. The inhibition of pathogenic bacteria such as *Staphylococcus aureus* by lactic microflora was reported by Heikkila and Saris (2003). The technological characterization of the lactic acid bacteria leads to the development of well defined bacterial strains with specific characteristics. The latter gradually replace the non-definite and non-defined mixtures of starters traditionally used in dairy industry (Crow et al., 1993; Papagianni and Anastasiadou, 2009).

Lately, the interest for the use of the bacteriocins or strains of lactic acid bacteria for applications as bio-preservatives in order to avoid the side effects of chemical preservatives has resulted to many research tasks (Schillinger and Lücke, 1989; Budde et al., 2003; Jacobsen et al., 2003; Vermeiren et al., 2004; Cocolin et al., 2007; Alvarez-Martin et al., 2008; Pringsulaka et al., 2011a). Several bacteriocin producing lactic acid bacteria offer potential applications in food preservation. The use of bacteriocins in the food industry can help to reduce the addition of chemical preservatives as well as the intensity of heat treatments, thus resulting in foods that are more naturally preserved and richer in organoleptic and nutritional properties (Galvez et al., 2007). The aim of this work is to isolate lactic acid bacteria that produce antimicrobial substances belonging to bacteriocin-type, which are able to inhibit the bacteria causing food poisoning.

MATERIALS AND METHODS

Bacterial strains and growth conditions

The species of *Lactobacillus* were obtained from the collection of the laboratory of applied microbiology, Department of biology, Faculty of Science, University of Oran, Algeria. The three pathogenic bacteria responsible of food toxic infections (*S. aureus* ATCC 25923) were obtained from the collection of Medical Analysis Laboratory of the University- Hospital centre of Oran. The species of lactobacilli were cultivated in liquid or solid deMan, Rogosa and Sharpe (MRS) medium with pH 5.4, and then incubated at 30°C for 48 h (de Man et al., 1960). The selective enumeration of *S. aureus* was carried out on Chapman medium at 37°C. The other bacteria, *Escherichia coli* and *Bacillus* sp. were grown on Muller-Hinton medium and incubated at 37°C. The media used during this work were either liquid, solid (1.5% agar p/v) or soft agar medium (0.7% agar). The skimmed milk medium (11% p/v) was sterilized at 110°C for 10 min and all other media were sterilized at 121°C for 20 min (Kihal et al., 2006).

The isolated *Lactobacillus* strains were selected as bacteriocin producers because of their broad antimicrobial activity, and

subjected to phenotypic identification. Cell morphology and Gram-staining reaction were examined by light microscopy. Test for catalase activity and fermentation of different sugars were also tested as described by Badis et al. (2004).

Hemolytic activity

For testing hemolytic activity, fresh cultures of isolated strains were streaked on blood agar, containing 5% (w/v) sheep blood, and incubated for 48 h at 30°C. Blood agar plates were examined for signs of hemolysis. This test was performed in the hospital laboratory (Ghraiiri et al., 2008).

Antimicrobial activity detection and assay

The production of antimicrobial substance by *Lactobacillus* sp. was detected by the deferred antagonism assay as described by Barefoot et al. (1983). An overnight culture of *Lactobacillus plantarum* was spotted onto the surface of an MRS plate and incubated for 16 h at 30°C to allow colonies to develop. Approximately 10^7 cfu/ml of indicator strain were inoculated into 10 ml of MRS agar and poured over the plate on which the producer was grown. After anaerobic incubation for 24 h at 30°C, the bacterial lawns were examined for zones of inhibition surrounding the producer colonies. Inhibition was recorded as positive if the width of the clear zone around the colonies of the producer was 2 mm or larger (Tahara and Kanatani, 1996). Bacteriocin-like activity was assayed by the agar-well diffusion method of Tagg and McGiven (1971). Portions (100 µL) of serial twofold dilutions of culture filtrates of *L. plantarum* strain were placed into each 0.5 cm diameter well of a plate, which was inoculated with approximately 10^6 cfu/ml of a log-phase culture of *L. plantarum* and after a 24 h incubation at 30°C, plates were examined for inhibition on the indicator lawn. For the activity spectrum determination of bacteriocin-like produced by *L. plantarum* pathogenic bacteria *S. aureus* ATCC 25923 were used.

Preparation of crude bacteriocin: Sensitivity to enzymes, pH and heat treatment

MRS broth was used for the preparation of crude bacteriocin from the culture of *L. plantarum*. After incubation at 30°C for 18 h, culture supernatants were obtained by centrifugation at 8000 rpm/min for 10 min at 4°C. Crude bacteriocin was concentrated by precipitation with 5% tricarboxylic acid (TCA) and stored at 4°C. As for the sensitivity to enzymes, pH and heat treatment, the neutralized MRS culture supernatant was tested for susceptibility to proteolytic enzymes. Samples were incubated at 30°C for 1 h in the presence of trypsin, alfa-chymotrypsine and catalase (Jamuna et al., 2005; Kabuki et al., 2007; Tahara and Kanatani, 1996).

After the treatment, the reaction mixture was heated for 10 min in boiling water (100°C) for enzyme inactivation and then bacteriocin activity was measured. To determine the pH stability of bacteriocin, pH values of the culture supernatants were adjusted within the range of five to seven by HCl or NaOH, and each sample was held for 1 h at 30°C. After incubation, the pH of each sample solution was adjusted to 6.5 by adding HCl or NaOH and the bacteriocin activity was measured as described above. To examine thermal stability of bacteriocin, crude bacteriocin was treated at the various temperatures mentioned above. After the treatment, the samples were rapidly cooled and the bacteriocin activity was assayed (Tahara and Kanatani, 1996).

Assay for antimicrobial activity

The search for possible production of inhibiting substances by the

isolated bacteria was carried out according to two methods: The first is the direct method of Tagg and Mc Given (1971), Fleming et al. (1975) and Tahara and Kanatani (1996), which consists of cultivating the two strains in the same medium with double-layer. The inhibitory activity spectrum was obtained using the agar spot test (Casalta et al., 1995) against the other strains. For this procedure, 5 μ L aliquots from an overnight culture of *Lactobacillus* sp. strain grown in MRS broth being spotted onto buffered MRS agar plates (1.5% agar) and incubated for 24 h were used. Subsequently the plates were then overlaid with 6 ml of soft agar medium (Muller-Hinton) seeded with actively growing cells of the test organisms (or pathogenic strain (*S. aureus*) after solidification and then incubation at 30°C. The antimicrobial activity of *Lactobacillus* sp. and the sensitivity of the pathogenic strain in question were evaluated by checking for clear zones around the spots (Tahara and Kanatani, 1996; Kaban and Kaya, 2006; Guessas et al., 2006).

The second is the indirect method of Barefoot et al. (1983) and Schillinger and Lücke (1989), which makes it possible to put in contact the supernatant of lactic acid bacteria strains that produce antimicrobial substances with the test strains. Producing strains of inhibiting substances were cultivated in liquid medium MRS and incubated for 18 h. After growth, the culture was centrifugalized at 8000 rpm/min during 10 min and the supernatant was stored at 4°C. The supernatant fluid was filtered through a syringe filter with a pore size of 0.22 μ M (Millipore Corporation, Bedford, USA) and adjusted to pH 6.0 with sterilized NaOH (2 M), so as to rule out inhibition through the production of organic acids. This supernatant was placed into the wells and the medium was inoculated by the test strains. These wells received 100 μ L of tested supernatant strains and were then incubated for 24 h. Inhibition of growth was determined by an area of inhibition surrounding each agar well (Miteva et al., 1998; Guessas, 2007).

Determination of the spectrum activity

The direct confrontation method of Shillinger and Lücke (1989) and Sookkhee et al. (2001) was used for the determination of the spectrum activity of antimicrobial substances strains production. The selected producing species belong to *L. plantarum* (58 and 68), while the test species belong to *S. aureus*.

Characterization of the nature of the inhibiting agent

The antimicrobial activity of *L. plantarum* can be caused by several factors such as acidity, hydrogen peroxide, phages and bacteriocins. To determine whether the inhibitory substances produced by the bacteria were proteinaceous and thus could be designated as bacteriocins, sensitivity to proteolytic enzymes (trypsin and α -chymotrypsin) was assessed in assays as described by Vaughan et al. (1992, 1994). Blank experiments were also performed using also no enzyme or inactivated enzyme. Gram positive *S. aureus* was used as the indicator strain in the experiments (Ghraiiri et al., 2008).

Kinetics of growth and acidification

Associative growth of *S. aureus* and the bacteriocin producer strain *L. plantarum* (58) was performed in sterilized skim milk at 30°C. The acidity evaluation of the pure strain was carried out by titration and pH metrical measurement (Campos et al., 2006). Each strain was inoculated in 10 ml of sterile skim milk (10% p/v). The pre-culture of *L. plantarum* was prepared by incubation at 30°C until coagulation. An amount of 3% of pre-culture inoculated in 100 ml of skim milk and was immediately homogenized. 10 ml of the mixed

culture was distributed in sterile tubes. The growth kinetics and acidification were carried out simultaneously within the regular lengths of time intervals.

The pathogenic strain of *S. aureus* (ATCC.25923) which was inoculated in skim milk at 2% which gives approximately 10³ cfu/ml, in pure and mixed culture with *L. plantarum* was used as test. The antimicrobial effect of the latter strain against *S. aureus* was measured by microbiological techniques as described by Kaban and Kaya (2006). Serial decimal dilutions were prepared in sterile physiological saline water and 0.1 ml samples of appropriate dilutions were spread in duplicate on selective agar plates. The number of *S. aureus* and *L. plantarum* was determined in Chapman and MRS agar media, respectively (Ananou et al., 2007).

Measurement of acid production

A deduction of 10 ml of the culture was transferred in a conical flask of 100 ml and five drops of phenolphthalein indicator (2 mg/ml in ethanol 60°) were added. The acidity was neutralized by NaOH 1/9 N until the appearance of a persistent pink color. The volume of the titrating solution was measured in order to indicate the production of acidity which was estimated in Dornic degrees (Kihal et al., 1996; Moulay et al., 2006).

Growth kinetics measurement in pure and mixed culture in milk

The enumeration of *S. aureus* was carried out on Chapman medium. Only plates that contained 30 to 300 colonies were taken into account. The enumeration in mixed culture was done by sowing 0.1 ml of serial dilutions in two selective acidified MRS media for *Lactobacillus* and Chapman for *S. aureus* (Guessas et al., 2006; Kaban and Kaya, 2006; Otero and Macias, 2006).

Effect of the crude supernatant of *L. plantarum*

The crude supernatant of *L. plantarum* (58) was tested against the growth of *S. aureus* in milk. The crude supernatant was obtained by the centrifugation of a culture of 18 h in MRS pH 6.8 with 8000 rpm/min for 10 min. Protein concentration of the culture supernatant was determined by the micromethod of Bradford (1976) using bovine serum albumin fraction as standard. The supernatant was heated at 100°C. The residual activity of the crude supernatant was immediately tested after freezing and neutralization by NaOH. *S. aureus* was tested for determining the minimal inhibiting concentrations of this crude supernatant. After incubation with various dilutions, the growth was estimated by reading the optical density at 670 nm (Tahara et al., 1996; Blay et al., 2007; Pal and Ramana, 2010).

RESULTS AND DISCUSSION

A total of 128 strains of *Lactobacillus* were isolated from raw goat's milk. The strains which produce antimicrobial substances were detected by confrontation on solid culture medium. From the 128 isolates only two strains showed inhibiting activity. These latter strains were identified to species level by microbiological and biochemical methods as described by Stiles et al. (1997), Klein et al. (1998) and Carr et al. (2002).

Table 1. First identification characters of *Lactobacilli* isolates from raw goat's milk.

Character	<i>Lactobacilli</i> strain	
	58	68
Form	Rod	Rod
Gram	+	+
Catalase	-	-
Fermentation ribose	+	+
CO ₂ from glucose	-	-
CO ₂ from gluconate	+	+
Arginine Hydrolysis	-	-
Growth at 15/45	+/+	+/+
Group	Lactobacillus Group II, facultative heterofermenter (Streptobacteria)	Lactobacillus Group II, facultative heterofermenter (Streptobacteria)

Characterization of the isolates

The strains retained gave small colonies of approximately 1 mm diameter, lenticular with a white or milky colour, smooth surface and regular circular circumference were observed on solid medium. The microscopic examination revealed that the tested strains were Gram positive, with a cellular rod form associated in pairs or in chains (Table 1). Table 2 shows the characteristics for a better identification of the species used in this study. These data guide us to classify the isolated bacteria to the genus level according to their cellular morphologies and their association mode and the type of Gram staining (Joffin and Leyral, 1996). On the basis of microbiological (Table 1), physiological and biochemical (Table 2) analysis results, we were able to establish the percentage of reliability of each strain in comparison with previous reports (Stiles et al., 1998; Klein et al., 1998; Carr et al., 2002).

Hemolytic activity

In eight strains of *Lactobacillus* isolated from the raw goat's milk and used in this work, no hemolysis activity was observed. However, absence of hemolytic activity should be a selection criterion for (bacteriocin-producing) starter strain for dairy use; absence of hemolytic activity in *Lactobacillus* indicates that these bacteria are non-virulent (De Vuyst et al., 2003).

Detection of inhibitory activity

All of the eight *Lactobacillus* species retained were tested for their antimicrobial activity (Table 3 and Figure 1). The zones of inhibition in the spot in the lawn method were more easily visualized. Our results show that the isolated strains (*L. plantarum* and *Lactobacillus rhamnosus*) had

inhibitory effect against *S. aureus*, with the most inhibiting species was *L. plantarum* (58). The inhibiting activity of *Lactobacillus* sp. could be as a result of two factors; the first is the production of lactic or acetic acid; indeed, the *Lactobacillus* are known for a great resistance to acid pH (with pH close to 3.5) contrarily to the other genera of lactic acid bacteria (Wong, and Chen, 1988; Podolak et al., 1996; Wilson et al., 2005); whereas the second comes from the production of another substance and probably of the bacteriocins (Larsen et al., 1993; Oyetayo et al., 2003; Avila et al., 2005; Cocolin et al., 2007).

Determination of the antimicrobial compounds

Our results of the inhibition study show that several strains can give an inhibition zone on solid medium. All tests of the determination of the antimicrobial compounds showed that the eight *Lactobacillus* strains cannot regenerate H₂O₂. The effect of acidity and phage lyses were also studied and no effect was observed. Juilliard et al. (1987) reported that the production of H₂O₂ occurred in aerobic conditions. The phages can be the origin of bacterial growth inhibitions (McGrath et al., 2002; Lewis et al., 1991). The test of the phage appeared negative for the strains of *Lactobacillus*. Moreover, the synergism effect of the antibacterial action cannot exclude the effect of acids (Deegan et al., 2006), hydrogen peroxide (Piard and Desmazeaud, 1991), diacetyl (Condon, 1987) or bacteriocin-like substances (Alexandre et al., 2002) by the strains of *Lactobacillus*.

The effect of the proteolytic enzymes on the inhibiting substance

All bacteriocins compounds are of protein nature (Callewaert and Vuyst, 2000; Aslim et al., 2005). To identify this nature of the antimicrobial substances, the


Table 2. Physiological and biochemical characters of *Lactobacillus* strains having an antimicrobial activity isolated from fresh raw goat's milk.

Carbohydrate	58	68	<i>Lactobacillus plantarum</i> ATCC 14917
Blanc	-	-	-
Glycerol	-	-	-
Erythritol	-	-	-
D-Arabinose	-	-	-
L-Arabinose	+	+	+
D-Ribose	+	+	+
D-Xylose	-	-	-
L-Xylose	-	-	-
D-Adonitol	-	-	-
MDX	-	-	-
D-Galactose	+	+	+
D-Glucose	+	+	+
D-Fructose	+	+	+
D-Mannose	+	+	+
L-Sorbose	-	-	-
L-Rhamnose	-	-	-
Dulcitol	-	-	-
Inositol	-	-	-
D-Mannitol	+	+	+
D-Sorbitol	+	+	+
MDM	+	-	+
MDG	-	-	-
NAG	+	+	+
Amygdaline	+	+	+
Arbutine	+	+	+
ESC	+	+	+
Salicine	+	+	+
D-Cellobiose	+	+	+
D-Maltose	+	+	+
D-Lactose ¹	+	+	V
D-Melibiose	+	+	+
D-Saccharose	+	+	+
D- Trehalose	+	+	+
Inulin	-	-	-
D-Melezitose	+	+	+
D-Raffinose	+	+	+
Amidon	-	-	-
Glycogen	-	-	-
Xylitol	-	-	-
Gentiobiose	+	+	+
D-Turanose	+	+	+
D-Lyxose	-	-	-
D-Tagatose	-	-	-
D-Fucose	-	-	-
L-Fucose	-	-	-
D-Arabitol	-	-	-
L-Arabitol	-	-	-
GNT	+	+	+
2KG	-	-	-
5KG	-	-	-
Identified as;	<i>Lactobacillus plantarum</i>	<i>Lactobacillus plantarum</i>	<i>Lactobacillus plantarum</i>

MDX: Methyl-β-D-xylopyranoside; MDM: methyl-α-D-mannopyranoside; MDG: methyl-α-D-glucopyranoside; NAG: N-acetylglucosamin; ESC: esculin citrate iron; 2KG: potassium 2-cetogluconat; 5KG: potassium 5-cetogluconat; GNT: potassium gluconate. ¹Bovine origin.

Table 3. Interactions between 8 selected *Lactobacillus* strains on solid medium and the diameter of inhibition zones measured (mm).

Code strain	58	55	68
58	0	7	5
55	8	3	5
68	6	3	4
2	6	4	2
13	3	3	5
52	4	7	2
31	7	4	3
54	5	4	5

**Figure 1.** Interactions between the isolates of the *Lactobacillus* species with the *Lactobacillus plantarum* (Strain 58) on solid medium.**Table 4.** The action of the proteolytic enzymes and the heat treatment on the antimicrobial activity of the crude supernatants of *Lactobacillus* strains towards the growth of *Staphylococcus aureus*.

Strain	α -Chymotrypsin	Trypsin	Heath treatment (100°C)
58	-	-	+
68	-	-	-

action of proteolytic enzymes (trypsin and α -chymotrypsin) was tested and the results are noted in Table 4, showing the effect of enzyme and heat treatment on the antimicrobial activity of the crude supernatant. The actions of α -chymotrypsin totally reduced the antimicrobial activity of the *Lactobacillus* strain, whereas only one *Lactobacillus* strain (58) could resist to the heat treatment at 100°C. Several bacteriocin of *Lactobacillus* have a large antimicrobial spectrum for Gram positive and Gram negative bacteria; for example the plantaricin C19 produced by *L. plantarum* (Nettles et al., 1993). The antimicrobial substances produced by lactic acid bacteria strains isolated from the goat's milk were investigated by Klaenhammer (1988). Our results indicate the case of the

inhibiting substances produced by *L. plantarum* (58) strain.

During the characterization of the bacteriocin, important variations in the spectra of activity were noted. It was also observed that the sensitivity of a strain depends on the genera, the species and even on the subspecies (Kalchayanand et al., 1992). The activity of strain supernatant was lost after the treatment with proteolytic enzyme, indicating that active component secreted extracellularly was proteinaceous in nature and confirmed that growth inhibition of sensitive strain of *Lactobacillus* was caused by bacteriocin (Ghrairi et al., 2008).

Inhibitory effect of *Lactobacillus* against *S. aureus*

The inhibitory spectrum of culture supernatants of eight *Lactobacillus* strains was assayed by the agar well diffusion method. The inhibitory activity directly evaluated against food spoilage bacteria and food-borne pathogen including three strains (*S. aureus*, *E. coli* and *Bacillus* sp.) is shown in Table 5. The results obtained in the interaction between the most powerful *Lactobacillus* shows an inhibition of eight strains (Figure 3). A strong inhibition

Table 5. Interactions and inhibition spectrum of *Lactobacillus* strains towards food spoilage bacteria and food-born pathogen *Staphylococcus aureus*.

Strain	<i>Staphylococcus aureus</i>		Inhibition (%)
	MNT	MT	
58	28	20	28.5
68	19	14	26.3
Average	21.87 ± 4.85	15 ± 2.26	31.41

MNT, Medium not buffered, MT, buffered medium.




Figure 2. The action of the proteolytic enzymes and the heat treatment on the antimicrobial activity (inhibition zones) of crude supernatant towards the growth of *Staphylococcus aureus*. a, α -Chymotrypsin; b, crude supernatants; c, crude supernatants with heat treatment at 100°C; d, Trypsin.

of *S. aureus* growth was obtained by *L. plantarum* (58). This inhibition was due to an inhibiting substance like bacteriocin. All the tests used in this study confirmed the nature of this substance as shown by Tagg et al. (1971) and Todorov and Dicks (2005). *Lactobacillus* strains 52, 54, 31, 2 and 13 expressed an inhibition growth of *S. aureus*, *Bacillus* sp. and *E. coli*, with various diameters but lower than those obtained by strains 58, 55 and 68. *L. plantarum* strains (58) gave an inhibition diameter of 20 mm for *S. aureus*. This strain (58) was retained as producing bacteriocin for carrying out this study.

Generally, the median values recorded by measuring the diameter of the inhibition zones showed that the Gram positive bacteria (*S. aureus*, 15 ± 2.26 mm) were sensitive to the inhibiting substances produced by the *Lactobacilli* compared to the Gram negative bacterium (*E. coli*, 8.62 ± 1.84 mm) (Figure 2). Todorov and Dicks (2005) observed that a high level of bacteriocin was produced when the cells were grown in the presence of K_2HPO_4 . However, little is known about the influence of

potassium ions on the production of bacteriocins. In the case of a bacteriocin produced by *L. plantarum*, level of K_2HPO_4 is needed to increase bacteriocin production (Todorov et al., 2005). The growth conditions used in this study were also similar to those cited above.

Growth kinetics and acidification

Evolution of pH and acidity in mixed culture in skim milk

The technological aptitude of the lactic acid bacteria is often based on the study of the acidifying capacity. All the species of *Lactobacillus* retained showed a high acidifying activity, which exceeded that produced by *S. aureus* which was 36.3°D in 24 h (Figure 4). *L. plantarum* (58) produced the highest level of lactic acid 50°D in 24 h. The other species of *Lactobacillus*, *L. plantarum* (68) produced 48 and 43.5°D, respectively in 24 h. In mixed culture of *L. plantarum* and *S. aureus*, a high production of lactic acid (69.4°D) was observed, whereas the decrease in the production of lactic acid in mixed culture with *L. plantarum* (68) (59.4°D) was noted.

Furthermore, the kinetics of pH evolution (Figure 4) showed that *L. plantarum* decreased the pH up to 4.32 in 24 h, whereas the pH of *L. plantarum* (68) was 4.76. The final pH of the culture of *S. aureus* was 5.13. In the mixed cultures, the pH reached 4.57 in *L. plantarum* (68) and 4.6 in *L. plantarum* (58). Lactic acid production in the culture of *L. plantarum* and *Lactobacillus brevis* studied by Kask et al. (1999) and Katina et al. (2002) was 50°D. These results are in agreement with our strains and in particular with *L. plantarum*. The latter produces a quantity of lactic acid higher than 40°D and almost reaches a maximum value of 100°D (Katina et al., 2002). Moreover, the most important quantity of lactic acid production was observed in *L. plantarum* (52°D) in pure culture after 72 h. Whereas, in mixed culture with *S. aureus* the species of *Lactobacillus* produced a higher quantity of acid, which was 71°D in *L. plantarum*, followed by *L. plantarum* (68) with 64°D. In synthetic medium saturated with glucose, Callewaert and De Vuyst (2000) reported that *L. reuteri* produces an acidity of 400°D in 24 h.




Figure 3. The inhibiting activity of *Lactobacillus* towards *Staphylococcus aureus* by the appearance of the clear zones around the colonies in none buffered (A) and buffered medium (B).




Figure 4. Acidity (shaded symbol) and pH (empty symbol) evolution in pure culture and mixed culture of *Lactobacillus plantarum* (58) and *Staphylococcus aureus* in milk.

Hence, the results of the present study support the controlling of *S. aureus* at the beginning of the dairy process when pH and temperature conditions may favor

enterotoxin production. Our results suggest that the growth in mixed culture of *L. plantarum* and *S. aureus* can take into account their interactions.




Figure 5. Kinetics of growth of *L. plantarum* (■) and *Staphylococcus aureus* (◆) in pure culture and *Staphylococcus aureus* (▲) in mixed culture in milk.

Growth evolution in mixed culture in milk medium

Enumeration of *S. aureus* in pure and mixed culture with *Lactobacillus* sp.

The initial enumeration of the two *Lactobacillus* strains *L. plantarum* (58), and *L. plantarum* (68) was 3.19 and 3.9 log cfu/ml, respectively. After 24 h of incubation the number of *L. plantarum* was 8.19 and 5.07 log cfu/ml for *S. aureus* in pure culture. Moreover, the number of the latter bacteria increased by 1.85 log cfu/ml after 24 h of incubation. In mixed culture after 24 h of incubation, a reduction of 1.6 log cfu/ml in the number of *S. aureus* was observed. This reduction proved the inhibiting effect of *L. plantarum* (Figure 5). A decrease of inhibition effect was observed for the other strains of *Lactobacillus* sp. towards *S. aureus* (results not showed).

After 72 h of incubation, viable cell number of *L. plantarum* (58) and *L. plantarum* (68) reached 8.05 and 8 log cfu/ml respectively. However, no growth could be detected for *S. aureus* after 72 h of incubation in the presence of *L. plantarum*. The inhibiting activity was slightly weak in the two other *Lactobacillus* species than *L. plantarum* (68). This variation of the inhibiting effect of the species of *Lactobacillus* towards *S. aureus* was also observed by Rodriguez et al. (2005). Production of multiple bacteriocins by *L. plantarum* caused an important inhibition of *S. aureus* (Hernandez et al., 2005).

The research work of Arquès et al. (2005) showed that after 72 h incubation, the number of *S. aureus* falls down to 0.46 log cfu/ml.

Effect of crude supernatant of *L. plantarum* on the growth of *S. aureus*

The influence of *L. plantarum* (58) supernatant was tested for evaluating the growth of the test strain of *S. aureus* after 6 h of incubation (Figure 6). The results show clearly the absence of growth of *S. aureus* in first dilutions (1/2 and 1/4) in which death rate was higher than 90% (Figure 6), whereas for dilutions of 1/256 and 1/1024 death rates were in close proximity to 50%. Moreover, after 24 h of incubation, death rate in the first three dilutions (1/2, 1/4 and 1/8) could reach 95.1, 93.4 and 92.5, respectively, compared to the test strain growth, while the death rate decreased considerably in dilutions 1/256 and 1/1024 and reached 4 and 4.9%, respectively.

In optimal growth conditions of the bacteriocin producer strain, *Lactobacillus acidophilus*, the culture supernatant contains 4.9 mg/ml of peptides (Kanatani et al., 1995). This concentration of protein in the culture supernatant was two times higher compared to that of *L. plantarum* (58). These results are, however, not surprising since it is well known that the culture medium and incubation




Figure 6. Effect of various dilutions of crude supernatant of *Lactobacillus plantarum* (58) heated with 100°C towards the growth of *Staphylococcus aureus*.

conditions affect the bacteriocins production in the genus of *Lactobacillus* (Todorov et al., 2005 and Babic et al., 2011). The death rate observed was inversely proportional to the dilutions level. Inhibition was high in the first dilutions where the concentration of peptides was higher than 2.5 mg/ml. The concentrations of peptides (0.31 mg/ml) in 1/8 dilution produced a middle inhibiting effect near 40% of death. In the last dilution (1/1024) which represents peptides concentration of 2.4 µg/ml, the death rate at 24 h was near 4.9%. Ananou et al. (2007) reported that the addition of enterocin AS-48 had an inhibitory effect on the growth of *Staphylococcus* sp., reducing viable counts below detection limits for the highest bacteriocin concentration tested (40 µg/ml). However, the bacteriocin concentration required to inhibit *Staphylococcus* growth during prolonged incubation was markedly higher compared to the minimal bactericidal concentration value of 15 µg/ml. It is well known that the activity of bacteriocin can be influenced by the chemical composition and the physical conditions of food.

In conclusion, the lactic acid bacteria originally isolated from raw goat's milk are probably the best candidates for improving the microbiological safety of dairy product because they are well adapted to the condition of milk and should be more competitive than lactic acid bacteria from other sources. Therefore, the results from the present study suggest that the bacteriocin-producing strain *L. plantarum* (58) could be used to improve the

safety of traditional fermented foods of dairy origin where *Lactobacillus* commonly occur. The selective use of this bacteriocinogenic strain may improve the microbiological quality of such foods.

REFERENCES

- Alexandre DP, Silva MR, Souza MR, Santos VLM (2002). Artisanal Antimicrobial activity of lactic acid bacteria from undermined cheese against indicator microorganisms. *Arch. Bra. Med. Veter. Zoot.* 54: 424-428.
- Alvarez-Martin P, Florez, AB, Barranco H, Mayo B (2008). Interaction between dairy yeasts and lactic acid Bacteria strains during milk fermentation. *Food Control.* 19: 62-70.
- Ananou S, Maqueda M, Martinez-Bueno M, Gavez A, Valdivia E (2007). Bactericidal synergism through enterocin AS-48 and chemical preservatives against *Staphylococcus aureus*. *Lett. Int. Appl. Microbiol.* 45: 19-23.
- Arquès JL, Rodriguez E, Gaya P, Medina M, Guamis B, Nunez M (2005). Inactivation of *Staphylococcus aureus* in raw milk cheese by combination of high-pressure treatments and bacteriocin-producing lactic acid bacteria. *Int. Dairy. J.* 15: 893-900.
- Aslim B, Yuksekdog ZN, Sarikaya E, Beyatli Y (2005). Determination of the bacteriocin-like substances produced by nap lactic acid bacteria isolated from Turkish dairy products. *LWT.* 38: 691-694.
- Avila M, Guard S, Medina M, Nunez M (2005). Effects of milk inoculation with bacteriocin-producing lactic acid bacteria on have djunct cheese culture. *J. Food Prot.* 68: 1026-1033.
- Babić I, Markov K, Kovačević D, Trontel A, Slavica A, Đugum J, Čvek D, Svetec IK, Posavec S, Frece J (2011). Identification and characterization of potential autochthonous starter cultures from a Croatian "brand" product "Slavonski". *J. Meat. Sci.* 02: 003.

- Badis A, Guetarni D, Moussa-Boudjema B, Henni DE, Kihal M (2004). Identification and technological properties of lactic acid bacteria isolated from raw goats milk of furnace Algerian races. *Food Microbiol.* 21(5): 579-588.
- Barefoot SF, Klaenhammer TR, 1983. Detection and activity of lactacin B, has bacteriocin produced by *Lactobacillus acidophilus*. *Appl. Env. Microbiol.* 45(6): 1808-1815.
- Blay G, Lacroix C, Zihler A, Fliss I (2007). In vitro Inhibition activity of nisin has nisin Z intestinal pediocin PA-1 and antibiotics common bacteria., *Lett. in Appl. Microbiol.* 45: 252-257.
- Bradford M (1976). A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding. *Annal. Biochemistry*, 72: 248-254.
- Budde BB, Hornbaek T, Jacobsen T, Barkholt V, Koch AG (2003). *Leuconostoc carnosum* 4010 has the potential for uses ace has protective culture for vacuum-packed meatuses: culture insulation, bacteriocin identification, and meat's experiments application. *Int. J. Food Microbiol.* 83: 171-184.
- Callewaert R, De Vuyst L (2000). Bacteriocins production with *Lactobacillus amylovorus* DCE 471 is improved and stabilized by EDF-batch fermentation. *Appl. Env. Microbiol.* 66: 606-613.
- Campos A, Rodriguez O, Calo-mata P, Prado M, Barros-velazquez J (2006). Preliminary characterization of bacteriocins from *Lactococcus lactis*, *Enterococcus faecium* and *Enterococcus mundtii* strain isolated from turbot (*Psetta maxima*). *Food. Res. Int.* 39: 356-364.
- Carr FJ, Chill D, Maida NR (2002). The lactic acid bacteria: With literature Survey. *Current. Rev. Microbiol.* 28(4): 281-370
- Casalta E, Vassal Y, Desmazeaud MJ, Casabianca F (1995). Comparaison de l'activité acidifiante de souche de *Lactococcus lactis* isolées de lait et de fromage Corse. *Lebensmittel-Wissenschaft Technol.* 28: 291-299.
- Cleveland J, Montville TJ, Nes IF, Chikindas ML (2001). Bacteriocins: Safe, natural antimicrobials for food preservation, *Int. J. Food Microbiol.* 71: 1-20.
- Cocolin L, Foschino R, Comi G, Fortina MG (2007). Description of the bacteriocins produced by two strains of *Enterococcus faecium* isolated from Italian goat milk. *Food Microbiol.* 31: 753-758.
- Condon S (1987). Responses of lactic acid bacteria to oxygen. *FEMS Microbiol. Rev.* 46: 269-280.
- Crow VL, Coolbear T, Holland R, Pritchard GG, Martley FG (1993). Starters as finishers starter properties relevant to cheese ripening. *Int. Dairy. J.* 3: 423-460.
- De Man JC, Rogosa M, Sharpe ME (1960). A medium for the cultivation of *Lactobacillus*. *J. Appl. Bacteriol.* 23: 130-135.
- De Vuyst L, Foulquie MR, Revets H (2003). Screening for enterocins and detection of hemolysin and vancomycin resistance in *Enterococci* of different origins. *Int. J. Food Microbiol.* 84: 299-318.
- Deegan L H, Cotter PD, Hill C, Ross P (2006). Bacteriocins: Biological tools for bio-safeguarding and shelf-life extension. *Int. Dairy. J.* 16: 1058-1071.
- Desmazeaud M, Cogan TM (1996). Role of cultures in cheese ripening. In: Cogan TM, Joined JP (Eds) *Dairy Starter Cultures*. VCH Publishers. Inc. New York. pp. 207-231.
- Ennahar S, Sashihara T, Sonomoto K, Ishizaky A (2000). Class IIa bacteriocins: biosynthesis, structure and activity. *Microbiol. Rev.* 24: 85-106.
- Fitzsimmons NA, Cogan TM, Condon S, Beresford T (1999). Phenotypic and genotypic characterization of not-choke lactic acid bacteria in mature cheddar cheese. *Appl. Env. Microbiol.* 65: 318-3426.
- Fleming HP, Erchells JL, Caslilow R (1975). Microbiol inhibition on isolate *Pediococcus* from cucumber bunc. *Appl. Env. Microbiol.* 30: 1040-1042.
- Gálvez A, Abriouel H, Lopez RL, Ben Omar N (2007). Bacteriocin-based strategies for bio-preservation. *Int. J. Food Microbiol.* 120: 51-70.
- Ghriri T, Frere J, Berjeaud JM, Mania M (2008). Purification and characterisation of bacteriocins produced by *Enterococcus faecium* from Tunisian rigouta cheese. *Food. Control*, 19: 162-169.
- Guessas B (2007). Les potentialités métaboliques des bactéries lactiques isolées du lait cru de chèvre dans le bio-contrôle de *Staphylococcus aureus*. Thèse de doctorat d'état. Département de biologie, Faculté des sciences, Université d'Oran, Es-Senia, p.165.
- Guessas B, Hadadji M, Saidi N, Kihal M (2006). Inhibition of *Staphylococcus aureus* growth in milk by lactic acid bacteria. *Dirassat*, 32(3): 53-60.
- Heikkila MP, Saris PEJ (2003). Inhibition of *Staphylococcus aureus* by the commensal bacteria of human milk. *J. Appl. Microbiol.* 95: 471-478.
- Hernandez D, Cardell E, Zarate V (2005). Antimicrobial activity of lactic acid bacteria isolated from Tenerife cheeses: Initial characterization of plantaricin TF 711, bacteriocin like substance produced by *Lactobacillus plantarum*. TF711. *J. Appl. Microbiol.* 99: 77-84.
- Jacobsen T, Budde BB, Koch AG (2003). Application of *Leuconostoc carnosum* for biopreservation of cooked meat products. *J. Appl. Microbiol.* 95: 242-249.
- Jamuna M, Babusha ST, Jeevaratnam K (2005). Inhibitory efficacy of nisin and bacteriocins from *Lactobacillus* isolates against food spoilage and pathogenic organisms in model and food systems. *Food Microbiol.* 22: 449-454.
- Joffin JN, Leyral G (1996). Technical microbiology. Teaching regional center of Documentation of Aquitaine Bordeaux. France, pp.219-223.
- Kaban G, Kaya M (2006). Effect of choke culture on growth of *Staphylococcus aureus* in sucuk. *Food. Control*, 17: 797-801.
- Kabuki T, Uenishi H, Watanabe M, Seto H, Nakajima H (2007). Characterization of a bacteriocin, Thermophilin 1277, produced by *Streptococcus thermophilus* SB1277. *Let. Appl. Microbiol.* 102: 971-980.
- Kaihei O, Kumar R, Sumie S, Watanabe K, Prakash TJ (2011). Lactic acid bacteria isolated from ethnic preserved meat products of the Western Himalayas. *Food Microbiol.* 28: 308-1315.
- Kalchayanand NR, Hanilin MB, Ray B (1992). Gram-negative Sublethal injury makes and resisting Gram-positive bacteria sensitive to the bacteriocins pediocin AcH and nisin. *Lett. Appl. Microbiol.* 15: 239-243.
- Kanatani K, Oshimura M, Sano K (1995). Isolation and characterization of acidocin A and cloning of the acteriocin gene from *Lactobacillus acidophilus*. *Appl. Env. Microbiol.* 61: 1061-1067.
- Kask S, Laht MT, Stake P, Paalme T (1999). A study on growth characteristics and nutrient of consumption *Lactobacillus plantarum* in A-stat culture. *Antonie van Leeuwenhoek.* 75: 309-320.
- Katina K, Sauri M, Alakomi HL, Mattila-Sandholm T (2002). Potential of lactic acid bacteria to inhibit rope spoilage in wheat sourdough bread. *Lebensmittel-Wissenschaft Technol.* 35(1): 38-45.
- Kihal M, Henni JD, Prevost H, Diviès C (2006). A new manometric method for measuring carbon dioxide production by starter culture: a case of *Leuconostoc mesenteroides*. *Afr. J. Biotechnol.* 5: 378-383.
- Kihal M, Prevost H, Lhotte ME, Huang DQ, Diviès C (1996). Instability of plasmid-encoded citrate permease in *Leuconostoc*. *J. Appl. Microbiol.* 22: 219-223.
- Klaenhammer TR (1988). Bacteriocins of lactic acid bacteria. *Biochemical*, 70: 37-49.
- Klaenhammer TR (1993). Genetics of bacteriocins produced by lactic acid bacteria. *Microbiol. Rev.* 12: 39-86.
- Klein G, Pack A, Bonaparte C, Reuter G (1998). Taxonomy and physiology of probiotic lactic acid bacteria. *Int. J. Food Microbiol.* 41: 103-125
- Larsen AG, Vogensen FK, Josephsen J (1993). Antimicrobial activity of lactic acid bacteria isolated from sour doughs: purification and characterization of bavaricin has, has bacteriocin produced by *Lactobacillus bavaricus* MI401. *J. Appl Bacteriol.* 75: 113-22.
- Lewis CB, Kaiser A, Montville TJ (1991). Inhibition of food-limits bacterial pathogens by bacteriocins from lactic acid bacteria isolated from meat. *Appl. Environ. Microbiol.*, 57: 1683-1688.
- Mc Auliffe O, Ross RP, Hill C (2001). Lantibiotics structure, biosynthesis and mode of action. *FEMS Microbiol. Rev.* 25: 285-308.
- Miteva V, Ivanova I, Budakov I, Pantev A, Stefanova T, Danova S, Monchev P, Mitev V, Dousset X, Boyaval P (1998). Detection and characterization of has Novell antibacterial substance produced by *Lactobacillus delbrueckii* strain 1043. *J. Appl. Microbiol.* 85: 603-614.
- Möll GN, Van der Akker E, Hauge HH, Nissen-Meyer J, Nes IF, Konings WN, Driessen AJM (1999). Complementary and overlapping selectivity of the two-peptide bacteriocins plantaricin EF and JK. *J. Bacteriol.* 181: 4848-4852.

- Moulay M, Aggad H, Benmechene Z, Guessas B, Henni DE, Kihal M (2006). Proteolytic activity of cultivable lactic acid bacteria isolated from Algerian raw goat's milk. *World. J. Dairy Food. Sci.* 1: 12-18.
- Nettles CG, Barefoot SF (1993). Biochemic and genetic characteristics of bacteriocins of food-associated lactic acid bacteria. *J. Food. Protein.*, 56: 338-356.
- Otero CM, Macias NR (2006). Inhibition of *Staphylococcus aureus* by H₂O₂-producing *Lactococcus gasserii* isolated from the vaginal leaflet of cattle. *Anim. Reprod. Sci.* 96: 35-46.
- Oyetayo VO, Adetuyi FC, Akinyosoye FA (2003). Safety and Protective effect of *Lactobacillus acidophilus* and *Lactobacillus casei* used as probiotic agent in-vivo. *Afr. J. Biotechnol.* 2: 448-452.
- Pal A, Ramana KV (2010). Purification and characterization of bacteriocin from *Weissella paramesenteroides* Dfr-8, an isolate from cucumber (*Cucumis sativus*). *J. Food. Biochem.* 34(5): 932-948.
- Papagianni M, Anastasiadou S (2009). Pediocins: the bacteriocins of *Pediococci*. Sources, production, properties and applications. *Microbial. Cell Factories*, 8: 3.
- Piard JC, Desmazeand M (1991). Inhibiting factors produced by lactic acid bacteria share L. oxygen metabolites and catabolism end-products. *Lait.* 71: 525-541.
- Podolak PK, Zayas JF, Kastner CL, Fung DYC (1996). Inhibition of *Listeria monocytogenes* and *Escherichia coli* O157:H7 on beef by application of organic acids. *J. Food. Prot.* 59: 370-373.
- Pringsulaka O, Patarasinpaiboon N, Suwannasai N, Atthakor W, Rangsiruji A (2011a). Isolation and characterisation of a novel Podoviridae-phage infecting *Weissella cibaria* N 22 from Nham, a Thai fermented pork sausage. *Food. Microbiol.* 28: 518-525.
- Pringsulaka O, Thongngam N, Suwannasai N, Atthakor W, Pothivejkul K and Rangsiruji A (2011b). Partial characterisation of bacteriocins produced by lactic acid bacteria isolated from Thai fermented meat and fish products. *J. Food. Control.* In press. pp. 1-5.
- Rodriguez E, Calzada J, Arquès JL, Rodrigues JM, Nunez M, Medina M (2005). Antimicrobial activity of Pediocin-producing *Lactococcus lactis* on *Listeria monocytogenes*, *Staphylococcus aureus* and *Escherichia coli* O157:H7 in cheese. *Int. Dairy. J.* 15: 51.
- Schillinger U, Lücke K (1989). Antimicrobial activity of *Lactobacillus sake* isolated from meat. *Appl. Env. Microbiol.* 55: 1901-1906.
- Sookkhee S, Chulasiri M, Prachyabru W (2001). Oral Lactic acid bacteria from healthy cavity of Thai volunteers: inhibition of oral pathogens. *J. Appl. Microbiol.* 90: 172-179.
- Stiles ME (1996). Biopreservation by lactic acid bacteria. *Antonie. Leeuwenhoek. J.* 70: 331-345.
- Stiles ME, Wilhelm H, Holzappel WH (1997). Lactic acid bacteria of foods and their current taxonomy. *Int. J. Food. Microbiol.* 36: 1-29.
- Tagg JR, Dajani AS, Wannamaker LW (1976). Bacteriocins of Gram-positive bacteria. *Bacteriol. Rev.* 40: 722-756
- Tagg JR, Mc Given AR (1971). Assay system for bacteriocins. *J. Appl. Microbiol.* 21: 943.
- Tahara T, Kanatani K (1996). Isolation, partial characterization and mode of action of acidocin J1229, a bacteriocin produced by *Lactobacillus acidophilus* JCM 1229. *J. Appl. Bacteriol.* 81: 669-677.
- Todorov SD, Dicks LMT (2005). *Lactobacillus plantarum* isolated from molasses produces bacteriocins active against Gram-negative bacteria. *Enzy. Microbiol. Technol.* 36: 318-326.
- Vaughan EE, Caplice E, Looney R, O'Rourke N, Coveney H, Daly C, Fitzgerald GF (1994). Isolation from food sources, of lactic acid bacteria that produced antimicrobials. *J. Appl. Bacteriol.* 75: 118-123.
- Vaughan EE, Daly C, Fitzgerald GF (1992). Identification and characterization of helveticin V-1829, a bacteriocin produced by *Lactobacillus helveticus* 1829. *J. Appl. Bacteriol.* 73: 299-308.
- Vermeiren L, Devlieghere F, Debevere J (2004). Protective evaluation of meatus born lactic acid bacteria ace cultures for the biopreservation of cooked meatus products. *Int. J. Food. Microbiol.* 96: 149-164.
- Wilson AR, Sigee D, Epton HAS (2005). Anti-bacterial activity of *Lactobacillus plantarum* strain SK1 against *Listeria monocytogenes* is due to lactic acid production. *J. Appl. Microbiol.* 99: 1516-1522.
- Wong HC, Chen YL (1988). Effects of lactic acid bacteria and organic acids on growth and germination of *Bacillus cereus*. *Appl. Env. Microbiol.* 54: 2179-2184.