



[www.ajbrui.org](http://www.ajbrui.org)

*Afr. J. Biomed. Res. Vol. 25 (January, 2022); 33 - 38*

*Research Article*

# **Unsafe Sexual Practices and Associated Factors Among Female Sex Workers in Sagamu Local Government, Ogun State, Nigeria**

**\*Olu-Abiodun O<sup>1</sup> and Edet O.B.<sup>2</sup>**

<sup>1</sup>*School of Nursing, Ijebu-Ode, Ogun state, Nigeria.*

<sup>2</sup>*Department of Nursing Science, College of Medical Sciences, University of Calabar, Cross River State, Nigeria.*

## **ABSTRACT**

One of the public health challenges that have gripped the universe is the rapid spread of the Human Immunodeficiency Virus. Unprotected heterosexual intercourse remains central to HIV transmission in Nigeria, as in most of sub-Saharan Africa and female sex workers play a crucial role in the spread of the virus through unsafe sex practices. Globally, sex workers make up 9% of the total number of new HIV infections and while HIV prevalence among the population in Nigeria has been declining from its peak of 5.8% in 2001 to 1.4% in 2018, prevalence among sex workers remains high at 14.4%. A descriptive cross-sectional study of 81 Female Sex Workers out of the estimated total population of 360 FSWs based in brothels only, in Sagamu local government was selected. The majority 63(77.8%) of the respondents engaged in unsafe sex practices. The majority 65(80.3%) had a low to moderate risk perception of STI/HIV/AIDS. Almost half participants 36(44.4%) had the erroneous belief that they could not contract HIV. Statistically significant association was found between unsafe sexual practices and previous history of STIs among FSWs ( $p = 0.021$ ), reported current HIV status ( $p = 0.019$ ), alcohol use ( $p = 0.004$ ), and binge alcohol ( $p = 0.037$ ). Unsafe sex practice is highly prevalent among female sex workers in Sagamu with low to moderate risk perception for HIV/AIDS / STI.

**Keywords:** *Female sex workers, HIV/AIDS, Perception, Unsafe sexual practices, sexually transmitted infections*

\*Author for correspondence: Email: [tmtosin@yahoo.co.uk](mailto:tmtosin@yahoo.co.uk); Tel: +234 8061992744

*Received: June 2021; Revised version accepted: November 2021*

## **INTRODUCTION**

One of the public health challenges that have gripped the universe is the rapid spread of the Human Immunodeficiency Virus (HIV), the main cause of AIDS. In many countries, the HIV epidemic is concentrated in subgroups of the population like homosexuals, sex workers, etc. whose behavior exposes them to the high risk of acquiring HIV infection.

According to UNAIDS (2018) report, Sex workers accounted for about one in 10 infections in eastern Europe and central Asia and the Middle East, and North Africa. Population size estimates suggest there are nearly one million sex workers in need of preventive services. Available data on prevention program coverage for this key population ranged from 38% in South Sudan to 74% in Kenya (UNAIDS, 2018). Unsafe sex practices are responsible for the rapid spread of HIV among female sex workers, therefore, understanding unsafe sexual practices is one of the most important strategies in preventing the spread of HIV/AIDS and STIs among sex workers. Certain barriers and unsafe sexual practices increase the vulnerability of female sex workers to the risk of HIV infection.

Existing studies estimate the prevalence of HIV to be 14.4% among sex workers in Nigeria (UNAIDS, 2017),

compared with a prevalence of (1.3%) in the adult population (NACA,2017). Notwithstanding increased funding for HIV prevention activities among sex workers in Nigeria since the 1990s, HIV prevalence among sex workers has remained high over the past two decades. HIV prevalence among sex workers in 2007 was still over 30%, and in some cities like Kano and Abuja, 16.8% of all brothel-based sex workers are HIV-infected (NARHS,2012). Similarly, brothel-based sex workers face greater HIV risk in Nigeria, with a prevalence of 27.4%. (FMOH, 2013; NARHS, 2012) Thus, while HIV prevalence among the general population in Nigeria has been declining from its peak of 5.8% in 2001 to 4.1% in 2011, (Federal Ministry of Health, 2010), HIV prevalence among sex workers is still eight times higher than the general population (NACA,2015). Although the population of brothel-based sex workers is very small in size compared with the general Nigeria population, recent modeling of apportionment of new infections estimated that in some states (such as Ogun in south-west Nigeria and Benue in north-central Nigeria), sex workers and their clients account for about 18% of all new infections. (Oladembe, Adeniji, Omilaju, Adebayo, Somda, Fajemisin, 2010; Baba-Manu, Nwankwo & Wende, 2010). This study was therefore designed to determine the prevalence of unsafe sexual

practices, identify their perceived risk of contracting HIV/AIDS/ STI, and to access the beliefs of brothel-based sex workers about sex work in Sagamu Local Government of Ogun State.

## MATERIALS AND METHODS

**Study design:** This was a cross-sectional study designed to investigate the prevalence of unsafe sexual practices, identify their perceived risk of contracting HIV/AIDS/STI, and access the beliefs of brothel-based female sex workers about sex work in Sagamu Local Government of Ogun State.

**Study setting:** The study was carried out in the Sagamu local government of Ogun State, southwest Nigeria. Sagamu is a well-known commercial center in Ogun State. It is a densely populated area, dominated by different tribes in Nigeria specifically, Yoruba, Hausas, Igbos, and other minor tribes (Isekiri, Irobo, and Fulanis). According to the National Population Commission project population report, Sagamu had an estimated population number of 355, 900 and comprises males and females (NPC, 2016 and NBS, 2018).

**Data collection and sampling procedure:** The study employed an interview schedule to interview 81 female sex workers derived from the Cochran formula for estimating proportion in a cross-sectional study at a 95% confidence level. (Cochran, 1985). The participants were selected using the cluster sampling technique. First, we used a simple random sampling by balloting to select three out of the six brothel locations in Sagamu. Each Brothel had between 50 and 55 sex workers living in them. Then, we recruited all eligible consenting sex workers within the selected Brothels to participate in the study. Consenting female sex worker, resident and duly registered in the selected study brothel who had been in the business for at least three months were eligible to participate in the study.

The respondents were approached for an interview during their break period in the daytime when they had no or less few clients. A well-structured questionnaire was used to investigate the social demographics of the respondents, the prevalence of unsafe sexual practices, their perceived risk of contracting STIs and HIV/AIDS, and their beliefs about sex work. The questionnaire was pretested in a brothel in Ijebu ode, a town with similar characteristics and identity to the main study setting. The validity of the study instrument was ascertained by experts in the field. The face and content validity of the instrument was established. The reliability of the instrument was good with a Cronbach alpha value of 0.872.

**Ethical consideration:** Ethical approval was sought and obtained from Ogun State Human Research Ethics Committee with ref no: HPRS/381/324. Written consent was used to obtain consent for participation in the study. Approvals were obtained from the management of the selected brothels before the commencement of the study. The researcher read the consent form to the respondents and they were given time to ask questions. The respondents agreed to participate by

signing the consent form. Confidentiality was also assured by removing all identifiers of respondents from the questionnaire.

**Data analysis:** The data was sorted, arranged, and input into the spreadsheet and analyzed using the Statistical Product for Social Sciences version 20.0 (SPSS Inc. USA). Descriptive and inferential statistics were used to analyze the data. Proportions and mean (with standard deviation) were used to summarize the data. Chi-square test, student's t-test, and logistic regression analysis were carried out. Findings were presented with charts and tables.

## RESULTS

**Respondents' background characteristics:** The female sex workers who participated in the study had a mean age of 31.16 years with a standard deviation of 7.71 years.

**Table 1:**  
Socio-demographic characteristics of respondents (N = 81)

Characteristics	Frequency	Percent
<b>Age in years</b>	31.16 (7.71)*	
<b>Age group in years</b>	16-25	27.2
	26-35	49.4
	36-45	22.2
	46-55	1.2
<b>Ethnicity</b>	Yoruba	16.0
	Hausa	16.0
	Igbo	29.6
	Others	38.3
<b>Highest level of education completed</b>	Non- formal	9.9
	Primary	17.3
	Secondary	65.4
	Tertiary	7.4
<b>Religion</b>	Christianity	75.3
	Islam	23.5
	Traditional	1.2
<b>Marital status</b>	Never married	34.6
	Co-habiting	3.7
	Married	19.8
	Divorced	28.4
	Separated	11.1
	Widowed	2.5
<b>Other occupation</b>	None	35.8
	Self-employed	12.3
	Restaurant/bar	43.2
	Others	8.7
<b>Duration of sex work in years*</b>	3.40	2.41
<b>The average number of steady male partners*</b>	1.54	1.50
<b>Customer pick up point</b>	Brothel alone	95.1
	Brothel and street	4.9
<b>Customer service point</b>	Brothel alone	86.4
	Brothel and other private spaces	7.5
	Brothel and other public places	1.2
	Brothel, public, and private spaces	4.9

The most predominant age group was 26 to 35yrs (49.4%), followed by 16 to 25yrs (27.2%). Concerning ethnicity,

29.6% (24) were Igbos, while 16% (13) each were Hausa and Yoruba while 38.3 % (31) of the respondents came from the other ethnic groups. Fifty-three (65.4%) of the FSWs had a secondary education, 61 (75.3%) 58, were Christians, and 28 (34.6%) were never married. Twenty-three (28.4%) of the respondents were divorced while 16 (19.8%) were married. Concerning other occupations engaged in by the respondents, 35 FSWs (43.2%) work in the restaurant/bar, while 29 (35.8%) of them had no other occupation. The mean years of duration of sex work are 3.40 years with a standard deviation of 2.41 years. Also, the average number of steady partners per participant is 1.54 with a standard deviation of 1.50. Seventy-seven (95.1%) of the respondents picked up their customers in the brothels alone and 57 (83.8%) serviced their customers in the brothels only.

As shown in Table 2, 63 (77.8%) of the respondents engaged in unsafe sex practices. Fifty-four (66.7%) of them reported that they had concurrent sexual relationships, 37 (45.7%) reported the practice of anal sex, while 36 (44.4%) of them reported alcohol use before sex. Few of the respondents reported drug use before sex (14.8%) and injection of drugs before sex (8.6%).

**Table 2:**  
Prevalence of unsafe sexual practices

Unsafe sexual practices	Yes (%)	No (%)
Non-use of condom during sex, regularly	3 (3.7)	78 (96.3)
Non-use of condom during the last sexual act	1 (1.2)	80 (98.8)
The practice of anal sex	37 (45.7)	44 (54.3)
Concurrent sexual relationship	54 (66.7)	27 (33.3)
Drug use before sex	12 (14.8)	69 (85.2)
Injection of drugs before sex	7 (8.6)	74 (91.4)
Alcohol use before sex	36 (44.4)	45 (55.6)
Practice of unsafe sex (overall)	63 (77.8)	18 (22.2)

Table 3 shows what the respondents thought were the risk and protective factors for STI/HIV. The majority (87.7%) of the respondents agreed that unsafe sex practices constitute a risk for HIV/STI transmission. Sixty-five (80.3%) of them are aware of their chances of contracting HIV/AIDs and STIs. Sixty-eight (84.0%) of the FSW perceived that sexual intercourse with multiple partners increased the risk of contracting HIV/AIDs. Concerning risk perception for HIV, only 44 (54.3%) of the respondents perceived themselves to be at risk of contracting HIV/AIDs.

Table 4 shows the result of bivariate analysis between participant characteristics and unsafe sexual practices. There

**Table 3:**  
Perceived risk and protective factor for STI/HIV

Risk of unsafe sex practices (no of respondents)	Agree (%)	Undecided (%)	Disagree (%)
Risk behaviour linked to STI/HIV transmission	71 (87.7)	9 (11.1)	1 (1.2)
Awareness of chances of contracting STI/HIV	65 (80.3)	10 (12.3)	6 (7.4)
Sexual intercourse with multiple partners can increase the risk of contracting STI/HIV	68 (84.0)	9 (11.1)	4 (4.9)
My sexual practices put me at risk of contracting STI/HIV	44 (54.3)	20 (24.7)	17 (21.0)

was statistically significant association between unsafe sexual practices and previous history of STIs ( $\chi^2 = 5.313, p = 0.021$ ), reported current HIV status ( $\chi^2 = 7.972, p = 0.019$ ), alcohol use ( $\chi^2 = 11.058, p = 0.004$ ), and binge alcohol ( $\chi^2 = 4.349, p = 0.037$ ). Participants who had a previous history of STIs, currently HIV positive, take alcohol, practice binge alcoholism were more likely to engage in unsafe sexual practices. The other participant characteristics did not show a significant relationship with unsafe sexual practices.

## DISCUSSION

The result of the study indicated that the majority (77.8%) of the study participants had engaged in unsafe sex practices such as anal sex (45.7%) concurrent sexual relationship (66.7%), alcohol use before sex (44.4%) and drug use before sex (14.8%). This is similar to a study by Mahapatra, Mohanty, Gurav and Ramesh, et al. (2013) on the prevalence of inconsistent condom use, concurrent sexual relationships, anal sex, and alcohol consumption before sex which reported that about half of the sex workers (51%) had engaged in risky sexual practices; 71% of FSWs reported inconsistent condom use with any client, 32% reported concurrent sexual relationships, 12% had engaged in anal sex and 54% reported consuming alcohol before sex. In this study, consistent condom use during sexual intercourse was self-reported by almost all the FSWs.

Majority (91.7%) of the FSWs perceived themselves to be at low/moderate risk of contracting STIs/HIV/AIDs despite their high risk of sexual activity which could, in turn, be a risk factor of contracting the infection. The perceived risk of HIV was found to be an important trigger among FSWs negotiating condom use with clients. Furthermore, risky sexual behavior despite preventive knowledge is less surprising if FSWs are unaware of the level of risk of contracting HIV when having unprotected sex or practicing unsafe sex, i.e., knowing how to reduce risk may not change behavior if the level of risk is perceived as low.

Most (87.7%) of the FSWs are also aware of the risk behaviour linked to HIV/STI transmission, 80.3% of the participant are aware of their chances of contracting STI/HIV/AIDs, (54.3%) of the respondents perceive themselves as being at risk of contracting HIV/AIDs. A study by Bruce, Bauai, Sapuri, Kaldor, Fairley, and Keogh, (2011), reported that although most FSWs were aware of HIV risk through sexual contacts, most perceived themselves as more at risk of infection through sex work than through their private sexual relationships.

**Table 4:**  
Relationship between participants' characteristics and unsafe sex practices

Characteristics (no of respondents)	unsafe sex practices		$\chi^2$ or t-test (*)	p-value	
	Yes (%)	No (%)			
<b>Age in years*</b>	30.82 (7.60)	32.44 (8.38)	0.779*	0.438	
<b>Age group in years</b>	16-25	18 (81.8)	4 (18.2)	0.672	0.879
	26-35	30 (75.0)	10 (25.0)		
	36-45	14 (77.8)	4 (22.2)		
	46-55	1 (100.0)	0 (0.0)		
<b>Ethnicity</b>	Yoruba	12 (92.3)	1 (7.7)	4.230	0.237
	Hausa	8 (61.5)	5 (38.5)		
	Igbo	20 (83.3)	4 (16.7)		
	Others	23 (74.2)	8 (25.8)		
<b>Highest level of education completed</b>	Non- formal	8 (100.0)	0 (0.0)	5.316	0.150
	Primary	8 (57.1)	6 (42.9)		
	Secondary	40 (75.5)	13 (24.5)		
	Tertiary	5 (83.3)	1 (16.7)		
<b>Religion</b>	Christianity	46 (75.4)	15 (24.6)	0.411	0.814
	Islam	15 (78.9)	4 (21.1)		
	Traditional	1 (100.0)	0 (0.0)		
<b>Marital status</b>	Single	22 (78.6)	6 (21.4)	0.370	0.946
	Co-habiting	2 (66.7)	1 (33.3)		
	Married	13 (81.2)	3 (18.8)		
	Divorced, Separated, Widowed	26 (76.5)	8 (23.5)		
<b>Other occupation</b>	None	24 (82.8)	5 (17.2)	6.792	0.079
	Self-employed	5 (50.0)	5 (50.0)		
	Restaurant/bar	26 (74.3)	9 (25.7)		
	Others	7 (100.0)	0 (0.0)		
<b>Duration of sex work in years*</b>	3.35 (2.18)	3.64 (3.18)	0.438*	0.663	
<b>Number of steady male partners*</b>	1.52 (1.62)	1.61 (1.11)	0.207*	0.837	
<b>History of STI</b>	No	14 (59.1)	9 (40.9)	5.313	0.021
	Yes	49 (85.5)	9 (14.5)		
<b>Knowledge of HIV Status</b>	No	18 (81.8)	4 (18.2)	0.468	0.494
	Yes	44 (74.5)	15 (25.5)		
<b>HIV status</b>	Negative	21 (61.8)	13 (38.2)	7.972	0.019
	Positive	17 (94.4)	1 (5.6)		
	Did not disclose	24 (82.8)	5 (17.2)		
<b>Alcohol use (76)</b>	Never	5 (41.7)	7 (58.3)	11.058	0.004
	Previous use	6 (75.0)	2 (25.0)		
	Current use	52 (85.2)	9 (14.8)		
<b>Binge alcohol consumption (70)</b>	No (Non-Binge)	23 (71.9)	9 (28.1)	4.349	0.037
	Yes (Binge)	44 (89.8)	5 (10.2)		
<b>Customer pick up point</b>	Brothel alone	59 (76.4)	18 (23.6)	1.202	0.272
	Brothel and street	4 (100.0)	0 (0.0)		
	Customer service point				
	Brothel alone	50 (71.4)	20 (28.6)	2.050	0.562
	Brothel and other private spaces	4 (66.7)	2 (33.3)		
	Brothel and other public places	1 (100.0)	0 (0.0)		
	Brothel, public, and private spaces	4 (100.0)	0 (0.0)		

\*t-test

This data also revealed that most FSWs had recently engaged in unsafe sex with clients despite adequate knowledge. According to Lammers, Wijnbergen and Willebrands, (2013), most FSWs also perceived themselves as more at risk of infection through sex work. In contrast, in a recent survey among the most at-risk populations in six states in Nigeria, over half of the sex workers did not consider themselves at risk of HIV infection. In Lagos state, with the highest concentration of sex workers in Nigeria, only 16% of brothel-based sex workers felt they were at risk, even though each has on average 34 clients per week. Despite their high-risk sexual

activity, many sex workers perceive their risk of HIV infection as low. (Messersmith, Kane, Odebiyi & Adewuyi, 2000). The respondents seemed not to connect knowledge and risk perception with their behaviour. This also confirms the results of similar studies in Nigeria by Messersmith et al. (2000), Federal Ministry of Health (2007), and Ankomah et al. (2011).

Many (44.4%) of the FSWs believed that they cannot contract HIV/AIDS. For many of these sex workers, belief and trust in God were important attributes that offered safety and security from infection Many were of the view that they will

never get infected because of their faith: "I will not catch it in Jesus' name."

There was a statistically significant association between unsafe sexual practices and previous history of STIs. Participants who had a previous history of STIs, currently HIV positive, take alcohol, practice binge alcoholism were more likely to engage in unsafe sexual practices. The other participant characteristics did not show a significant relationship with unsafe sexual practices. Similarly, reports from scientific research indicate a positive association between alcohol consumption and HIV risk. Moreover, FSWs who consume alcohol are more likely to engage in anal sex than those who do not, recent advances in HIV research have also highlighted concurrent sexual relationships as another important factor in increasing the rate of infection in a population. (Hutton, McCaul, Santora & Erbeling, 2008; Verma, Saggurti, Singh & Swain, 2010; Samet, Pace, Cheng, Coleman & Bridden, 2010).

The result shows that there was no evidence of a statistically significant relationship between the female sex workers' perception of their risk of contracting HIV and the practice of unsafe sex. ( $\chi^2 = 1.852$ ,  $p = 0.592$ ). According to Bruce, Bauai, Sapuri, Kaldor, Fairley, & Keogh. (2011), Studies on the influence of HIV knowledge on safer sex practice have reported mixed results. Although some have found conclusive associations, others have found mixed results consistent with our findings. Although some studies found an association between individual perception of HIV risk and safer sex practices, others have found mixed results. This implies that safer sex is complex and determined by several factors in different settings.

In conclusion, the findings highlight that safe sex practice is not determined by the availability of, access to, and promotion of condoms alone, but also by complex contextual, emotional, and other environmental factors. Until effective strategies are implemented to address these factors, no progress on HIV will be sustained. For FSWs, the vital element is empowerment. Overall, it means efforts should be targeted on health education, economic independence, and community development schemes to sustain their livelihoods. Furthermore, safer sex messages have reached most FSWs and they do not need to be convinced about it; however, they need to be enabled to practice it. Current health promotion messages are inadequate in addressing the complexities of sex work in Nigeria. For example, advising sex workers to limit the number of partners or always to use condoms, when the fee for one session of unprotected sex can equal two or three days' income, is an impractical and difficult behaviour to adapt. To succeed, interventions aimed at vulnerable groups need to engage them in developing programs that are relevant to their circumstances (Kaddour, Hafez & Zurayk, 2005; Obermeyer, 1999). Therefore, more efforts should be devoted to engaging and encouraging sex workers to be involved in developing effective sexual health messages that take account of the circumstances surrounding their lives. Without this, economic survival and the contextual circumstances of FSWs lives will continue to make current safe-sex messages obsolete and impractical. (Muñoz, Adedimeji & Alawode. 2010)

These results suggest the need for expanded intensive educational interventions targeting the FSWs and their clients

with messages emphasizing the importance of safe sex practices and simultaneously address the barriers to safe sex practice identified by some of the respondents. More efforts should be devoted to engaging and encouraging sex workers to be involved in developing effective sexual health messages that take account of the circumstances surrounding their lives. Without this, economic survival and the contextual circumstances of FSWs lives will continue to make current safe-sex messages obsolete and impractical. There is a need for intervention to address their poor risk perception about contracting STI/HIV/AIDs. Governments and organizations need to create an environment where sex workers can protect themselves against HIV, and easily access HIV prevention, testing, and treatment services. Effective interventions for sex workers must recognize the realities of FSWs' lives regarding the genuine constraints to adopting protective measures. In particular, our findings support the urgent need to move beyond a solely individual-level HIV-prevention approach, such as condom distribution, to structural environmental HIV prevention that facilitates female sex workers' ability to negotiate their risk environment in safer sex-work settings. Finally, we strongly recommend that programs must revisit the areas of misperceptions, avoid promoting messages such as abstinence and being faithful to one uninfected sexual partner that is considered unrealistic in the context of sex work, and intensify prevention efforts on illicit drugs and alcohol use for FSWs.

#### **Acknowledgements**

*The authors acknowledge all the female sex workers based in selected brothels in Sagamu local government area, Ogun state Nigeria, their managers, owners of the brothels and other workers who have contributed in one way or the other to the success of the research. We also acknowledge the research assistants who worked tirelessly for the success of the project.*

#### **REFERENCES**

- Adeneye, AK., Adeneye, AA., Mafe, MA., & Adeiga, AA. (2013)** Pattern of condom use and perceived risk of HIV infection among female sex workers in selected Brothels in Ogun State, Nigeria. *Int J Public Health Epidemiol.* 2(5):90– 100.
- Ankomah, A., Omoregie, G., Akinyemi, Z., Anyanti, J., Ladipo, O., & Adebayo, S. (2011).** HIV related risk perception among female sex workers in Nigeria. *HIV/AIDS-Research and Palliative Care* 3, pp. 93-100
- Baba-Manu, A., Nwankwo, E., & Wende, G. (2010).** HIV modes of transmission in Benue State: an analysis of the distribution of new infections. Abuja, Nigeria: Enhancing Nigeria Response to HIV/AIDS Project.
- Braunstein, S. L., Ingabire, E., Kestelyn, AU., Uwizire, L., Mwamarangwe, J., & Ntirushwa. (2011).** High human immunodeficiency virus incidence in a cohort of Rwandan female sex workers. *sexually transmitted diseases* 38:385–394.
- Bruce, E., Bauai, L., Sapuri, M., Kaldor, K., Fairley, K., & Keogh, L. (2011).** HIV knowledge, risk perception and safer sex practices among female sex workers. *Int J Womens Health.* 3: 53-61 doi: 10.2147/IJWH.S14669
- Cochran, W. G. (1985)** *Sampling Techniques*, 3rd Ed., New York: John Wiley and Sons, Inc.
- Cook, RL., Comer, DM., Wiesenfeld, HC., Chang, CC., & Tarter, R. (2006).** Alcohol and drug use and related disorders: An under recognized health issue among adolescents and young adults

- attending sexually transmitted disease clinics. *Sex Transm Dis* 33: 565–570
- Cook, RL., Clark, DB. (2005).** Is there an association between alcohol consumption and sexually transmitted diseases? A systematic review. *Sex Transm Dis* 32: 156–164. 12.
- Cusik, L. (2006). Widening the harm reduction agenda: From drug use to sex work. *International Journal of Drug Policy*, 17, 3–11.
- Federal Ministry of Health. (2010).** National HIV seroprevalence sentinel survey among pregnant women attending antenatal clinics in Nigeria. Abuja, Nigeria: Department of Public Health; 2011.
- Fisher, JC., Cook, PA., & Kapiga, SH. (2010).** Alcohol use before sex and HIV risk: situational characteristics of protected and unprotected encounters among high risk African women. *Sex Transm Dis* 37: 571–578.
- George, A., Sabarwal, S., & Martin, P. (2011).** Violence in contract work among female sex workers in Andhra Pradesh, India. *J Infect Dis* 2011;204:S1235-40.
- Kalichman, S., Pinkerton, S., Carey, M., Cain, D., & Mehlomakulu, V. (2011)** Heterosexual anal intercourse and HIV infection risks in the context of alcohol serving venues, Cape Town, South Africa. *BMC Public Health* 11: 807–817
- Lammers, J., Wijnbergen, S., & Willebrands, D. (2013)** Condom use, risk perception, and HIV knowledge: a comparison across sexes in Nigeria. *HIV/AIDS Auckl* 5: 283–293. doi: 10.2147/HIV.S31687
- Lane, T., Pettifor, A., Pascoe, S., Fiamma, A., & Rees, H. (2006)** Heterosexual anal intercourse increases risk of HIV infection among young South African men. *AIDS* 20: 123–125.
- Low, N., Broutet, N., Adu-Sarkodie, Y., Barton, P., Hossain, M., & Howkes, S. (2006).** Global Control of Sexually transmitted infections. *Lancet*. 2006; 368:2001–2016. [PubMed]
- Mahapatra, B., Lowndes, CM., Mohanty, SK., Gurav, K., & Ramesh, BM. (2013).** Factors Associated with Risky Sexual Practices among Female Sex Workers in Karnataka, India. *PLoS ONE* 8(4): e62167. doi: 10.1371/journal.pone.0062167
- Mayhew, S., Collumbien, M., Qureshi, A., Platt, L., Rafiq, N., Faisal, A., Lalji, N., & Hawkes, S. (2009):** Protecting the unprotected: mixed-method research on drug use, sex work and rights in Pakistan's fight against HIV/AIDS. *Sex Transm Infect* 2009;85 Suppl 2:31-6.
- Messersmith LJ, Kane TT, Odebiyi AI, Adewuyi AA. (2000):** Who's at risk? men's STD experience and condom use in southwest Nigeria. *Stud Fam Plann*. 2000; 31:203–216.
- Morisky, DE., Chiao, C., Stein, JA., & Malow, R. (2005)** Impact of social and structural influence interventions on condom use and sexually transmitted infections among establishment-based female bar workers in the Philippines. *J Psychol Human Sex*;17(1/2):45–63.
- National AIDS/STI Control Programme, Federal Ministry of Health (2010).** HIV/STI Integrated Biological Behavioural Surveillance Survey (IBSS) Abuja, Nigeria: Federal Ministry of Health; 2010.
- NACA (2017)** 'National Strategic Framework on HIV and AIDS: 2017 -2021' [pdf]
- NACA (2015)** 'Nigeria GARPR 2015' [pdf]
- National Reproductive Health Survey-NARHS 2012.** Shadow report on situation of sex workers in Nigeria.
- NAIIS. (2018).** The 2018 Nigeria HIV/AIDS Indicator and Impact Survey. Retrieved from <https://www.naiis.ng/>
- Okafor, UO., Crutzen, R., Ifeanyi, O., Adebajo, S. & Borne, HVD. (2017).** HIV prevalence and high-risk behaviour of young brothel and non-brothel based female sex Workers in Nigeria. *BMC Res Notes*10:380 DOI 10.1186/s13104-017-2712-8.
- Oladembe, R., Adeniji, GO., Omilaju, I., Adebayo, S., Somda, Z., & Fajemisin, O. (2010).** HIV modes of transmission in Ogun state: an analysis of the distribution of new infections. Abuja, Nigeria: Enhancing Nigeria Response to HIV/ AIDS Project.
- Open Society Foundations (2012).** Criminalizing condoms: How policing practices put sex workers and HIV services at risk in Kenya, Namibia, Russia, South Africa, USA, and Zimbabwe. Sexual Health and Rights Project. New York
- Panchanadeswaran, S., Johnson, S.C., Sivaram, S., Srikrishnan, A.K., Latkin, C.,& Bentley, M.E. (2008).** Intimate partner violence is as important as client violence in increasing street-based female sex workers' vulnerability to HIV in India. *International Journal of Drug Policy*, 19, 106-112.
- Patra, RK., Mahapatra, B., Kovvali, D., Proddutoor, L., & Saggurti, N. (2012).** Anal sex and associated HIV-related sexual risk factors among female sex workers in Andhra Pradesh, India. *Sex Health* 9: 430–437.
- Pettifor, A., & Rosenberg, N. (2011).** The Need to Focus on Sex Workers in Generalized Epidemic Settings. *Sexually Transmitted Diseases* 38(4):324–325.
- Price, J., & Cates, W. (2011).** Sex Workers Studies: The Science, Semantics and Politics of Targeting our HIV Prevention Response. *Sexually Transmitted Diseases* 38(5):395–397.
- Ramesh, B.M., Moses, R., Washington, S., Isac, B., Mohapatra., & Mahagaonkar, SB. (2008).;** IBBA Study Team. Determinants of HIV Prevalence Among Female Sex Workers in Four South Indian States: Analysis of Cross-Sectional Surveys in TwentyThree Districts. *AIDS* 5(Suppl. 22): S35–S44.
- Sarkar, NN. (2008)** Barriers to condom use. *Eur J Contracept Reprod Health Care*;13(2):114–122. DOI: 10.1080/13625180802011302.
- Suratt, H., & Inciardi, J. (2010).** An effective HIV risk reduction protocol for drug-using women sex workers. *Journal of Prevention and Intervention in the Community* 38(2):118–131.
- UNAIDS (2018)** 'Miles to go: closing gaps, breaking barriers, fighting injustices. [http://www.unaids.org/sites/default/files/media\\_asset/miles-to-go\\_en.pdf](http://www.unaids.org/sites/default/files/media_asset/miles-to-go_en.pdf) Accessed 2019 May 10
- UNAIDS(2017)** [http://www.unaids.org/sites/default/files/media\\_asset/20170720\\_Databook.pdf](http://www.unaids.org/sites/default/files/media_asset/20170720_Databook.pdf)
- UNAIDS/the Global Fund/WHO (2016)** 'Key Population Atlas' as referenced in UNAIDS (2016) 'Prevention Gap Report'
- UNAIDS, (2009).** Global summary on Aids epidemic Available from: <http://www.unaids.org> Accessed 2019 May 10.
- Verma, R., Saggurti, N., Singh, A., & Swain, S. (2010):** Alcohol and Sexual Risk Behavior among Migrant Female Sex Workers and Male Workers in Districts with High In-Migration from Four High HIV Prevalence States in India. *AIDS Behav* 14: S31–39. 15.
- World Health Organization (WHO) (2013):** 'Implementing comprehensive HIV/STI programmes with sex workers: practical approaches from collaborative interventions', [https://apps.who.int/iris/bitstream/handle/10665/90000/9789241506182\\_eng.pdf;jsessionid=2A83413C5E9B140396BD52E7179B083F?sequence=1](https://apps.who.int/iris/bitstream/handle/10665/90000/9789241506182_eng.pdf;jsessionid=2A83413C5E9B140396BD52E7179B083F?sequence=1) Accessed 2019 May 10.
- Zhang, H., Liao, M., Nie, X., Pan, R., & Wang C. (2011):** Predictors of consistent condom use based on the Information-Motivation-Behavioral Skills (IMB) model among female sex workers in Jinan, China. *BMC Public Health* 11: 113.