

<https://ojshostng.com/index.php/ajbr>

Afr. J. Biomed. Res. Vol. 27 (May 2024); 243- 251

Research Article

Geospatial Analysis of Malaria Prevalence among Children Under Five Years in Jigawa State, North West, Nigeria

Yakudima I.I.¹, Adamu Y.M.², Samat N.³, Mohammed M.U.², Abdulkarim I.I.² and Hassan N.I.².

¹*Department of Geography, Aliko Dangote University of Science and Technology, Wudil, Kano, Nigeria*

²*Department of Geography, Bayero University, Kano, Nigeria*

³*Department of Geography, University Sains Malaysia.*

ABSTRACT

Malaria is one of the leading causes of illness and death in developing countries. Despite growing international concern and efforts to provide effective treatment through the development and improvement of vector control mechanisms, malaria infection continues to remain a leading health problem in Africa, particularly south of the Sahara. At present, however, very few studies have been undertaken that investigate the spatial distribution of malaria cases. This study, therefore, aims to employ geospatial tools to map out areas of high and low malaria prevalence among children under the age of five years. Annual malaria prevalence was computed using confirmed cases from 2014 to 2019 in Jigawa state. Spatial autocorrelation techniques using global Moran's I and Getis-Ord G_i^* statistics were applied. Overall malaria prevalence ranged between 2,743 and 12,916 per 100,000 populations. Results of the overall global Moran's I indicate a statistically significant degree of positive autocorrelation ($I = 0.358122$, $Z = 3.721018$, $P = 0.000198$) and all the years under study showed clustered patterns. Hotspot analysis was further explored to show the location of clusters. The results of the analysis detected high prevalence clusters in central and north-western parts of the state. The study recommends targeting hotspot areas in the design and implementation of malaria control activities.

Keywords: *Cluster, Coldspot, Hotspot, Jigawa, Prevalence, Spatial statistics.*

*Author for correspondence: Email: iaaliyu.geo@buk.edu.ng;

Received: September 2023; Accepted: March 2023

DOI: <https://doi.org/10.4314/ajbr.v27i2.7>

© 2024 The Author(s).

This article has been published under the terms of Creative Commons Attribution-Noncommercial 4.0 International License (CC BY-NC 4.0), which permits noncommercial unrestricted use, distribution, and reproduction in any medium, provided that the following statement is provided. "This article has been published in the African Journal of Biomedical Research"

INTRODUCTION

Malaria is one of the leading causes of illness and death in developing countries, especially in sub-Saharan Africa (Sachs and Malaney, 2002; WHO, 2018; Rumisha *et al.* 2019). It is a vector-borne infectious disease transmitted from one person to another through the bite of an infected female anopheles mosquito (Kim *et al.*2020). It therefore, remains one of the highest public health concerns in tropical and subtropical regions of the world, especially in South and Latin America, Africa, and Southeast and Central Asia (Prothero, 1995; WHO, 2008). Most cases occur in Sub-Saharan Africa (Carter and Mandis, 2002; Sachs and Malaney, 2002; Patz *et al.* 2005; WHO, 2014). Malaria is caused by protozoan parasites of the genus plasmodium. Human malaria is normally caused by four different species of plasmodium; they are *p. falciparum*, *p.*

malariae, *p. ovale*, and *p. vivax*. Among these species, *p. falciparum* is the most significant and deadly form of malaria presenting severe health risks, especially in sub-Saharan Africa, which is preventable (WHO, 2011; Paton, *et al.* 2021).

An estimated 3.4 billion people in 92 countries are at risk of contracting malaria, with 1.1 billion at high risk (a likelihood of obtaining malaria in a year greater than one in 1000) (WHO, 2019). A more recent report indicated that there are expected to be 241 million cases of malaria worldwide, with 627 000 malaria fatalities in 2020 (WHO, 2021) indicating an increase in the number of cases and deaths reported in 2019. The WHO African Region is responsible for a disproportionately large amount of the worldwide malaria burden. In 2020, 95 percent of malaria cases and 96 percent of malaria deaths occurred in the region (WHO, 2021). In 2017, it was estimated that there were 219 million clinical malaria

cases worldwide. 92% of the cases occurred in the WHO African region. Global malaria-related deaths in 2017 were estimated at 435,000, of which 93% were in Africa, and children under the age of five years are particularly high risk accounting for 61% of all deaths (WHO, 2018). According to a WHO report in 2013, more than 90% of the population in Africa lives in areas exposed to the risk of malaria infection, with 74% of the population living in highly endemic areas where malaria transmission is greatest and perennial (WHO, 2014).

The vector *Anopheles* mosquito is generally influenced by environmental conditions. Climate, through the influence of factors such as temperature, precipitation, and relative humidity, has a significant influence on malaria transmission (Dahal, 2008; Oluleye and Akinbobola, 2010). It influences the development, reproduction, and survival of *Anopheles* mosquitoes and malaria parasites. Other environmental factors including altitude, topography, land use/land cover, and human settlement patterns have a significant impact on the spatial distribution of vectors and the disease they transmit (Abeku *et al.* 2004; Van Der Hoek *et al.* 2003; Kibret *et al.* 2019).

The malaria situation in Africa is becoming more and more complex as the disease is now extended to include previously non-endemic or low transmission areas of the eastern and southern parts of the continents (Chala, 2007). In fact, malaria globally is responsible for about 7.8 % disease burden (WHO, 2021) with Africa accounting for the majority. A recent study reveals that malaria transmission is higher in rural areas than in urban areas (Doumbe-Belisse, *et al.* 2021). Concerning Nigeria, malaria is endemic in all parts of the country, with year-round transmission and the rates of Transmission differ from place to place being slightly lower in Sahelian areas and high mountain areas of the plateau (National Population Commission, National Malaria Control Programme and ICF International, 2012). Nigeria accounted for 25% and 19% of the global estimate of malaria cases and deaths in 2017 (WHO, 2018). The system for control and prevention of malaria now faces major challenges in endemic countries like Nigeria due wide range of factors. Among the factors are increasing resistance of the parasite to anti-malarial drugs, the resistance of mosquito to insecticides, poor coverage of health infrastructure for diagnosis and treatment of the disease especially in rural areas, shortage of well-trained personnel, scarce financial resources, high cost of the control program and weak strategic planning effective malaria control (Traore, 2003; Alemu, 2006; Ranson 2018).

In all malaria-endemic regions, malaria tends to form a cluster known as “hot spots” and “hot populations” that become sources of continued infections (Rulisa *et al.* 2003; Shahandeh and Basseri, 2019). Active and prompt identification of clusters and specific locations of clusters, as well as associated risk factors, is essential for assisting government and private partners interested in malaria control and prevention to provide appropriate intervention strategies and resource allocation (WHO, 2007).

Recent development in geographical information system (GIS) and spatial statistical tools provide new methodologies to analyze the generated clinical data by detecting spatial patterns of disease distribution and delineation of hot spots to

assess the true situation for better public surveillance and for improving our understanding of the transmission dynamics of disease such as malaria (Osterholt *et al.* 2006; Yeshiwondim *et al.* 2009; Saxena *et al.* 2009). Geospatial statistics have been increasingly applied in analyzing and mapping malaria case data (Gebreslasie, 2015; Kirk, *et al.* 2015; Ahmad, Goparaju, and Qayum, 2017; Zandian *et al.* 2019; Gambo, *et al.* 2020). They provide an opportunity for medical geographers to better understand and explain the spatial distribution of diseases and the possible association between risk factors. Spatial autocorrelation statistics analyze the correlation of a variable in relation to the location that that variable is. Moran’s I is a more popular tool for measuring spatial autocorrelation (Huo *et al.* 2012). Global Moran’s I measure and analyze the degree of dependency among observations in geographic space. On the other hand, local spatial autocorrelation statistics such as local Moran’s I and Getis and Ord are useful in identifying locations of spatial clusters or “Hot spots” and outliers (Huo *et al.* 2012).

Currently, despite the endemic situation of malaria in Jigawa State, the application of GIS and spatial statistics to study malaria and provide information to assist in the control and elimination of the disease in the state is sparse in published literature. Thus, this study applied GIS and spatial statistical techniques on confirmed malaria cases to identify spatio-temporal patterns of hotspots of malaria cases in Jigawa state.

MATERIALS AND METHODS

Study Area: Jigawa State is located in the northwest geopolitical zone of Nigeria, consisting of 27 local government areas. The state lies roughly between latitude 10° N and 12° N and longitude 7° E to 10° E. It shares its western border with Kano and Katsina States, Bauchi State to the east, and Yobe State to the northeast. To the north, Jigawa State shares an international border with the Niger Republic. It covers an approximate land area of about 22, 410 Km² (Fig 1). The study area experiences a tropical wet and dry type of climate coded as “Aw” in the Koppen classification system




Figure 1: Jigawa State (the study area)

. The wet season commences from June through September, although mid-May and mid-October are usually regarded as part of the wet season. The normal annual rainfall is higher in the southern parts, where it is between 1000 and 1100 mm and decreases to about 500 to 550 mm as one moves to the north and north-eastern parts of the state (Olofin, 2008). Maximum temperatures (above 42° C) are recorded during the hottest months of March and April and may extend to mid-May, while lower temperatures as low as 10o C are normally recorded in the coolest months (December to February). The mean annual temperature is about 26° C, but the mean monthly values range between 21° C and 23° C in the coolest months and over 30° C in the hottest months (Olofin, 2008). The Hadejia river system forms the major drainage accounting for about 80% of the historic flow of the river system in the area (Abdulhamid, 2014) and drains essentially north-eastwards into Lake Chad. However, in some parts of the north, the river ceases to flow in a proper channel but wanders across the formation. As a consequence, the area gets flooded in the rainy seasons, which becomes a breeding site for malaria vectors.

Jigawa state ranked 8th among the most populous states in Nigeria based on the 2006 population census. Its population stood at 4,348,649 (Federal Republic of Nigeria, 2007) with an average density of about 194 persons/km². The projected population figure for the area as of 2019, stood at 6, 526, 432 (NPC, 2020). Agriculture, which is regarded as one of the occupational malaria risk factors, is the dominant economic activity in the state employing 70% of the inhabitants (Jigawa State Ministry of Health {JSMoH}, 2010).

Malaria and Population Data: Data on malaria were obtained from the Jigawa State Ministry of Health. The data includes all malaria cases among under-five years old, confirmed by microscopic or Rapid Diagnostic Test (RDT) kits, and clinically diagnosed at various public health facilities across the 27 LGAs. Malaria cases were recorded daily for each facility and later transferred to National Health Management Information System (NHMIS) monthly summary form for health facilities version 2013 at the local government level for onward submission to the state ministry of health. Monthly malaria cases for the period of six years (2014 - 2019) were extracted from the database of the ministry. The choice of this period was based on data availability for the entire state.

Data on population were collected from National Population Commission, Jigawa state. Data from population censuses conducted in 1991 and 2006 were collected and used to compute the inter-census growth rate for each local government area of the state. The population was projected for the years 2014 to 2019 and the data were used to calculate malaria prevalence rates and produce hotspot maps.

Prevalence Rate: Malaria case data were standardized as prevalence rather than total cases. The essence was to give a better representation of the data due to the varying population size of the LGAs in the state. Prevalence is one of the measures of disease frequency and burden. It is often useful to assess the burden of disease in a given population. This is not limited to burden in relation to resources; it also reflects burden in terms of life expectancy, morbidity, quality of life, or other

indicators (Noordzij, *et al.* 2010). For this study, malaria prevalence data were generated by dividing the total number of malaria cases for each year of study for the 27 LGAs by the population of each LGA. The following formula was used to compute the prevalence rate:

$$\text{Prevalence rate} = \frac{\text{Total Number of Malaria cases}}{\text{Total Population of the area}} \times 100,000 \dots (1)$$

Cluster Analysis: Spatial and temporal cluster analyses were conducted to detect spatial clustering of malaria prevalence using the spatial autocorrelation technique. The technique represents the correlation between a variable at a location and the values of the same variable at a neighbouring location (Zhang *et al.* 2009). Spatial autocorrelation statistics measure and analyze the degree of similarity among observations in geographic space. Two forms of analysis were employed. The first was to determine the degree of the spatial pattern of malaria cases in the area, which may either be clustered; dispersed, or random using Global Moran's I. The global Moran's I statistics provide a summary for each feature in the dataset consisting of I, z-score, and p-value. The tool compares the values of neighbouring locations and strong positive autocorrelation (clustered) existed when neighbouring areas over the entire area under study have similar values. If neighbouring areas have dissimilar values, then the tool yielded strong negative spatial autocorrelation (dispersed). The value of I, therefore, ranges from +1 (indicating positive autocorrelation) to -1 (representing negative autocorrelation). A 'Z' value is computed to assess whether clustering/dispersing is statistically significant or otherwise. The statistical test, therefore, is capable of determining the degree of spatial autocorrelation of a given phenomenon. It measures the degree to which a phenomenon is clustered in space (Osayomi, 2015). However, the tool is not powerful to indicate the location of clusters and outliers. The formula for Moran's I statistics is given below:

$$I = \frac{N}{\sum_i \sum_j W_{ij}} \frac{\sum_i \sum_j W_{ij} (X_i - \bar{X})(X_j - \bar{X})}{\sum_i (X_i - \bar{X})^2} \dots (2)$$

Where N is the number of spatial units (LGAs) indexed by *i* and *j*. X represents the study variable (malaria prevalence

rate), \bar{X} is the mean value of malaria cases. W_{ij} is an element of a matrix of spatial weights which defines the degree of proximity between local governments areas *i* and *j*.

The second analysis is to show the location of hotspots/cold spots using Local Getis-Ord statistics (G_i^*). The Getis-Ord G_i^* statistical analysis, on the other hand, was employed to supplement Moran's I. The tool identifies different spatial clustering patterns like hot spots and cold spots, with statistical significance (Getis and Ord, 2010). The statistics reflect whether differences between the local mean (i.e., cases for a local government and it's nearest neighbouring local government in the state) were significantly different from the global mean (i.e., the overall cases of all local governments in the state). The resultant "z" score is an indicator of a high or low malaria prevalence cluster. High and positive z-score

values above 1.96 indicate the occurrence of malaria hotspots, while values below 1.96 represent cold spot areas. Therefore, the higher/lower the z-score is, the stronger or weaker the relationship would be. Scores closer to zero indicate the absence of clusters. The formula for calculating hot spots using Getis-ord G_i^* is given as:

$$I_i = \frac{(y_i - \bar{y}) \sum_{j=i}^n w_{ij} (y_j - \bar{y})^2}{m_2} \dots\dots\dots (3)$$

Where y_i is the value of the variable (malaria prevalence rate) at the location i th, n is the number of LGAs, w_{ij} is a weight representing the spatial relationship of LGAs i , and j , m_2 is the average of the squared deviations from the mean of malaria cases.

RESULTS AND DISCUSSION

Descriptive Statistics: The statistical summary of malaria prevalence over the period 2014 - 2019 is presented in Table 1. According to the Table, malaria prevalence among children below the age of five years ranged from 2743/100,000 population to 12,916/100,000 with an average rate of 5952/100,000 individuals. The standard deviation of malaria prevalence ($\delta = 2261.92$) which measures the degree of dispersion of cases from the mean, shows that there is wide variation in the prevalence rate of malaria under five years.

Table 1: Descriptive statistics of malaria prevalence among under-five years old

Mean	Minimum	Maximum	Std. Deviation
5952	2743	12,916	2261.92

Source: JSMoH, 2020 and Author’s analysis, 2021

The mean annual prevalence rate among those under-five years (5,952 per 100,000 or 5.95%) observed in this study is higher than 1.8% and 1.92% reported by Dawaki *et al.* (2016) and Solomon *et al.* (2020) respectively. However, our result (5.95%) is close to that of Alemu *et al.* (2012) and Nanvyat *et al.* (2017) who found 7.8% and 7.4% in Kola Diba, North Gondar, Ethiopia, and Jos Plateau State, Nigeria. In contrast, other studies have reported a high prevalence of 27.7% by Elechi *et al.* (2015) in Maiduguri, Borno State, Nigeria; 61.7% by West and Okari (2018) in Port Harcourt, Nigeria and 63% by Simon-Oke *et al.* (2019) in Ekiti State, Nigeria. Similarly, other high prevalence rates were reported in Africa and Asia. They include: 11.97% in Tanzania (Paul and Msengwa, 2018), 22.1% in Ghana (Yankson *et al.* {2019}), 22.1% in Ethiopia (Abossie *et al.* {2020}), 37.4% in Malawi (Gaston and Ramroop, 2020), 35.4% in Malawi (Chilanga *et al.* {2020}) and 42.3% in Indonesia (Jiero and Pasaribu, 2021). This variation in the results might be due to differences in climatic conditions, altitudinal variations, type of the study design employed, variations in the methods of malaria detection, differences in the coverage of intervention activities, and other factors that affect transmission.

The low prevalence rate among under-five children reported in this study might be related to increased awareness of the inhabitant of the state on the use of ITNs. This finding is not surprising considering the 2018 Nigeria Health Demography Survey Report, which revealed that household ownership of ITNs in the country is highest in Jigawa (the study area) and Kebbi States (98%) which is above the national target (NPC and ICF International, 2019). The report further disclosed that in Jigawa State, 90.5% of the surveyed children who are below five years of age slept under ITNs the night before the survey. This shows that the vast majority of the younger children are well protected from the infective bites by mosquitoes.

Spatial Patterns of Malaria Prevalence: Global spatial statistics using the Moran’s I measure were used to test for the significant pattern in the prevalence of malaria among under-five children in the study area. The test results (Table 2) showed overall positive and statistically significant Moran’s I (Moran’s I = 0.358122, Z score = 3.721018. P = 0.000198), suggesting a significant concentration of high prevalence rates (clustering). The result in Table 2 further shows a similar level of autocorrelation for the years under study. The clustering was strongest in the year 2018 (Moran’s I = 0.406459, Z score = 4.023251. P = 0.000057) followed by 2019 (Moran’s I = 0.310412, Z score = 3.263420, P = 0.001101).

Table 2: Spatial pattern of malaria prevalence

Years	Moran’s Index	Z-score	P value	Pattern
Under-five years:				
2014	0.235427	2.614904	0.008925	Clustered
2015	0.214774	2.461729	0.013827	Clustered
2016	0.237812	2.501937	0.012352	Clustered
2017	0.287145	2.914179	0.003566	Clustered
2018	0.406459	4.023251	0.000057	Clustered
2019	0.310412	3.263420	0.001101	Clustered
2014-2019	0.358122	3.721018	0.000198	Clustered

Source: Author’s computation (2021)

The presence of significant positive autocorrelation for the years under study shows the extent to which neighbouring rates are correlated. Thus, LGAs sharing a border is more similar with respect to malaria than those that are distantly apart. However, the occurrence of malaria prevalence in these years in specific areas is not by chance. The statistically established spatial dependency of the disease implies the presence of similar risk factors in neighbouring areas thereby influencing the spatial transmission of the disease. This finding is not surprising since malaria is an environmental disease, and environmental variables are spatially related. Other similar studies (Sexena *et al.* 2012; Osayomi, 2014; Osei and Yibile, 2015; Yakudima and Adamu, 2019; Bizimana and Nduwayezu, 2020) had also revealed spatial clustering of malaria cases in their studies. However, these global test results indicate a need for further investigation using local spatial statistics (hotspot analysis).




Figure 2:
Hotspot of malaria prevalence (2014-2019)

Hotspot Analysis of Malaria for Children Under Five Years: Local spatial statistics using Getis-Ord G_i^* was applied to detect the locations of clusters over the study periods. Figure 2 represents the result of the cumulative (2014 – 2019) hotspot analysis for the under-five years population group. The figure showed central and north-western parts of the state as the hotspot regions. Hotspots of 90% and 95% confidence levels were noted in the following LGAs: Kiyawa, Jahun, Dutse, Gwiwa, Roni, Kazaure, and Yan-kwashi. North-eastern part of the state (comprising Kaugama, Auyo, Hadejia, Malam Madori, Kiri Kasamma, Guri, and Birniwa LGAs) on the other hand formed the cold spot region.

The year-wise hotspot analysis results (figure 3a) showed a hotspot cluster at all three confidence levels in 2014. LGAs like Gwiwa, Roni, Yan-kwashi, Kazaure, and Kiyawa LGA formed the hotspot clusters. In 2015, only three LGAs (Jahun, Gwiwa, and Roni) showed hotspots at 95% and 90% confidence levels (figure 3b). For the remaining years of the study (2016-2019) four LGAs exhibit hotspot clusters (figures 3c-3f). Gwiwa, Roni, and Kiyawa LGAs appear as hotspot areas at 99% confidence level for two years of the analysis, while four LGAs including Birnin Kudu, Buji, Kazaure, and Yan-kwashi appear in the group (99% C.I.) in only one year. For 95% confidence level Dutse LGA consistently appears for the three years of the analysis, Jahun LGA appears in two years while Yan-kwashi, Kazaure, and Kiyawa each appeared in only one year. Six LGAs formed hotspot clusters at 90% confidence level, they are Kiyawa, Gwiwa, Roni, Jahun, Ringim, and Birnin Kudu, and each appeared in only one year. On the other hand, the overall result (2014-2019) of hotspot analysis established a cold spot cluster in the north-eastern

part of the state with Guri, Birniwa, and Kiri Kasamma at 99% confidence interval, while Kaugama, Malam Madori, and Hadejia fall under 95% confidence level (figure 2). Auyo LGA belongs to the cold spot at 90% confidence interval (figure 2). The yearly results show that Birniwa LGA consistently emerges as a significant cold spot (99%) for five consecutive years, Kiri Kasamma appears in three years, Malam Madori and Hadejia two years each, while Guri and Kaugama each appeared in only one year (figure 3a-3f). At 95% confidence level Guri LGA occurs under a cold spot for five years, Kiri Kasamma and Kaugama (3 years), Malam Madori (2 years), while Hadejia, Auyo, and Birniwa had emerged in only one year.

Identification of a hotspot cluster consistently located in the central and the north-western parts of the state was the most significant outcome of cluster analysis. A possible explanation for the consistent clustering of malaria in these areas could be the presence of favourable conditions for malaria transmission. Some LGAs like Jahun and Kiyawa shared borders with other LGAs that have wetlands where traditional rice cultivation is practiced. Rice fields according to Gurthmann *et al.* (2002) are the most favourable sites for the mosquito to breed. In addition, most of the LGAs that fall within and surrounding the clusters have water streams that allow for the cultivation of vegetables and other crops. These water agro-systems may provide significant habitat for mosquito breeding and thus, increase the vector population for malaria transmission. This finding is confirmed by the works of Rulisa *et al.* (2013) and Bizimana & Nduwayezu (2020) that discovered significant malaria hotspots located close to water-based agro-ecosystems.




Fig 3a: Hotspot of malaria prevalence (2014)




Fig3b: Hotspot of malaria prevalence (2015)




Fig 3c: Hotspot of malaria prevalence (2016)




Fig 3d: Hotspot of malaria prevalence (2017)




Fig 3e: Hotspot of malaria prevalence (2018)




Fig 3f: Hotspot of malaria prevalence (2019)

The second likely factor is urbanization. Jigawa State is made up of five emirate districts. The headquarters of these districts are considered the major towns in the state and are therefore providing employment opportunities and other services, hence attracting people from other areas. Three out of the five emirate headquarters were part of the identified clusters. Osayomi (2014) established an association between malaria and urbanization in Nigeria. This association was due to the fact that most towns are characterized by poor sanitary conditions, such as improper disposal of waste, poor condition of drainage, and an unclean housing environment among

others. All these encourage the breeding of mosquitoes and thus, promote the transmission of malaria.

The availability of formal healthcare centres in these 11 LGAs may partly explain the emergence of hotspot clusters in the areas. This finding was supported by a study on the distribution of healthcare facilities in Jigawa state by Yakudima (2018) who found that nearly 60% of modern health facilities in the state are concentrated in these areas. Another likely reason for a stable hotspot in these areas is that the existing malaria control and intervention measures might have not been taken correctly or might have not been applied appropriately.

Table 3:
Number of LGAs per type and intensity of malaria cluster

Confidence Interval	2014	2015	2016	2017	2018	2019	2014-2019
Under-five years:							
Coldspots 99% C.I.	1	3	1	1	5	3	3
Coldspots 95% C.I.	2	-	5	4	2	3	3
Coldspots 90% C.I.	-	-	1	1	1	1	1
Not significant	19	21	16	17	15	16	13
Hotspots 90% C.I.	1	2	2	1	-	-	6
Hotspots 95% C.I.	2	1	2	2	1	-	1
Hotspots 99% C.I.	2	-	-	1	3	4	-

Source: Author's computation (2021)

Our analysis further showed that the coldspots are majorly concentrated in the north-eastern part of the state. This area is characterized by low rainfall and high temperature. Adequate rainfall is required to recharge existing water bodies or create new ones which serve as breeding sites. The low rainfall receives in this area makes mosquito breeding habitats so scarce thus, low transmission intensity of malaria. Temperature is also considered a key determinant of transmission. This area (north-east) is associated with high temperatures for most of the year. The high temperature at 40oc reduces mosquito abundance due to the long larval duration (Dale *et al.* {2005}). Low rainfall coupled with high temperature affects the development of vegetation cover in the area. Adequate vegetation cover provides favourable conditions for malaria vectors to rest. The sparse nature of vegetation cover in the area can reduce mosquito density. All these factors could partly explain the low transmission of malaria in the area leading to cold spot conditions.

Table 3 summarized the number of LGAs identified as hotspots and coldspots for each year. From the table, the number of LGAs reported as hotspots cluster across the state was highest in the year 2014 and slightly lower during the 2016-2019 periods.

High-intensity clustering was reported in 2018 and 2019 in three and four LGAs respectively. This pattern of high-intensity clustering in these years could be related to high rainfall which brings about serious flooding in many parts of the state. Thus, stagnant water points tend to occur in many places and favoured ecological conditions suitable for malaria pathogens and vectors to develop and proliferate. These findings corroborate the works of Okunola and Oyeyemi, (2019) who reported malaria clusters in the 6 geopolitical zones of Nigeria, and Gambo, *et al.* (2020) who reported environmental influences, for example, Climate change, which has a favourable impact on the vector's ecology, habitat, and breeding grounds, as well as the study area's geographical position and poorly coordinated settlement pattern and environmental cleanliness, to have all contributed to an increase in malaria cases in Nigeria.

In this study, the spatial autocorrelation technique has been explored to identify and map the geographical distribution of malaria prevalence clusters. Further, the study highlighted the role of GIS and spatial data analysis in

explaining disease dynamics. The results discovered statistically significant clustering of high malaria prevalence in the central and north-western parts of the state while coldspot LGAs were persistently detected in the north-eastern part of the area. This finding suggests the provision of adequate malaria intervention services including vaccination to the identified high malaria prevalence clusters.

Acknowledgment:

We would like to thank the ministry of health, Jigawa State Nigeria, for providing malaria incidence data used in writing this paper

REFERENCES

- Abdulhamid, A. (2014)** Drainage, Hydrology and Water Resources. In Tanko, A. I. And Momale S. B. (eds) 2014: KANO – Environment, Society and Development. Adonis and Abbey Publishers Ltd UK pp 21-34
- Abeku, T. A., De Vlas, S. J., Borsboom, G. J. J. M., Tadege, A., Gebreyesus, Y., Gebreyohannes, H., ... & Habbema, J. D. F. (2004).** Effects of meteorological factors on epidemic malaria in Ethiopia: a statistical modeling approach based on theoretical reasoning. *Parasitology*, 128(6), 585-593.
- Abossie, A., Yohanes, T., Nedu, A., Tafesse, W., & Damitie, M. (2020).** Prevalence of malaria and associated risk factors among febrile children under five years: a cross-sectional study in Arba Minch Zuria district, South Ethiopia. *Infection and Drug Resistance*, 13, 363-372
- Ahmad, F., Goparaju, L., & Qayum, A. (2017).** Studying malaria epidemic for vulnerability zones: Multi-criteria approach of geospatial tools. *Journal of Geoscience and Environment Protection*, 5(5), 30-53.
- Alemu, A., Muluye, D., Mihret, M., Adugna, M., & Gebeyaw, M. (2012).** Ten-year trend analysis of malaria prevalence in Kola Diba, North Gondar, Northwest Ethiopia. *Parasites & vectors*, 5(1), 1-6.
- Alemu, G. (2006)** Prevalence of malaria and its influencing factors in Awassa district Southern Ethiopia. M Sc thesis Addis Ababa University, Ethiopia
- Bizimana, J.P. and Nduwayezu, G. (2020)** Spatio-temporal patterns of malaria incidence in Rwanda. *Transaction in GIS*, 00:1-17. <https://doi.org/10.1111/tgis.12711>

- Bosman, A. and Teklehaimanot, A. (1997)** Malaria control strategies in Africa. In Carosi, G., and Casteli, F. (ed): Handbook of malaria infection in the tropics. Health co-operation papers – 15 pp 217-226
- Carter R. and Mendis, K. N. (2002)** Evolutionary and historical aspect of the burden of malaria. *Clin. Microbial Rev.* 15: 564-594
- Chala, B. (2007)** Assessment of malaria as a public health problem in Finchaa sugar factory based on clinical records and parasitological surveys. M Sc thesis School of graduate studies, Addis Ababa University
- Chilanga, E., Collin-Vezina, D., Maclntosh, H., Mitchell, C. and Cherney, K. (2020)** Prevalence and determinants of malaria infection among children of local farmers in Central Malawi. *Malaria Journal*, 19:308
- Dahal, S. (2008).** Climatic determinants of malaria and kala-azar in Nepal. *Regional Health Forum*, 12:32-37.
- Dale, P., Sipe, N., Anto, S., Huntajulu, B., Ndoen, E., Papayungan, M., Salikhu, A. and Parabowo, T. (2005).** Malaria in Indonesia: a summary of recent research into its environmental relationships. *Southeast Asian Journal of Tropical Medicine and Public Health.* 36(1): 1-13.
- Dawaki, S., Al-Mekhlafi, H. M., Ithoi, I., Ibrahim, J., Atroosh, W. M., Abdulsalam, A. M., ... & Lau, Y. L. (2016).** Is Nigeria winning the battle against malaria? Prevalence, risk factors, and KAP assessment among Hausa communities in Kano State. *Malaria Journal*, 15(1), 1-14.
- Doumbe-Belisse, P., Kopya, E., Ngadjieu, C. S., Sonhafou-Chiana, N., Talipouo, A., Djamouko-Djonkam, L., ... & Antonio-Nkondjio, C. (2021).** Urban malaria in sub-Saharan Africa: dynamic of the vectorial system and the entomological inoculation rate. *Malaria Journal*, 20(1), 1-18.
- Elechi, H. A., Rabasa, A. I., Muhammad, F. B., Garba, M. A., Abubakar, G. F., & Umoru, M. A. (2015).** Prevalence and pattern of malaria parasitaemia among under-five febrile children attending paediatric out-patient clinic at University of Maiduguri Teaching Hospital, Maiduguri. *Nigerian Journal of Paediatrics*, 42(4), 319-324.
- Federal Republic of Nigeria Official Gazette, (2007):** No.24 Vol. 94
- Gambo, J., Shafri, H. Z. B. M., Abubakar, A., Yusuf, Y. A., & Adam, M. B. (2020).** Comparative Analysis of Malaria Cases Using Geospatial-statistical Approach in Hadejia Metropolis, Jigawa State, Nigeria. In IOP Conference Series: Earth and Environmental Science (Vol. 540, No. 1, p. 012035). IOP Publishing
- Gaston, R. T., & Ramroop, S. (2020).** Prevalence of and factors associated with malaria in children under five years of age in Malawi, using malaria indicator survey data. *Heliyon*, 6(5), e03946.
- Gebreslasie, M. T. (2015).** A review of spatial technologies with applications for malaria transmission modelling and control in Africa. *Geospatial health*, 10(2).
- Getis, A., & Ord, J. K. (2010).** The analysis of spatial association by use of distance statistics. In *Perspectives on spatial data analysis* (pp. 127-145). Springer, Berlin, Heidelberg.
- Guthmann, J.P, Llanos-Cuentas, A., Palacios, A., Hall, A.J. (2002)** Environmental factors as Determinants of Malaria Risk. A descriptive study on the Northern coast Peru. *Tropical Medicine and International Health*, 7(6):51-525
- Huo, X., Li, H., Sun, D., Zhou, L., & Li, B. (2012)** Combining geostatistics with moran's I analysis for mapping soil heavy metals in Beijing, China. *Int. J. Environ. Res. Public Health*, 9: 995-1017
- Jiero, S. & Pasaribu, A.P. (2021)** Haematological profile of children with malaria in Sorong, west Papua, *Indonesia. Malaria Journal*, 20:126
- Jigawa State Ministry of Health {JSMoH}(2010)**Strategic Health Plan (2010-2015)
- Kibret, S., Glenn Wilson, G., Ryder, D., Tekie, H., & Petros, B. (2019).** Environmental and meteorological factors linked to malaria transmission around large dams at three ecological settings in Ethiopia. *Malaria journal*, 18(1), 1-16.
- Kim, H. Miller, F.D., Hernandez, A., Tanser, F. Mogeni P. & Cuadros. D.F. (2020):** Spatiotemporal analysis of insecticide-treated net use for children under 5 in relation to socioeconomic gradients in Central and East Africa. *Malar J* (2020) 19:163
- Kirk, K. E., Haq, M. Z., Alam, M. S., & Haque, U. (2015).* Geospatial Technology: a tool to aid in the elimination of malaria in Bangladesh. *ISPRS International Journal of Geo-Information*, 4(1), 47-58.
- National Population Commission {NPC}(2020)** Jigawa state population projection
- Nanvyat, N., Mulambalah, C.S., Ajiji, J.A., Dakul, D.A. & Tsingalia, M.H. (2017)** Prevalence of human malaria infection and its transmission pattern in the highlands and lowlands of Plateau State, Nigeria. *Indian Journal of Science and Technology*, 10(32):1-9
- National Population Commission (NPC){Nigeria} & ICF International (2019)** Demographic and Health Survey 2018. Abuja Nigeria and Rockville, Maryland, USA: NPC and ICF International
- National Population Commission (NPC){Nigeria}, National Malaria Control Programme (NMCP){Nigeria} and ICF International (2012)** Nigeria malaria indicator survey. Abuja Nigeria: NPC, NMCP and ICF International
- Noordzij, M., Dekker, F.W., Zoccali, C. & Jager, K.J. (2010)** Measures of disease frequency: prevalence and incidence. *Nephron Clinical Practice*, 115:c17-c20
- Oesterholt, M. J. A. M., Bousema, J. T., Mwerinde, O. K., Harris, C., Lushino, P., Masokoto, A., ... & Drakeley, C. J. (2006).** Spatial and temporal variation in malaria transmission in a low endemicity area in northern Tanzania. *Malaria Journal*, 5(1), 1-7.
- Okunlola, O. A., & Oyeyemi, O. T. (2019).** Spatio-temporal analysis of association between incidence of malaria and environmental predictors of malaria transmission in Nigeria. *Scientific Reports*, 9(1), 1-11.
- Olofin, E.A. (2008)** The physical setting. In Olofin, E.A., Nabegu, A.B. & Dambazau, A.M. (eds). *Wudil within Kano region: a geographical synthesis*. A publication of the department of Geography, Kano University of Science and Technology Wudil. Adamu Joji Publishers Kano City
- Oluleye, A. & Akinbobola, A. (2010).** Malaria and pneumonia occurrence in Lagos, Nigeria: Role of temperature and rainfall. *African Journal of Environmental Science and Technology* 4: 506-516.

- Osayomi, T. (2015)** Spatio–Temporal Cluster analysis of Diabetes mellitus in Oyo state, Nigeria (2000-2014). *The Nigerian Geographical Journal*, 10(2):191-179
- Osei, F.B. & Yibile, M.M. (2015)** Geographic patterns of malaria in the Brong Ahafo Region of Ghana. *Austin Journal of Public Health and Epidemiology*, 2(2):1020
- Paton, R. S., Kamau, A., Akech, S., Agweyu, A., Ogero, M., Mwandawiro, C., ... & Snow, R. W. (2021).** Malaria infection and severe disease risks in Africa. *Science*, 373(6557), 926-931.
- Patz, J.A Campbell-Lendrum, D., Holloway, T. & Foley, J.A. (2005)** Impact of regional climate change on human health. *Nature*, 438, 17: 310-318
- Paul, E. & Msengwa, A.S. (2018)** Prevalence and socio-demographic factors associated with malaria infection among children under five years in Tanzania. *Journal of Public Health and Epidemiology*, 10(11):387-394
- Prothero, R.M. (1995)** Malaria in Latin America: Environmental and human factors. *Bulletin of Latin American Research*, 14 (3): 357-365
- Ranson, H. (2018).** Malaria: Key challenges and potential solutions. K4D Reading Pack. Brighton, UK: Institute of Development Studies.
- Rietveld, A. E. C. & Kouznetsov, R. L. (1997)** Epidemiology of Human malaria plasmodium. In Carosi, G. And Castelli, F. (eds) Hand book of malaria infection in the Tropics. Health co-operation papers No.15 pp 39-52
- Rulisa, S., Kateera, F., Bizimana, J. P., Agaba, S., Dukuzumuremyi, J., Baas, L., ... & de Vries, P. J. (2013).** Malaria prevalence, spatial clustering and risk factors in a low endemic area of Eastern Rwanda: a cross sectional study. *PloS one*, 8(7), e69443.
- Rumisha, S. F., Shayo, E. H., & Mboera, L. E. (2019).** Spatio-temporal prevalence of malaria and anaemia in relation to agro-ecosystems in Mvomero district, Tanzania. *Malaria Journal*, 18(1), 1-14.
- Sachs, J. & Malaney, P. (2002)** The economic and social burden of malaria. *Nature*, 415(6872): 680-685.
- Saxena, R., Nagpal, B. N., Das, M. K., Srivastava, A., Gupta, S. K., Kumar, A., ... & Baraik, V. K. (2012).** A spatial statistical approach to analyze malaria situation at micro level for priority control in Ranchi district, Jharkhand. *The Indian journal of medical research*, 136(5), 776.
- Saxena, R., Nagpal, B.N., Srivastava, A., Gupta, S.K. & Dash, A.P. (2009)** Application of spatial technology in malaria research and control: some new insights. *Indian J. Med. Res.* 130: 125-132
- Shahandeh, K., & Basseri, H. R. (2019).** Challenges and the path forward on malaria elimination intervention: a systematic review. *Iranian Journal of Public Health*, 48(6), 1004.
- Simon-Oke, I.A., Ogunsemi, M.F., Afolabi, O.J. & Awosolu, O.B. (2019)** Prevalence of malaria parasites among pregnant women and children under five years in Ekiti State, southwest Nigeria. *Journal of Biomedicine and Translational Research*, 5(1):5-11
- Solomon, A., Kahase, D., & Alemayehu, M. (2020).** Trend of malaria prevalence in Wolkite health center: an implication towards the elimination of malaria in Ethiopia by 2030. *Malaria journal*, 19(1), 1-8.
- Traore, C. (2003)** Epidemiology of malaria in a haloendemic area of rural Burkina Faso. Ph D Dissertation Ruprecht-karls – University Heidelberg institute of Hygiene
- Van Der Hoek, W., Konradsen, F., Amerasinghe, P. H., Perera, D., Piyaratne, M. K., & Amerasinghe, F. P. (2003).** Towards a risk map of malaria for Sri Lanka: the importance of house location relative to vector breeding sites. *International journal of epidemiology*, 32(2), 280-285.
- West, B.A. and Okari, T.G. (2018)** Prevalence of malaria in febrile under five children in Port Harcourt, Nigeria. *Journal of Dental and Medical Sciences*, 17(12):46-51
- WHO (2007)** Malaria elimination: a field manual for low and moderate endemic countries. Geneva.
- WHO (2008)** World Malaria Report, 2008. Geneva, World Health Organization.
- WHO (2014)** World Malaria Report, 2014. Geneva, World Health Organization.
- WHO (2018)** World Malaria Report, 2018. Geneva, World Health Organization.
- WHO (2019)** The Global Health Observatory, The Global Health Observatory, Geneva, World Health Organization.
- WHO (2020)** World Malaria Report, 2020. Geneva, World Health Organization.
- WHO (2021)** World Malaria Report, 2021. Geneva, World Health Organization.
- World Health Organisation (2002)** Roll Back Malaria. Fact sheet 2. www.rbm.who.int
- World Health Organization.** World malaria report 2021. <https://www.who.int/teams/global-malaria-programme/reports/world-malaria-report-2021> (Accessed on February 16, 2022)
- Yakudima, I.I. (2018)** Distribution pattern of healthcare services in Jigawa state, Northwest, Nigeria. In Nabegu, A.B., Na'ibbi, A.I. and Garba, A.S. (eds) Northwest A Book of Readings. Ahmadu Bello University Press Limited, Zaria, Kaduna State, Nigeria. Pp 227-244
- Yakudima, I.I. & Adamu Y.M. (2019)** Spatio-temporal clustering of malaria cases among children in Kano State, Nigeria. *Wudil Journal of Pure and Applied Sciences*, 1(2):10-22
- Yankson, R., Anto, E.A., & Chipeta, M.G. (2019)** Geostatistical analysis and mapping of malaria risk in children under 5 using point-referenced prevalence data in Ghana. *Malaria Journal*, 18:67
- Yeshiwondim, A. K., Gopal, S., Hailemariam, A. T., Dengela, D. O., & Patel, H. P. (2009).** Spatial analysis of malaria incidence at the village level in areas with unstable transmission in Ethiopia. *International Journal of Health Geographics*, 8(1), 1-11.
- Zandian, E., Nodez, S., Khosravani, M., Rafatpanah, A. & Latifi, R. (2019)** GIS determination of malaria hot spots in Qeshm Island. *Journal of Zoonotic Diseases & Public Health*, 3(1):1-5
- Zhang, L., Ma, Z. & Guo, L. (2009)** An evaluation of spatial autocorrelation and heterogeneity in the residuals of six regression models. *Forest Sciences*, 55(6): 533-48.