



*Afr. J. Biomed. Res. 13 (January 2010) 55 - 59*

*Research article*

## **Anti-hyperglycemic Activity of a Polysaccharide fraction from *Lycium barbarum***

**Guowen Cui<sup>1</sup>, Longjun Jing<sup>2\*</sup>, Qiang Feng<sup>2</sup>, Yuanshi Xiao<sup>2</sup>, Ramesh Putheti<sup>3</sup>**

<sup>1</sup>*Yuxi Normal University, Yuxi, Yunnan Province, 653100, P.R. China.*

<sup>2</sup>*Qujing Normal University, Qujing, Yunnan Province, 655011, P.R. China.*

<sup>3</sup>*10314-E, Malcolm Circle, Cockeysville, MD, 21030.*

---

**ABSTRACT:** The antihyperglycemic activity of a polysaccharide fraction from *Lycium barbarum* (LBP) was investigated. The various parameters studied included body weight, fasting blood glucose levels, total cholesterol (TC) and triglyceride (TG) in diabetic and normal mice. LBP treatment (20, 40 mg/ kg body weight) for 28 days resulted in a significant decrease in the concentration of fasting blood glucose (FBG), total cholesterol (TC) and triglyceride (TG) in diabetes mellitus mice. Furthermore, LBP significantly increased body weight (BW). The data demonstrated LBP at the dose of 40 mg/kg bw exhibited the optimal effect.

**Key words** *polysaccharide from Lycium barbarum; antihyperglycemic; activity*

---

### **INTRODUCTION**

Diabetes mellitus is a disease affecting approximately 3% of the population worldwide (Skyler, 2004). For a long time, diabetics have been treated with several medicinal plants or their extracts based on folklore medicine (Akhtar and Ali, 1984). Synthetic hypoglycemic agents can produce serious side effects and in addition, they are not suitable for use during pregnancy (Pari and Saravanan, 2002). Therefore, the search for more effective and safer hypoglycemic agents has continued to be an important area of active research.

*Lycium barbarum* species are deciduous woody perennial plants, produce a bright orange-red, ellipsoid berry 1-2 cm long. The berries have been used in traditional Chinese medicine for about 1,900 years (Jin et al., 2006). They have been used for the treatment of cerebral arteriosclerosis, liver or heart diseases, hypercholesterolemia and diabetes (Zhao and Liu, 2008). The active components of *Lycium barbarum* primarily contain water-soluble polysaccharides (Chen and Mu, 2007). They could be extracted with hot water followed by precipitation with ethanol to obtain high quantity of polysaccharides (Zhi et al., 2004; Gan et al., 2004; Zhang et al., 2004). LBP have been recently studied for their physiological and pharmaceutical activities. The purpose of this study was to investigate the hypoglycemic effect of LBP in alloxan-induced diabetic mice.

### **MATERIAL AND METHOD**

#### **Plant materials**

Dried *Lycium Barbarum* were purchased from a local drug market and the material was identified by

---

Manuscript received: September 2009; Accepted: December 2009

\*Address for correspondence: Tel.: +8613170662021

E-mail address: jinglongjun666@yahoo.cn

Mr. King Li, a botanist of Qujing Normal University. A voucher specimen has been deposited in herbarium of Qujing Normal University.

#### **Drugs and reagents**

Alloxan was purchased from Sigma Co. (USA). Glucose Analyzer and strips were purchased from Roche Diagnostic Co. (USA). Reagents for total cholesterol (TC) and triglyceride (TG) were obtained from Beijing Chengxinde Biochemistry Reagent Company (Beijing, China). Reagents for serum insulin was purchased from Adlitteram Diagnostic Laboratories Co. (USA).

#### **Extraction of LBP**

Dried *Lycium Barbarum* was crushed in an electrical grinder and then powdered, 1000 g of this powder was immersed in tenfold dH<sub>2</sub>O, boiled at 100°C for 12 h (Ayiguli et al., 2007; Luo et al., 2000) and then the water extract was collected. The process was repeated once, and the extracts were combined and concentrated with a vacuum rotary evaporator at 70 °C. The concentrated solution was precipitated with addition of 4 times volume 95% ethanol and the precipitation was washed in turn with 100 % ethanol, 100 % Ether and acetone, polysaccharide from *Lycium barbarum* was obtained by vacuum drying (Luo et al., 2000). The *Unico-7200* spectrophotometer (*Unico* Co., Shanghai, China) was used to determine the content of polysaccharides in the above extracted product at 490 nm (Ayiguli et al., 2007; Wang et al., 2007).

#### **Experimental animals**

Male mice of original Kun-ming strain (18-22 g each) were used for the study. The study was carried out according to the "Principles of Laboratory Animal Care" (World Health Organization (WHO) Chronicle, 1985). A standard pellet diet and water were given ad libitum. Animals were maintained under a constant 12-h light and dark cycle and an environmental temperature of 21–23 °C.

#### **Preparation of alloxan-induced diabetic mice**

Diabetes was induced in fasted mice (12 h) by intraperitoneal injection of 200 mg/kg bw of alloxan, freshly dissolved in sterile normal saline immediately before at a concentration of 40 g/L. Diabetes was confirmed by the determination of tail vein blood glucose levels on the third day after administration of alloxan. The mice with a blood glucose level above 11 mmol/L, as well as with polydipsia, polyuria, and polyphagia were selected for the experiment (Zhang et al., 2004; Yang et al., 2006).

#### **Experimental Design**

Forty Male mice were randomly divided into five equal groups as follows:

- i) Normal control group (NC): normal control mice administered water daily for 28 days;
- ii) Diabetic control group (DC): diabetic control mice administered water daily for 28 days;
- iii) Diabetic + LBP (20mg/kg) group (DLL): diabetic mice administered LBP (20mg/kg) daily for 28 days.;
- iv) Diabetic + LBP (40 mg/kg) group (DLH): diabetic mice administered LBP (40 mg/kg) daily for 28 days.;
- v) Diabetic + glibenclamide (4 mg/kg) group (DG): diabetic mice administered reference drug glibenclamide (4 mg/kg) daily for 28 days.

Animals in the control, NC and DC groups were subjected to forceful feeding of 0.5ml distilled water/100 g bw daily for 28 days to keep all the animals at same type of treatment condition with respect to BLP supplemented groups.

During LBP and Glib supplement for 28 days, fasting blood glucose level was measured for once every week. Blood was collected from tip of the tail vein and fasting blood glucose level was measured by using a glucose analyzer. At the same time, the body weight of each mouse was measured by balance. On 28th day of experiment, the mice were sacrificed by decapitation under light ether anesthesia and blood was collected from dorsal aorta and serum was separated by centrifugation for 5min and was kept at -20 °C for the biochemical assay of total cholesterol (TC), and triglyceride (TG). TC and TG were determined by enzyme methods.

#### **Acute toxicity studies**

LBP was tested for its acute toxicity in male mice. The test was carried out by single oral administration of LBP at doses of 80, 240, 400 mg/kg to different groups of mice (5 mice in each group). The mortality and general behavior was observed continuously for one hour, four hour, and intermittently for next six hour, and again at 24 h and 48 h. The parameters were observed are gross behavioral changes, grooming, alertness, sedation, loss of righting reflex, tremors convulsions (Mukund et al., 2008).

#### **Statistical analysis**

All results were expressed as means ± SEM for each group (N = 8). Data were analysed statistically by one-way analysis of variance (ANOVA). The significance of the difference between the means of test and control studies was established by student's t-test. P values of less than 0.05 were considered significant.

**RESULTS**

**Acute toxicity studies**

In the present study, toxicity test was carried up to high concentration of 400 mg/kg (10 times more than chosen dose). Even at this dose extract did not exhibit any sign of toxicity. Since the main purpose of this test is to get some idea on conspicuous behavioral changes and death, if any, and the LBP did not exhibit any toxic symptoms in the limited toxicity evaluation in male mice.


**Effect of LBP on body weight**

The alloxan-induced diabetic mice exhibited loss of body weight. Before embarking on the experiment, all the groups had no significant difference in body weight ( $P > 0.05$ ). A significant ( $P < 0.05$ ) decrease in body weight was detected in the DC, DLL and DLH groups

as compared to the normal control group from 7 days after alloxan injection. However, the body weights in the DLH groups were significantly ( $P < 0.05$ ) and dose-dependently increased as compared to those of the diabetic control from 14 days after administration, which is comparable to that of the DG group. The results were shown in Fig.1.

**Effect of LBP on fasting blood glucose levels**

The alloxan-induced diabetic mice exhibited hyperglycemia. At the beginning, a significant ( $P < 0.05$ ) increase in FBG was detected in the diabetic groups as compared to the normal control group. But these abnormal increases in blood glucose levels significantly ( $P < 0.05$ ) and dose-dependently decreased in the LBP-administered groups as compared to the diabetic control group from 7 days after administration.



**Fig.1:** Effect of *Lycium barbarum* on body weight (g) in mice

**Table 1**  
Effect of LBP on blood glucose Level (mmol/L) in mice

Groups	Days after dosing (day)				
	0	7	14	21	28
NC	5.03±0.14	5.02±0.16	4.98±0.17	4.92±0.08	5.04±0.11
DC	15.24±0.39 <sup>①</sup>	15.21±0.26 <sup>①</sup>	15.21±0.35 <sup>①</sup>	15.03±0.15 <sup>①</sup>	15.12±0.29 <sup>①</sup>
DLL	15.13±0.30 <sup>①</sup>	9.43±0.21 <sup>①②</sup>	8.41±0.21 <sup>①②</sup>	7.45±0.15 <sup>①②</sup>	6.79±0.16 <sup>①②</sup>
DLH	15.28±0.28 <sup>①</sup>	8.62±0.11 <sup>①②</sup>	7.61±0.28 <sup>①②</sup>	6.12±0.17 <sup>①②</sup>	5.23±0.16 <sup>①②</sup>
DG	15.17±0.29 <sup>①</sup>	8.96±0.12 <sup>①②</sup>	7.12±0.23 <sup>①②</sup>	6.35±0.24 <sup>①②</sup>	5.56±0.34 <sup>①②</sup>

n=8; (mean±S.D., g); <sup>①</sup>P < 0.05 as compared with normal control group.; <sup>②</sup>P < 0.05 as compared with diabetic control Group

In the DG group, decrease was also significant ( $P < 0.05$ ) from 7 days after administration. NC and DC groups did not show any significant variation on the blood glucose level throughout the experimental period ( $p > 0.05$ ). The results were shown in Table 1.

**Table 2**  
Effect of LBP on blood lipids (mmol/L) in mice

Groups	TG	TC
NC	1.58±0.03	2.64±0.04
DC	1.99±0.03 <sup>①</sup>	3.28±0.05 <sup>①</sup>
DLL	1.71±0.03 <sup>①②</sup>	3.03±0.08 <sup>①②</sup>
DLH	1.64±0.03 <sup>①②</sup>	2.75±0.11 <sup>①②</sup>
DG	1.66±0.0 <sup>①②</sup>	2.097±0.05 <sup>①②</sup>

$n=8$ ; (mean±S.D., g); □ $P < 0.05$  as compared with normal control group; □ $P < 0.05$  as compared with diabetic control Group

#### Effect of LBP on blood lipids levels

Diabetes mellitus is usually complicated with hyperlipoproteinemia. The present results showed that the TC and TG levels were significantly elevated in the diabetic control group as compared to the normal control group ( $P < 0.05$ ). After supplementation with LBP, the alteration in lipid metabolism was partially attenuated as evidenced by decreased serum TG and TC levels in diabetic mice. The response was better in DLH group compared to the others group. The results were shown in Table 2.

#### DISCUSSION

Diabetes mellitus is a serious chronic disease. Although oral anti-hyperglycemic agents and insulin are often successful in diabetes treatment, they have prominent side effects and fail to significantly alter the course of diabetic complications. Effective control of the blood glucose level is a key step in preventing or reversing diabetic complications and improving the quality of life in both type 1 and type 2 diabetic patients (Chen et al., 2008). The present study shows that alloxan-induced diabetic mice presented obvious hyperglycemic symptoms, but LBP produces a significant antihyperglycemic effect when oral administration to alloxan-diabetic mice. The dosage of 40 mg/kg is more effective than that of 20 mg/kg.

Diabetes is also associated with hyperlipidemia. The serum TC and TG have been decreased significantly in diabetic mice after LBP supplementation. These effects

may be due to low activity of cholesterol biosynthesis enzymes and or low level of lipolysis which are under the control of insulin (Sharma et al., 2003).

*Lycium barbarum* have been used to treat diabetes in folk tradition for a long time. From this study, we could conclude LBP possesses hypoglycemic effects and the dose of 40 mg/kg bw represents the optimal level for effecting a positive diabetic response in mice. Toxicity data have already proved that the LBP did not show any toxic reactions. So, it can be said that *Lycium barbarum* is a good natural material to develop new agent to treat diabetes, maybe the effective constituent is polysaccharide.

#### REFERENCES

- Akhtar FM, Ali MR. (1984). Study of antidiabetic effect of a compound medicinal plant prescription in normal and diabetic rabbits. Journal of the Pakistan Medical Association, 34: 239–244.
- Ayiguli A, Wang Y, Yang XJ, Han HX, Bao XW, Zhu JF, Yu NN (2007). Study on Extraction Method and Assay of Polysaccharides of *Lycium barbarum* L. from Two Different Source. Xinjiang Agricultural Sciences, 44: 724–728.
- Chen H, Mu M. (2007). Studies on extraction of polysaccharides from *Lycium Barbarum*. Journal of Anhui Agricultural Sciences, 35: 3736–3737.
- Chen J, Li WL, Wu JL, Ren BR, Zhang HQ (2008). Hypoglycemic effects of a sesquiterpene glycoside isolated from leaves of loquat (*Eriobotrya japonica* (Thunb.) Lindl.). Phytomedicine, 15: 98–102.
- Gan L, Hua ZS, Liang YX, Bi XH (2004). Immunomodulation and antitumor activity by a polysaccharide-protein complex from *Lycium barbarum*. International Immunopharmacology, 4: 563–569.
- Jin LE, Xie XM, Chang LP (2006). Studies on Extraction of Antioxidant from *Lycium* and Its Antioxidant Activities. Chemistry and Industry of Forest Products, 26: 55–58.
- Luo Q, Yan J, Zhang SH (2000). Isolation and purification of *lycium barbarum* polysaccharides and its antifatigue effect. Journal of Hygiene Research, 29: 115–116.
- Mukund H, Rao CM, Srinivasan KK, Mamathadevi DS, Satish H (2008). Comparison between the effects of alloxan and streptozotoci on inducing diabetes in mice. Pharmacognosy Magazine., 15: 819–824.
- Pari L, Saravanan G (2002). Antidiabetic effect of Cogent db, a herbal drug in alloxan-induced diabetes mellitus. Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology, 131: 19–25.

- Sharma SB, Nasir A, Prabhu KM, Murthy PS, Dev G (2003).** Hypoglycemic and hypolipidemic effects and antioxidant activity of fruit extracts from *Lycium barbarum*. *Journal of Ethnopharmacology*, 85: 201–206.
- Skyler JS. (2004).** Diabetes mellitus: pathogenesis and treatment strategies. *Chemistry*, 47: 4113–4117.
- Wang HB, Bai HJ, Wang JL (2007). Studying on Ultrasonic2Microw ave Synergistic Extraction Polysaccharides in *Lycium rehnicum* Murr. *Acta Agriculturae Boreali-Occidentalis Sinica*, 16: 157–158.
- World Health Organization (1985).** Principles of laboratory animal care. *World Health Organization Chronicle*, 39: 51–56.
- Yang RJ, Li Qw, Zhao R (2006).** Comparison between the effects of alloxan and streptozotoci on inducing diabetes in mice. *Journal of Northwest Sci-Tech University of Agriculture and Forestry*, 2: 17–20.
- Zhang R, Zhou J, Jia Z, Zhang Y, Gu G (2004).** Hypoglycemic effect of *Rehmannia glutinosa* oligosaccharide in hyperglycemic and alloxan-induced diabetic rats and its mechanism. *Journal of Ethnopharmacology*, 90: 39–43.
- Zhang RX, Zhou JH, Jia ZP (2004).** Hypoglycemic effect of *Rehmannia glutinosa* oligosaccharide in hyperglycemic and alloxan-induced diabetic rats and its mechanism. *Journal of Ethnopharmacology*, 90: 39–43.
- Zhao YH, Liu H(2008).** Studies on extraction of polysaccharides from *Lycium Barbarum*. *China Food Additives*, 4: 73–75.
- Zhi F, Zheng W, Chen P, He M (2004).** Study on the extraction process of polysaccharide from *Lycium barbarum*. *Journal of Chinese Medicinal Materials*, 27: 948–950.