

INVITRO ANTIMICROBIAL SUSCEPTIBILITY PATTERN OF BACTERIAL ISOLATES FROM WOUND INFECTIONS IN UNIVERSITY OF ILORIN TEACHING HOSPITAL

¹TAWO S.S., ²OKESINA A.B., ¹ONILE B.A.

¹Department of Microbiology and Parasitology, ²Department of Chemical Pathology and Immunology
University of Ilorin Teaching Hospital, P.M.B. 1459, Ilorin, Nigeria.

The outcome of 532 wound swabs received from patients with wound infections in different units of the University of Ilorin Teaching Hospital Ilorin, Nigeria, over a one year period (July 2000 – June 2001), and routinely processed by Gram staining and culture in the Microbiology Laboratory, is reported. 444(83.5) of all samples cultured positive for bacterial pathogens while 88 (16.5%) were bacteriologically sterile. 272 swabs yielded single isolate while 172 yielded a mixture of two or more organisms. *Staphylococcus aureus* predominates (35.8%), followed by *Pseudomonas* spp (21.8%), *Escherichia coli* (15.3%), *Klebsiella* spp (13.4%), *Proteus* spp (5.6%), *Coagulase Negative Staphylococci* (3.1%), *Streptococcus faecalis* (2.8%), *Streptococcus pyogenes* (0.9%), Group B β -haemolytic *Streptococci* (0.9%), and *Acinetobacter* spp (0.3%).

Both Gram-positive and Gram-negative organisms demonstrated moderate to high in vitro sensitivity to Ofloxacin and Ciprofloxacin (sensitivity rate 70-94%). In vitro sensitivity to Cloxacillin, Erythromycin, Azithromycin and Ceftazidime by Gram positive organisms ranged between 55 and 90% while Gentamicin, Ceftrazidime and Azithromycin equally demonstrated moderate to high inhibitory effect on Gram negative organisms including *Pseudomonas* spp. (sensitivity rate 55-90%).

The Fluoroquinolones are the favoured antimicrobial agents nowadays, as demonstrated in this study. In our environment however, a combination of Cloxacillin and Gentamicin is an effective empiric alternative when cost is considered and this combination can be used. The need for continuous antimicrobial monitoring of clinical isolates of wound infection for drug resistance, which is of paramount importance in the empiric selection of antibiotics, is emphasized.

INTRODUCTION

Every individual carries a large resident microbial population on the skin surfaces, and in the openings of the hair follicles, sweat glands and sebaceous glands. This population comprises mainly Gram positive cocci of the genera *Staphylococcus* and *Micrococcus*, and Gram positive rods of the genera *Propionibacterium* and *Corynebacterium* together with the yeast, *Pityrosporum* (1). The skin is also host to a variable number of transient or contaminating bacteria. Although the resident flora produce antibacterial substances that provide some protection against colonization by potential pathogens, any breach in the skin surface, whether accidental or surgical, provides an open door for bacterial infection.

Surgical wound infection rates have been found to vary between 3 and 11% and wound, skin and burns are areas after genito-urinary tract, where nosocomial infections tend to occur more commonly in surgical practice (2,3,4). The risk of infection increases with the degree of contamination and it has been estimated that about 50% of wound contaminated with bacteria become clinically infected. The prevalent organisms that have been associated with hospital – acquired wound infection include *Staphylococcus aureus* which from various studies have been found to account for 20-40% (3), and *Pseudomonas aeruginosa* 5-15% of the nosocomial infection, with infection mainly following surgery and burns. Other pathogens such as enterococci and members of the enterobacteriaceae have been implicated, especially in immuno-compromised patients and following abdominal surgery (3).

It is also known that aside surgical units, intensive care units, nurseries, operating room theatre, and recovery rooms are units where

nosocomial wound infection frequently occurs (1,3). In the Accident and Emergency unit, accidental wound, clean or dirty, is one of the most common reasons for attendance by patients. In all these units, wound infection which are mainly due to nosocomial pathogens, tends to be associated with bacteraemia, septicaemia, shock and death in some patients, and prolong hospital stay in many others. This situation may be a serious matter for the patient and his family, as his maintenance in the hospital and treatment are expensive and meanwhile a bed space is occupied which might otherwise be used for other patients.

In view of this, there is a need for continuous monitoring of the hospital by infect control team, which should particularly be aware of not only nosocomial wound infection but the local prevalence of antibiotic resistant bacteria strains, as this varies greatly from place to place. The pattern of the bacteria pathogens isolated from wound swabs in this hospital and their antibiotic sensitivity pattern is intended to provide Clinicians and Surgeons valuable information upon which empiric antimicrobial therapy of wound infection can be predicated.

MATERIALS AND METHOD

This study was carried out over a period of one year (July 2000 – June 2001) at the University of Ilorin Teaching Hospital, Ilorin, Nigeria. All wound swabs from different units of the hospital were received on the swab bench of the Microbiology laboratory and subjected to routine Gram staining and culture.

Gram staining was done according to the standard techniques (5). Swabs were inoculated onto Blood, Chocolate and MacConkey agar, and plates incubated aerobically at 37°C for 24 to 48 hours. Anaerobic culture was not done, as this is not a routine in our laboratory.

*Corresponding Author

Growth on culture plates were identified by colony morphology, and confirmed by Gram staining reaction, standard biochemical and serological tests(5).

Antibiotic susceptibility of pure culture of confirmed isolates were performed on Diagnostic Sensitivity Test Agar by the Kirby Bauer disc diffusion method (6) using the appropriate Gram positive and Gram negative discs, and Staphylococcus aureus, Escherichia coli and Pseudomonas aeruginosa as control strains. Isolate was considered sensitive or resistant by comparing zone diameter of inhibition to the zone diameter interpretive standard of the National Committee for Clinical Laboratory Standard (7).

Necessary Patient bio-data were obtained from the laboratory request forms and data were fed into EPI INFO version 6.0 computer with analysis done using the appropriate statistical methods where necessary.

RESULT

Of all the wound swabs received from 532 patients with clinical evidence of wound infection over the period of study, 346 (65%) were from in-patients, 132 (24.8%) from outpatients and 54 (10.2%) were from patients whose wards or clinics were not indicated on the request forms. The distribution of swabs and isolates by wards is shown in Table 1. Surgical wards accounted for the highest number of request and isolation rates, followed by outpatient units and lowest in Psychiatric and Obstetrics and Gynaecology units.

Of the 532 swabs, 444 (83.5%) cultured positive for bacterial pathogens while 88 (16.5%) were bacteriologically sterile. 272 (61.5%) of these yielded single, 152 (34.2%) yielded two while 20 (4.5%) yielded a mixture of three organisms (Table II).

The distribution of bacteria pathogens in pure and mixed cultures is as shown in Tables III and IV. A total of 642 bacterial isolates were obtained in all, 280 (43.6%) were Gram positive while 362 (56.4%) were Gram negative. Staphylococcus aureus was the predominant organism isolated accounting for 35.3%, followed by Pseudomonas spp (21.8%), Escherichia coli (15.3%), Klebsiella spp (13.4%), Proteus spp (5.6%), Coagulase Negative Staphylococci (3.1%), Streptococcus faecalis (2.8%), Streptococcus pyogenes (0.9%), Group B β -haemolytic Streptococci (0.9%), Acinetobacter spp (0.3%).

The antimicrobial profile of the pathogens is summarized in Table V and VI. The fluoroquinolones (Ofloxacin, Perfloxacin and Ciprofloxacin) showed increased activity against all the isolates. Cloxacillin, Erythromycin, Gentamicin and Azithromycin equally showed good activity against Staphylococcus aureus, the predominant Gram positive isolate, with 77.4%, 87.8%, 93.9% and 96.5% of isolates sensitive. Ceftazidime is the only Cephalosporin that

showed moderate activity against Staphylococcus aureus with 52.2% of the isolates sensitive. Ampicillin and Penicillin G were ineffective against Staphylococcus aureus with only 18.3% and 16.5% of the isolates sensitive, but Streptococcus pyogenes and Group B β -haemolytic Streptococci are highly sensitive to these agents.

WARDS	SWABS	ISOLATES
Surgical (W2, W5, W8)	168(31.6)	144(32.4)
Outpatient (SOP/MGP/GOP)	132(24.8)	124(27.9)
Medical (W1, W4, W6)	52(9.8)	44(9.9)
Paediatric (W3)	52(9.8)	32(7.2)
Emergency (A/E & EPU)	50(9.4)	36(8.1)
Obstetrics and Gynaecology	12(2.3)	10(2.3)
Intensive Care Unit	10(1.9)	8(1.8)
Psychiatric (W7)	2(0.4)	2(0.5)
Not Indicated	54(10.2)	48(10.8)
TOTAL	532(100)	444(100)

Key:

W = Ward

Number in parenthesis = Percentages

Table 1: Distribution of wound swabs and isolates by wards

Organism	No.	(%)
Staphylococcus aureus	230	35.8
Pseudomonas spp	140	21.8
Escherichia coli	98	15.3
Klebsiella spp	86	13.4
Proteus spp	36	5.6
CONS	20	3.1
Streptococcus faecalis	18	2.8
Group B β - haemolytic Strept	6	0.9
Streptococcus Pyogenes	6	0.9
Acinetobacter spp	2	0.3
Total	642	100%

Table 2: Distribution of Bacterial Pathogens Isolated from 44 wounds swabs

Azithromycin, Gentamicin and Ceftazidime respectively showed good activity against *Pseudomonas* spp, the most prevalent Gram negative pathogen, with 60%, 64.3% and 85.7% of isolates susceptible. Other Gram negative bacteria with the exception of *Acinetobacter* spp are equally susceptible to these antibiotics.

Organism	Number	(%)
<i>Staphylococcus aureus</i>	116	(42.6)
<i>Pseudomonas</i> spp	58	(21.3)
<i>Escherichia coli</i>	46	(20)
<i>Klebsella</i> spp	24	(8.8)
Coagulase Negative <i>Staphylococci</i>	12	(4.4)
<i>Proteus</i> spp	6	(2.2)
<i>Streptococcus pyogenes</i>	4	(1.5)
<i>Streptococcus faecalis</i>	4	(1.5)
Group B β -haemolytic <i>streptococci</i>	2	(0.7)
TOTAL	272	(100)

No in parenthesis = Percentages

Table 3: Distribution of bacteria pathogens from wound swabs in pure cultures

No in parenthesis = Percentages

Table 4: Mixed bacteria growth in wound swabs

ORGANISM	Number	(%)
<i>Staphylococcus aureus, Escherichia coli</i>	26	(15)
<i>Staphylococcus aureus, Klebsiella</i> spp	24	(14)
<i>Pseudomonas</i> spp, <i>Klebsiella</i> spp	24	(14)
<i>Staphylococcus aureus, Pseudomonas</i> spp	22	(13)
<i>Staphylococcus aureus, Proteus</i> spp	16	(9)
<i>Pseudomonas</i> spp, <i>Escherichia coli</i>	12	(7)
<i>Staphylococcus aureus, Pseudo. spp, Klebsiella</i> spp	8	(5)
<i>Staphylococcus aureus, Streptococcus faecalis</i>	4	(2)
<i>Escherichia coli, Coagulase Negative Staphylococci</i>	4	(2)
<i>Streptococcus faecalis, Proteus</i> spp,	4	(2)
<i>Staphylococcus aureus, Proteus</i> spp, <i>Pseudomonas</i> spp	4	(2)
<i>Staphylococcus aureus, Pseudomonas</i> spp, <i>E. coli</i>	4	(2)
<i>Pseudomonas</i> spp, <i>Proteus</i> spp	4	(2)
<i>Pseudomonas</i> spp, <i>Coagulase Negative Staphylococci</i>	2	(1)
<i>Klebsiella</i> spp, <i>Escherichia coli</i>	2	(1)
<i>Klebsiella</i> spp, <i>Proteus</i> spp	2	(1)
<i>Staphylococcus aureus, Streptococcus pyogenes</i>	2	(1)
<i>Escherichia coli, Streptococcus faecalis</i>	2	(1)
<i>Acinetobacter</i> spp, <i>Group B βhaemolytic Streptococci</i>	2	(1)
<i>Coagulase Negative Staphylococci, Strept. Faecalis</i>	2	(1)
<i>Pseudo</i> spp, <i>Group B βhaemolytic Strept / Klebs. Spp.</i>	2	(1)
<i>Streptococcus faecalis, Proteus</i> spp, <i>Pseudomonas</i> spp	2	(1)
TOTAL	172	(100)

Organism	Penicillin 1unit	Ampicillin 10µg	Streptomycin 10µg	Gentamicin 10µg	Cloxacillin 5µg	Erythromycin 10µg	Chlaramphenicol 10µg	Ceftazidime 30µg	Ceftaxime 30µg	Ceftazidime 30µg	O-floxacin 10µg	Ciprofloxacin 10µg	Azithromycin 10µg		
Staph aureus S	38(6.5%) 192(83.5%)	42(18.3%) 188(81.7%)	148(64.3%) 82(35.7%)	216(93.9%) 14(6.1%)	178(77.4%) 52(22.6%)	202(89.8%) 28(12.2%)	160(69.6%) 70(30.4%)	148(64.3%) 82(35.7%)	178(77.4%) 52(22.6%)	120(52.2%) 110(47.8%)	214(93.6%) 16(7%)	194(84.3%) 36(15.7%)	210(91.3%) 20(8.7%)	222(96.5%) 8(3.5%)	
N230 CONS S	0(0%) 20(100%)	0(0%) 20(100%)	12(60%) 8(40%)	10(50%) 10(50%)	6(30%) 14(70%)	14(70%) 6(30%)	16(80%) 4(20%)	20(100%) 0(0%)	8(40%) 14(60%)	20(100%) 0(0%)	14(70%) 6(30%)	14(70%) 6(30%)	20(100%) 0(0%)	14(70%) 6(30%)	
N20 Strept faecalis S	0(0%) 18(100%)	0(0%) 18(100%)	14(77.8%) 4(22.2%)	10(55.6%) 8(44.4%)	4(22.2%) 14(77.8%)	8(44.4%) 10(55.6%)	0(0%) 18(100%)	6(33.3%) 12(66.7%)	4(22.2%) 14(77.8%)	4(22.2%) 14(77.8%)	16(88.9%) 2(11.1%)	18(100%) 0(0%)	18(100%) 0(0%)	14(77%) 4(22.2%)	8(44.4%) 10(55.6%)
N18 Op Biflaemo. S	6(100%) 0(0%)	6(100%) 0(0%)	4(66.7%) 2(33.3%)	6(100%) 0(0%)	6(100%) 0(0%)	4(66.7%) 2(33.3%)	4(66.7%) 2(33.3%)	4(17%) 2(8.3%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	
N6 Streptococi	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	0(0%) 6(100%)	
N6 Strept pyogenes S	6(100%) 0(0%)	4(66.7%) 2(33.3%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	6(100%) 0(0%)	

Tables 5: Susceptibility pattern of Gram-Positive cocci isolated from wound swabs to antimicrobial agents

Organism	Ampicillin 10µg	Tetracycline 25µg	Streptomycin 10µg	Gentamicin 10µg	Cefuroxime 30µg	Ceftaxone 30µg	Ceftazidime 10µg	O-floxacin 10µg	Perfloxacin 10µg	Ciprofloxacin 10µg	Azithromycin 10µg
Pseudomonas spp S	56(40%) 84(60%)	28(20%) 112(80%)	86(61.4%) 54(38.6%)	90(64.3%) 50(35.8%)	20(14.3%) 120(85.7%)	58(41.4%) 82(58.6%)	120(85.7%) 20(14.3%)	110(78.6%) 30(21.4%)	104(74.3%) 36(25.7%)	120(85.7%) 20(14.4%)	84(60%) 56(40%)
N 140 Escherichia S	48(48.9%) 50(51.1%)	56(57.1%) 42(42.9%)	40(40.8%) 58(59.2%)	66(65.9%) 32(44.1%)	54(47.1%) 44(52.9%)	40(23.5%) 58(76.5%)	68(58.8%) 30(41.2%)	76(77.5%) 22(22.5%)	72(73.5%) 26(26.5%)	82(83.7%) 16(16.3%)	84(85.7%) 14(14.3%)
N98 Klebsiella spp S	52(60.5%) 34(39.5%)	36(41.9%) 50(58.1%)	70(81.4%) 16(18.9%)	56(65.1%) 30(34.9%)	42(48.8%) 44(51.2%)	40(46.5%) 46(53.5%)	62(72.1%) 24(37.9%)	78(90.7%) 8(9.3%)	68(79.1%) 18(20.9%)	74(85%) 12(14%)	74(85%) 12(14%)
N86 Proteus spp S	24(65.7%) 12(33.3%)	12(33.3%) 26(66.7%)	28(77.8%) 8(22.2%)	34(94.4%) 2(5.6%)	30(83.3%) 6(16.7%)	36(100%) 0(0%)	30(83.3%) 6(16.7%)	34(94.4%) 2(5.6%)	36(100%) 0(0%)	36(100%) 0(0%)	26(72.2%) 10(27.8%)
N36 Acinetobacter spp S	0(0%) 2(100%)	2(100%) 0(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)	0(0%) 2(100%)
N2											

Table 6: Susceptibility pattern of Gram-negative rods isolated from wound swabs to antimicrobial agents

DISCUSSION

Bacteria contamination of wound is a serious problem in the hospital especially in the surgical practice where clean operations can become contaminated and subsequently infected (2,8). Although it has been argued that wound swabs from surface of intact or ulcerated skin for culture, provides little or no clinically useful information, because of lack of correlation between surface colonization and below-the-surface infection (9), it is nonetheless known that the degree of wound contamination from surface wounds become clinically infected (2). 83.5% of wound swabs in this study cultured positive for bacteria pathogens. If 50% of these were indeed from infected wound, then the wound infection are will be 41.3%. This figure is slightly higher than the 39% recorded in Lagos (10).

Surgical wards posted the highest number of request and isolation rates of organism. This is in agreement with the trend world wide (3,8,10,11), which is attributable to the fact that patients here are likely to undergo surgical operation, and more likely to have breaks in their local defense systems. The low rate of request and isolation rate in intensive care unit as against the normal trend may be due to the fact that this unit is quite small and requests are therefore correspondingly small. It may also be a reflection of strict hygiene and good nursing practice in this unit.

The common pathogens isolated are *Staphylococcus aureus* (35.8%), *Pseudomonas* spp (21.8%), *Escherichia coli* (15.3%), *Klebsiella* spp (13.4%), *Proteus* spp (5.6%), and CONS (3.1%). The preponderance of *Staphylococcus aureus* is in keeping with other studies (3,9,11,12,13). The organism is a normal flora of the skin in most people and can easily contaminate wounds. 56.4% of all isolates are Gram negative organisms against 43.6% Gram positive bacteria. This is similar to the observation in some other centres (10) where *Pseudomonas* spp, *Klebsiella* spp, *Escherichia coli* and Coliforms are the predominant pathogens responsible for wound and other nosocomial infections. This pattern is best understood in terms of selective pressure exerted on the organism based on the current antibiotic use. In our environment, the third generation Cephalosporins are increasingly ssssss being used.

The susceptibility pattern of the organisms heavily favours the Quinolones, particularly Ciprofloxacin, and the new macrolide, Azithromycin, which are effective but expensive antibiotics in the treatment of wound infections in this environment. Ciprofloxacin has to be used with caution in the paediatric age group. In the light of 74.4% sensitivity of *Staphylococcus aureus* to Cloxacillin and 87.8% to Erythromycin, and greater than 60% sensitivity of the predominant Gram negative organisms to Gentamicin, a cost effective empiric combination of Cloxacillin and Gentamicin or Erythromycin and Gentamicin may be favourably considered for wound infection in this environment.

It is recommended that in addition to using the above antimicrobial therapy in the treatment of wound infection, adequate attention should be placed on preventive measures such as hand washing, disinfection, good nursing practice and good surgical techniques amongst others, to reduce bacterial contamination of wounds.

ACKNOWLEDGEMENT

We are grateful to Miss Fatimah Yusuf for typing the manuscript.

REFERENCES

1. Deurden BI, Reids TMS, Jewsbury JM, Turk DC (eds). A new short textbook of microbial and parasitic infection. Edward Arnold London. 1987.
2. Editorial Nosocomial infections Part II. The Ibadan Surgeons 1992; 19(4): 113-16.
3. Balows A, Hausler WJ jr., Hermann RL, et al. (eds). Manual of Clinical Microbiology. 5th edition American Society of Microbiology Press, Washington DC. 1991.
4. Hughes JM, Culver DH, White JW, et al. Nosocomial infections surveillances, 1980-82 morbid, Mortal, Weekly Rep. CDC Surveillance Summaries 1983; 32 (spec. supp 1.4) ISS-16SS.
5. Cheesborough M (ed) Medical Laboratory Manual for Tropical Countries. Vol. II Microbiology Cambridge University Press 1984.
6. Bauer AW, Kirby QMM, Sherris JC et al. Antibiotic susceptibility testing by a standardized single disk method. Am. J. Clin Path 1966; 45: 493-96.
7. National Committee for Clinical Laboratory Standards Performance standards for antimicrobial disk susceptibility tests. NCCLS documents M2-A6, Approved standard 6th edition; Wayne PA: NCCLS 1997.
8. Odelowo EOO, Onile BA Perioperative Infections in Nigerians: A seven year prospective study. E. Afr. Med. J. 1990; 67(3): 172-81.
9. Castiglia M, Smego RA jr. Skin and Soft Tissue infections in: Mahon CR, Manuselis G. (eds) Textbook of Diagnostic Microbiology W.B. Saunders Co. 1995; 970-91.
10. Ogunsola FT, Oduyebo O, Iregbu KC, et al. A review of Nosocomial infection at the Lagos University Teaching Hospital: Problems and strategies for improvement J. Nig. Infect. Contr. Ass. 1998; 1(1): 14-20.
11. Ayliffe GAJ, Lowsbury EJI, Geddes AM, Williams JD (eds) control of Hospital Infection (A practice Handbook) 2nd edition Chapman and Hall Medical, London 1992.
12. Dixon RE, effect of infections on Hospital care. Ann. Intern. Med. 1978; 89:749-53.
13. Couper RG, Summer C. Hospital Infection data from a children's hospital. Med. J. Aust. 1970; 2: 1110-1113.