

**APPLICATION OF FACTORIAL DESIGN FOR THE OPTIMIZED PRODUCTION OF ANTISTAPHYLOCOCCAL METABOLITE BY *AUREOBASIDIUM PULLULANS*****<sup>1</sup>Kalantar <sup>2</sup>Rajendra D. <sup>3</sup>Dr. Kalantar E, Sanandaj, K. I.****<sup>1</sup>Department of Medical Microbiology, School of Medicine, Kurdistan University of Medical Sciences, Sanandaj, Kurdistan- Iran <sup>2</sup>Department of Microbiology, University of Pune, Pune - India <sup>3</sup>Department of Medical Microbiology School of Medicine Kurdistan University of Medical Sciences****Phone : 0098 871 ..... Fax : 0098 871..... E-mail: [kalantar.cnavat@yahoo.com](mailto:kalantar.cnavat@yahoo.com) & [ekalantar@hotmail.com](mailto:ekalantar@hotmail.com)****Abstract****Background:** Antimicrobial substances are mainly produced by bacteria and lower fungi, and have great roles in the treatment of most infectious diseases.**Purpose:** Production of antistaphylococcal metabolite from *A. pullulans* by development of a cultural medium using response surface methodology.**Methods:** Production of antistaphylococcal metabolite from *Aureobasidium pullulans* was optimized in shake flasks using a statistical experimental design approach. Effect of various components in the basal medium, glucose, peptone,  $\text{KH}_2\text{PO}_4$ , as well as initial pH and temperature were statistically combined using an experimental  $2^4$  fractional factorial two-level design and tested for their influence on maximal antistaphylococcal metabolite production. results were analyzed using response surface methodology (RSM) software.**Results:** Optimum production of antistaphylococcal metabolite occurred at glucose 2.0 %, peptone 2.5%,  $\text{KH}_2\text{PO}_4$  0.15%, pH 4.0 and temperature 30°C. The maximal amount of antistaphylococcal metabolite 900 U/flask from about 0.85 g of dry weight biomass was extracted.**Conclusion:** The antistaphylococcal activity of *A. pullulans* seemed to be associated with primary metabolite rather than secondary metabolite. However, this conclusion should be taken with caution because both secondary metabolites as well as antibiotics are heterogeneous group and our knowledge regarding the exact definitions and of secondary metabolite / antibiotics are far from the perfection.**Key words:** *Aureobasidium*, antistaphylococcal activity, production, factorial design**INTRODUCTION**

*Aureobasidium pullulans* (de Bary) is cosmopolitan yeast like fungus that occurs in diverse habitats including the phyllosphere of many crop plants and due to production of melanin, it is popularly known as black yeast (1-3). Literature survey shows few reports on production of antimicrobial compounds from *Aureobasidium pullulans* (3-4).

Despite extensive use of antibiotics and vaccination programs, infectious diseases continue to be leading cause of morbidity and mortality worldwide. Widespread antibiotic resistance, the emergence of new pathogens in addition to the resurgence of old ones, and the lack of effective new therapeutics exacerbate the problem (5). The need for safe and

effective antimicrobial compounds increases in parallel with the expanding number of immunocompromised patients at risk for invasive fungal / bacterial infections.

One of the commonest operations in the study of production of antimicrobial agents by microorganisms is the development of a medium to obtain maximum cell and metabolic product yield (6-7). The selection of media for microorganism's growth and metabolic products is usually based on a combination of experimentation and logic (8). Often such medium screening strategies involve the "one factor at a time" technique. This approach is tedious and time consuming, especially for a large number of variables.

Moreover, it does not guarantee the determination of optimal conditions (9). The experimental design constitutes an efficient tool and is well adapted for treating problems with a large number of variables. In particular response surface methodology can be used when presence of complex interaction is suspected (9).

In our preliminary studies in the development of the production medium, various parameters were found to be important factors in enhancing the antistaphylococcal metabolite formation. However, no systematic study to achieve optimum medium composition and process conditions has been reported for the production of antistaphylococcal metabolite. This work reports production of antistaphylococcal metabolite from *A. pullulans* by development of a cultural medium using response surface methodology.

## MATERIALS AND METHODS

ANTISTAPHYLOCOCCAL METABOLITE FROM *A. PULLULANS* WAS CARRIED OUT AS DESCRIBED IN OUR PREVIOUS ARTICLE (3).

Optimization of media for production of antistaphylococcal metabolite from *A. pullulans* using factorial design

A TWO LEVEL FACTORIAL DESIGN-EXPERIMENT WAS CARRIED OUT FOR FIVE VARIABLES VIZ. GLUCOSE (0.4, 2.0 AND 4.0 G%), PEPTONE (0.5, 2.5 AND 5.0 G%),  $\text{KH}_2\text{PO}_4$  (0.015, 0.15 AND 0.3 G%), pH (3.0, 4.0 AND 5.0) AND TEMPERATURE (25, 30 AND 35°C) AFFECTING THE PRODUCTION OF ANTISTAPHYLOCOCCAL METABOLITE BY *A. PULLULANS*.

## RESULTS

Optimization of media for production of antistaphylococcal metabolite by *A. pullulans*

To observe the effect of five variables namely glucose, peptone,  $\text{KH}_2\text{PO}_4$ , pH and temperature on production of antistaphylococcal metabolite, statistically designed experiments were performed. The variables having significant effect on production were evaluated by conducting 40 experiments which included two replicates of a  $2^4$  factorial experiments with all the four factors and eight center points. Results were analyzed using response surface methodology (RSM) software. Effect of temperature could not be established because at 35°C there was no response in terms of production of antistaphylococcal metabolite. Therefore, we dropped it from the analysis. Table 1 gives the responses obtained in the form of production of antistaphylococcal metabolite (U/flask). The experimental results obtained showed that all the variables had significant effect on production of antistaphylococcal metabolite from *A. pullulans* strain. Based on the identification of variables by the 2-level factorial design, a central composite design was developed for variables significantly affecting production of antistaphylococcal metabolite.

These studies revealed that the optimum production of antistaphylococcal metabolite occurred at 2.0 % glucose, 2.5% peptone, and 0.15%  $\text{KH}_2\text{PO}_4$ , pH 4.0 and temperature 30°C.

The responses obtained were statistically evaluated and the model was built based on the variables with confidence levels more than 95% (Table 2). The model generated was of the quadratic type, the selected P-values of linear and interactive variables have been mentioned in Table 3. The standard error in production of antistaphylococcal metabolite was estimated to be 35.75.

Figs. 1 a, b, c indicate that glucose-peptone,  $\text{KH}_2\text{PO}_4$ - peptone, pH-peptone,  $\text{KH}_2\text{PO}_4$ - glucose, pH-glucose, pH-  $\text{KH}_2\text{PO}_4$  have a quadratic relationship. With increase in any of them the antistaphylococcal metabolite production increased.

The model has a high correlation coefficient ( $R^2=0.8719$ ), a significant F-value (17.3212), an insignificant lack of fit F-value (1.8966) and standard

error less than 10 in all the factors. Based on the model equation, three-dimensional surface plots were constructed, which gave the optimal level of the variables and their linear, interactive or quadratic responses. The plots represent interaction of two variables while keeping others constant.

**TABLE 1: 2-LEVEL FACTORIAL DESIGN FOR PRODUCTION OF ANTISTAPHYLOCOCCAL METABOLITES BY *A. PULLULANS***

RUN	A	B	C	D	RESPONSE* (UNITS / FLASK)
1	-1	-1	-1	-1	200
2	1	-1	-1	-1	170
3	-1	1	-1	-1	240
4	1	1	-1	-1	210
5	-1	-1	1	-1	190
6	1	-1	1	-1	170
7	-1	1	1	-1	230
8	1	1	1	-1	200
9	-1	-1	-1	1	190
10	1	-1	-1	1	150
11	-1	1	-1	1	190
12	1	1	-1	1	220
13	-1	-1	1	1	180
14	1	-1	1	1	0
15	-1	1	1	1	180
16	1	1	1	1	200
17	-1	-1	-1	-1	210
18	1	-1	-1	-1	180
19	-1	1	-1	-1	220
20	1	1	-1	-1	190
21	-1	-1	1	-1	200

22	-1	-1	1	-1	230
23	-1	1	1	-1	190
24	1	1	1	-1	240
25	-1	-1	-1	1	180
26	1	-1	-1	1	260
27	-1	1	-1	1	160
28	1	1	-1	-1	260
29	-1	-1	1	1	240
30	1	-1	1	1	0
31	-1	1	1	1	220
32	1	1	1	1	220
33	0	0	0	0	380
34	0	0	0	0	420
35	0	0	0	0	460
36	0	0	0	0	480
37	0	0	0	0	430
38	0	0	0	0	380
39	0	0	0	0	450
40	0	0	0	0	420

\*RESPONSE IS IN TERMS OF PRODUCTION OF ANTISTAPHYLOCOCCAL METABOLITE PER FLASK. THE DATA OF RUNS IS THE MEAN OF THREE INDEPENDENT EXPERIMENTS.

A= GLUCOSE

C=  $\text{KH}_2\text{PO}_4$

B= PEPTONE

D= pH

**TABLE 2: ESTIMATES OF THE MODEL COEFFICIENTS**

WITH THEIR P- VALUES:

VARIABLES		ESTIMATE OF COEFFICIENT	P- VALUES*
A	GLUCOSE	-10	0.4363
B	PEPTONE	19.375	0.2542
C	KH <sub>2</sub> PO <sub>4</sub>	-10.625	0.4175
D	pH	-13.125	0.3546
AB	GLUCOSE - PEPTONE	16.875	0.2872
AC	GLUCOSE - KH <sub>2</sub> PO <sub>4</sub>	-13.125	0.3546
AD	GLUCOSE - pH	-4.375	0.6872
BC	PEPTONE -	10	0.4363

	KH <sub>2</sub> PO <sub>4</sub>		
BD	PEPTONE - pH	8.75	0.4784
CD	KH <sub>2</sub> PO <sub>4</sub> - pH	-12.5	0.3687
AA	GLUCOSE - GLUCOSE	-59.0625	0.0492
BB	PEPTONE - PEPTONE	-59.0625	0.0492
CC	KH <sub>2</sub> PO <sub>4</sub> - KH <sub>2</sub> PO <sub>4</sub>	-59.0625	0.0492
DD	pH - pH	-59.0625	0.0492

A = GLUCOSE, B = PEPTONE, C = KH<sub>2</sub>PO<sub>4</sub>, D = pH

\* P- VALUES LESS THAN 0.05 INDICATE SIGNIFICANT VARIABLES.


**TABLE 3: ANALYSIS OF VARIANCES OF MODEL:**

SOURCE	D.F.	SUM OF SQUARES	MEAN SQUARE	F	P VALUE
MODEL	11	407435	37039.5455	17.3212	0
LINEAR	4	24337.5	6084.375	2.8453	0.0426
A	1	3200	3200	1.4965	0.2314
B	1	12012.5	12012.5	5.6175	0.2049
C	1	3612.5	3612.5	1.6894	0.2043
D	1	5512.5	5512.5	2.5779	0.1196
INTERACTIONS	6	25887.5	4314.5833	2.0177	0.0967
AB	1	5512.5	9112.5	4.2614	0.0484
AC	1	5512.5	5512.5	2.5779	0.1196
AD	1	612.5	612.5	0.2846	0.5967
BC	1	3200	3200	1.4965	0.2314
BD	1	2450	2450	1.1457	0.2936
CD	1	5000	5000	2.3382	0.1375
QUADRATIC	2	357210	178605	83.523	0.0003
AA	1	357210	357210	167.046	0
BB	1	0	0	0	1
CC	1	0	0	0	1
DD	1	0	0	0	1
PURE ERROR	7	8950	1278.5714	0	0
LACK OF FIT	21	50925	2425	1.8966	0.196
ERROR	28	59875	2138.3929	0	0
TOTAL	39	467310	0	0	0

\* P- VALUES LESS THAN 0.05 INDICATE SIGNIFICANT VARIABLES.

A =  
GLUCOSE  
B = PEPTONE

ACTIVITY (UNIT / ML)





ACTIVITY (UNIT / ML)

Fig. 1.a. Optimization of antistaphylococcal metabolite from *A. pullulans* using factorial design. A two level factorial design experiments was carried out for optimization of production of antistaphylococcal metabolite for four variables viz glucose (0.4, 2 and 4 %), peptone (0.5, 2.5 and 5%),  $\text{KH}_2\text{PO}_4$  (0.015, 0.15 and 0.3%) and pH (3, 4 and 5).

A = GLUCOSE  
B = PEPTONE  
C =  $\text{KH}_2\text{PO}_4$   
D = PH


ACTIVITY (UNIT / ML)



ACTIVITY (UNIT / ML)


FIG. 1.B. OPTIMIZATION OF ANTISTAPHYLOCOCCAL MET

UNDER FIG. 1.A.



B = PEPTONE  
C =  $\text{KH}_2\text{PO}_4$   
D = pH

ACTIVITY (UNIT / ML)



ACTIVITY (UNIT / ML)




FIG. 1.C. OPTIMIZATION OF ANTISTAPHYLOCOCCAL METABOLITE FROM *A. PULLULANS* USING FACTORIAL DESIGN. AS

UNDER FIG. 1.A.

## DISCUSSION

In order to optimize the parameters of production of antistaphylococcal metabolite by *A. pullulans* factorial design was used. In the traditional methods of optimization, since each parameter is independently investigated, the interaction effect is missed. Moreover, it is tedious and time consuming, especially for a large number of variables (10). According to Adinarayana and Ellaiah factorial experiments are a good way of judging the relative significance of the influencing factors and give a quantitative measure of the contribution of each factor to the overall response (11).

In the present investigation, a 2<sup>4</sup> factorial design was chosen to investigate the effect of parameters namely; glucose, peptone, KH<sub>2</sub>PO<sub>4</sub>, temperature and pH. Under optimized conditions *A. pullulans* produced 900 units of antistaphylococcal metabolite from the biomass grown in 100 ml medium under the optimum conditions.

The production of antistaphylococcal metabolite was parallel to log phase of growth, though, the initial lag of one day was observed before the beginning of accumulation of intracellular antistaphylococcal metabolite. The antistaphylococcal activity of *A. pullulans* seemed to be associated with primary metabolite rather than secondary metabolite. However, this conclusion should be taken with caution because both secondary metabolites as well as antibiotics are heterogeneous group and our knowledge regarding the exact definitions and of secondary metabolite / antibiotics are far from the perfection.

## ACKNOWLEDGEMENTS

The authors gratefully acknowledge the financial support of Jawaharlal Nehru Memorial Fund, New Delhi, India. The authors especially thank Dr. S.D. Gore, department of computer sciences, university of Pune for his assistance and advice in the development of this work.

## REFERENCES

1. Winokur, T.D. Color variants of *Aureobasidium pullulans* over produce xylanase with extremely high specific activity. *Appl. Environ. Microbiol.* 1986;52: 1026.
2. Takesako, K., K. Ikai., F. Haruna., M. Endo., K. Shimanaka., K. Sona., T. Nakamura and I. Kato. Aureobasidins, new antifungal antibiotics, taxonomy, fermentation, isolation and properties. *J. Antibiotics.* 1991;44:919-924.
3. Kalantar E, Deopurkar R, and Kapadnis B. Antistaphylococcal metabolite from *Aureobasidium pullulans*: production and characterization. *Afr. J. Clin. Expt. Microbiol.* 2005; 6(3), 177-187.
4. McCormack, P., G. Howard and P. Jefferies. Production of antistaphylococcal compounds by phyloplane inhabiting yeasts and yeast like fungi. *Appl. Environ. Microbiol.* 1994; 60: 927-931.
5. Ritu, Dhand. Microbial infection and immune defense. *Nature*, 2000; 106:7. 759.
6. Maddox IS, Richert SH. Use of response surface methodology for rapid optimization of microbiological media. *J. Appl. Bacteriol.* 1977;43: 197-204.
7. Marisca GA, Munguis AL. Production and characterization of dextranase from isolated *Paccilomyces lilacinus* strain. *Appl. Microbiol. Biotechnol.* 1991; 36: 327-331.
8. Mateles, R and E, Battat. Continuous culture used for media optimization. *Appl. Microbiol. Biotechnol.* 1974; 28: 901-905.
9. Hounsa, C., M. Aubry., H Dubourguier and P. Hornez. Application of factorial and Doehlert design for optimization of pectate lyase production by a recombinant *E. coli*. *Appl. Microbiol. Biotechnol.* 1996; 45: 764-770.
10. K. Adinarayana<sup>1</sup>, P. Ellaiah. Response surface optimization of the critical medium components for the production of alkaline

protease by a newly isolated *Bacillus* sp. J.  
Pharm Pharmaceut Sci. 2002; 5 (3):272-278.

11. Roberto, I.C., Sato, S., Mancilha, I.M. and Taqueda, M.E.S., Influence of Media

Composition on Xylitol Fermentation by  
*Candida guilliermondii* Using Response  
Surface Methodology, Biotechnology Letters.  
1995; 17: 1223-1228.

Visit our website <http://www.ajol.info/journals/ajem>