

Full Length Research Paper

# Effects of Soy-based allochthonous nutrient inputs on intensively cultured female *Clarias gariepinus* (Burchell) brooder's growth performance and on the in-dwelling net microplanktonic populations

Olaleye, V. F.\*, Adewumi, A. A., Adedeji A. A. and Ogbogu S. S.

Department of Zoology, Obafemi Awolowo University, Ile-Ife, Nigeria.

Accepted 21 January, 2009

Portions of soybean meal autoclaved at 116°C and 1.2 kg/cm<sup>2</sup> pressure for 10, 15, 20, 25 and 30 min were used to compound 4 experimental diets appropriately labeled SB<sub>10</sub>, SB<sub>15</sub>, SB<sub>20</sub>, SB<sub>25</sub> and SB<sub>30</sub> respectively. The experimental diets were fed to *Clarias gariepinus* female broodstocks intensively for 84 days. During the experimental period, the female fish broodstock growth performance as well as the water and the in-dwelling net microplanktonic quality were monitored. The feed inputs into the different culture media ranged between 13,759 g (SB<sub>15</sub>) and 15, 649 g (SB<sub>10</sub>). The highest weight gain was recorded in the female broodfish fed diet SB<sub>25</sub>. The assayed water quality parameters for the different treatment culture media were not statistically different from each other ( $P > 0.05$ ). Four (4) phytoplanktonic phyla containing 7 orders with 30 species were recorded in the various media receiving the different feed inputs. The richest floristic composition with 21 species and a Margalef's Richness Index ( $R^1$ ) of 1.66 was obtained in the culture medium that received SB<sub>25</sub> feed input. Twenty (20) micro-invertebrate species belonging to 2 phyla and 4 orders were recorded in the various media during the period of study. The rotifers (Order: Ploima) with 16 species dominated the microinvertebrate fauna. Results also showed that the culture medium which received SB<sub>25</sub> diet had the highest micro-invertebrate fauna with 19 species.

**Key words:** Soybean, feed input, *Clarias gariepinus*, water quality, microplankton.

## INTRODUCTION

*Clarias gariepinus* which is a choice fish of culture in Nigeria requires adequate nutrition for fast growth and reproductive success (Adewumi, 2005). Culturing *C. gariepinus* broodstock intensively require continuous introduction of highly nutritious artificial feed into the culture system. Systematic addition of artificial feed and subsequent nutrient leaching from it as well as microbial decomposition of uneaten feeds and fecal matters pose a potential risk to water quality, the established biota and biological productivity (Bergheim and Asgard, 1996).

The leached nutrients and organic decompositions rich

in nitrogen and phosphorus (Temporeti et al., 2000) subsequently play a fundamental role in the inherent biogeochemical dynamics of the aquatic ecosystems (Charrier et al., 2000). The ecological relevance of the released nutrients is therefore contingent on their recycling or transfer to other compartments of the aquatic ecosystem (Beyruth and Tanaka, 2000). Inadequate nutrient compartmentalization or recycling however negatively impacts the water quality. Low survivability, poor growth rate and low reproductive potential are some of the reported sequential physical damage and nutrient enrichment induced intoxication effects on fish and other aquatic organisms (Barreto and Uieda, 2000). A major constraint in fish culture improvement in Nigeria is the provision of good quality feeds (Ezenwa, 1994). Balanced nutrition for fish broodstock enhances egg production, milt quality

\*Corresponding author. E-mail: Olafoluwa\_59@yahoo.com, volaleye@oauife.edu.ng.

(FAO, 2000) and confer superior health and growth potential on the resultant progenies (Takeuchi et al., 1978). Fishmeal (imported wholly into Nigeria), is the choice protein source in compounded fish feed because it is rich in essential amino acids (Adewumi, 2005).

Most current fish nutrition studies are presently focused on the replacement and/or supplementation of scarce and expensive imported fishmeal component of the fish diet with alternative cheaper protein sources (Adewumi et al., 2005). Soybean meal is used as an economic substitute for fishmeal in animal diets worldwide because of the rich quality of its amino acid profile, high digestibility (Lim and Akiyama, 1992) and ready availability (Adewumi et al., 2006). The use of soybean meal in fish feed production however, is limited by its deficiency in lysine and the presence of several anti-metabolites such as phytase and trypsin inhibitors (Khalifa et al., 1992). Protease inhibitors in raw or inadequately heated soybean meals adversely affect growth and reproductive performances in fishes (Peres et al., 2003). Adequate heat treatment of soybean meal is known to reduce the inherent trypsin inhibitory activity to a level where it supports optimal growth and reproductive efficiency in *C. gariepinus* (Adewumi et al., 2005).

The present study reports the effects of continuous and intensive feeding of soy-based diets to the *C. gariepinus* brooders growth performance as well as effects on water quality enrichment and planktonic community.

## MATERIALS AND METHODS

### Culture facility and the fish

The study was conducted in outdoor concrete tanks (6 x 5 x 1.5 m) filled with water from Obafemi Awolowo University Reservoir between May and August, 2005. *C. gariepinus* yearlings for the experiment ( $182 \pm 10$  g) were obtained from a commercial fish farm. Three hundred (300) healthy fish specimens were selected from the original stock of laboratory acclimatized 480 fishes and were distributed in batches of 30 into 10 labeled fine-meshed (0.752 mm) nylon hapas designated for the feeding trials. Water in the culture tanks were replenished periodically as the need arose to compensate for evaporation.

### Experimental diet preparation

Portions of full-fatted, dried soybean seeds (SAMSOY 2 TGX 636-02D) winnowed, heat treated for different times (in an autoclave) and milled were used as the primary protein sources to compound experimental diets for *C. gariepinus* brooders (Adewumi, 2005). The soybean portions were autoclaved at 116°C and 1.2 kg/cm<sup>2</sup> pressure for 10, 15, 20, 25 and 30 min respectively to remove the inherent trypsin inhibitors in the soybean meal (Peres et al., 2003) and to determine the appropriateness and adequacy of the various heating times used to remove trypsin inhibition. The differently autoclaved soybean portions were separately used along with other ingredients like: yellow maize, Tilapia fish meal and brewery waste (Adewumi, 2005) to compound approximately iso-nitrogenous (31% protein) and iso-caloric (3.34 kcalg<sup>-1</sup>) experimental diets. The prepared diets were appropriately labeled SB<sub>10</sub>, SB<sub>15</sub>, SB<sub>20</sub>, SB<sub>25</sub> and SB<sub>30</sub> (Table 1) to correspond with the different soybean heat processing times.

Ingredients for each experimental diet were properly mixed and extruded through an 8 mm diet using a Hobart A-200 mixer. The proximate compositions of the extruded pelleted experimental diets were determined according to AOAC (1990) while the gross energy contents were determined using Ackermann et al. (1969) method. Each of the diet was also wet-digested using perchloric acid and nitric acid mixtures (Diks and Allen, 1983) and the resultant digests were diluted using distilled water. The concentrations of Ca<sup>2+</sup>, Mg<sup>2+</sup>, Zn<sup>2+</sup>, Cu<sup>2+</sup> and Fe<sup>2+</sup> in the diluted digest were then determined using atomic absorption spectroscopy (APHA et al., 1992). Phosphorus concentration in the digest was determined using molybdate-sulphuric acid reagent method (APHA et al., 1992).

### Experimental feeding and data collection

The experiment which was primarily designed to determine adequacy of soybean toasting time on some growth parameters in *C. gariepinus* female brooders lasted for 84 days. During the experiment, the five prepared experimental diets (calculated on a dry matter basis) were fed twice daily at the allotted rate of 5% of the fish body weight. Each feed treatment was replicated. Feeding was accomplished by offering the prepared diet to the fish at the same spot between 08.00 - 09.00 G.M.T. and 18.00 - 19.00 G.M.T. Body weight changes in the fish were compensated for through weekly re-calculations of the rations offered to the experimental fish. Fish mortalities in the experimental tanks were monitored daily.

Data collection started on the first day of the feeding trials. Length (in cm) and weights (in gm) of individual fish stocked for each of the five dietary treatments were taken using standard techniques. Length-weight data collection repeated subsequently, weekly, was used to calculate the amount of feed offered, body weight changes and the condition factor. The gross feed input was determined according to Page and Andrews (1973) while the mean weight gained and the percentage body weight gained was obtained using Pitcher and Hart (1982) method. Estimation of the specific growth rate (SGR) was done according to Brown (1957) while Fulton's condition factor ( $K_f$ ) indicating the state of fish well-being was calculated using the method of Le Cren (1951).

### Water quality studies

Temperature, pH and conductivity of the water samples were determined in situ using a multi-probe HACH portable water laboratory daily. Duplicate sub-surface water samples collected fortnightly from each of the culture tanks using a Friedinger water sampler brought into the laboratory were analysed for the phenolphthalein acidity, carbonate alkalinity, phosphorus, nitrate and dissolved oxygen content (APHA et al., 1992). Na<sup>+</sup> and K<sup>+</sup> levels were determined flame photometrically (Golterman et al., 1978) while the Ca<sup>2+</sup>, Mg<sup>2+</sup>, Fe<sup>2+</sup> concentrations in the water samples were read using Alpha 4, Model 4200, Chem Tech Analytical flameless atomic absorption spectrophotometer (AAS). The concentrations of PO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> were also determined according to Golterman et al. (1978).

### Phytoplankton and microinvertebrate studies

Thirty (30) liters of water samples collected from each treatment tank fortnightly were filtered and concentrated to 20 ml using a 45 µ plankton net. The plankton concentrate samples were preserved in 5% formalin and two drops of Lugol solution for quantitative and qualitative examination. Planktonic identification and enumeration were done by introducing 1 ml of the preserved concentrate plankton samples into a Sedwick-Rafter counting chamber for examination through an Olympus BH2 Microscope. Planktonic identification

**Table 1.** Experimental diet formulations, the dietary energy content and the proximate composition of the experimental diets.

	Diets				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
<b>Formulation (g/100 g dry diet)</b>					
Soybean	34.43	34.43	34.43	34.2	34.58
Fishmeal	3.34	3.34	3.34	3.57	3.19
Brewery wastes	37.78	37.78	37.78	37.78	37.78
Yellow maize	22.45	22.45	22.45	22.45	22.45
Mineral/Vitamins <sup>1</sup>	1.00	1.00	1.00	1.00	1.00
Vegetable oil	1.00	1.00	1.00	1.00	1.00
Total	100.00	100.00	100.00	100.00	100.00
<b>Computations</b>					
Crude Protein (%)	30.96	31.25	30.89	30.83	31.01
Dietary Energy (Kcal g <sup>-1</sup> )	3.46	3.37	3.86	3.93	3.06
<b>Proximate composition (%) (<math>\bar{X}</math> (SD))</b>					
Crude protein	30.45 (3.12)	29.51 (0.88)	30.55 (1.42)	31.20 (2.18)	30.99 (2.51)
Moisture content	6.59 (0.08)	5.49 (0.06)	5.06 (0.96)	4.94 (1.88)	4.46 (0.59)
Lipid Content	4.02 (1.05)	4.65 (0.03)	4.68 (1.21)	4.81 (0.88)	3.82 (0.08)
Ash	6.81 (2.03)	7.22 (1.41)	7.41 (0.80)	7.85 (0.28)	9.33 (0.44)
Crude fibre	7.91 (0.09)	7.81 (1.02)	5.73 (0.09)	5.86 (0.28)	6.22 (0.55)
NFE <sup>2</sup>	44.22 (1.44)	45.32 (1.51)	45.90 (0.11)	46.34 (1.22)	45.18 (1.42)
<b>Mineral Composition (mg/g) (<math>\bar{X}</math> (SD))</b>					
Iron	10.00 (2.31)	16.02 (1.19)	11.19 (2.11)	11.30 (2.19)	14.64 (1.19)
Copper	0.07 (0.01)	0.06 (0.01)	0.19 (0.01)	0.20 (0.10)	0.13 (0.01)
Magnesium	3.39 (1.08)	3.36 (1.02)	3.17 (1.08)	3.60 (1.21)	3.36 (1.08)
Calcium	62.98 (10.11)	72.96 (9.02)	66.36 (6.18)	76.36 (12.01)	73.02 (11.09)
Phosphorus	5.90 (1.25)	6.28 (1.15)	6.86 (1.20)	7.40 (2.01)	5.38 (1.91)

<sup>1</sup>Mineral inclusion (g Kg<sup>-1</sup> dry diet): Manganese sulphate (MnSO<sub>4</sub>.4H<sub>2</sub>O) 0.0660, Iron sulphate (Fe SO<sub>4</sub>.7H<sub>2</sub>O) 0.3500, Copper sulphate (CuSO<sub>4</sub>. 5H<sub>2</sub>O) 0.0290, Calcium iodate (Ca IO<sub>3</sub>.6H<sub>2</sub>O) 0.0112, Sodium chloride (NaCl) 1.1800, Potassium chloride (KCl) 0.0800, Calcium orthophosphate (CaHPO<sub>4</sub>)2.550, Sodium selenite (Na<sub>2</sub> SeO<sub>3</sub>) 0.005, Vitamin inclusion (g Kg<sup>-1</sup> dry diet): Vitamin A 9\*10<sup>6</sup>, Vitamin O<sub>3</sub>1.25\*10<sup>6</sup>, Vitamin E 7\*10<sup>3</sup>, Vitamin B<sub>3</sub> 22, Vitamin B<sub>5</sub> 14, Riboflavin 6, Choline chloride 240, Amino Acid inclusion (g Kg diet): Lysine 120, Methionine 65.

<sup>2</sup>NFE = Nitrogen-free Extract = 100 – (Crude protein + Crude fibre + Lipid content + Moisture content + Ash).

was done to specific levels according to Edmondson (1959), Adeniyi (1978) and Fernando (2002). Planktonic abundance, species richness, diversity and evenness were calculated according to Ludwig and Reynolds (1988).

### Statistical analysis

Data collected were subjected to a non-parametric t-test (Mann-Whitney U-Noncoxon Rank Sum W-test) for the comparison of means within treatments while the Kruskal-wallis one way ANOVA was used to compare means at 5% level of significance (SPSS statistical package programme, version 12.0)

## RESULTS

### Feed input and fish growth performance

The soybean-based feed inputs into the culture system

ranged between 13,759 g (SB<sub>15</sub>) and 15,649 g (SB<sub>10</sub>) (Table 2) during the period of culture. The highest feed input into the culture medium per gram of fish flesh produced (1.88) was obtained in the culture medium that received SB<sub>10</sub> feed, followed by the media receiving SB<sub>30</sub> (1.70) and SB<sub>15</sub> (1.69) feeds respectively. The lowest value (1.31) was obtained in the culture medium that received SB<sub>25</sub> feed.

The growth performance of *C. gariepinus* female broodstock fed the soybean-based diets showed that the highest growth rate was obtained in the broodfish fed SB<sub>25</sub> diet closely followed by the fish fed SB<sub>20</sub> diet. Also, the percentage weight gained by the female broodfish fed SB<sub>20</sub> and SB<sub>25</sub> diets were significantly higher ( $P < 0.5$ ) than those of the fish fed other soybean-based diets. The specific growth rate (SGR) of the female broodstocks in the different culture media varied between  $0.22 \pm 0.05$

**Table 2.** Growth performance<sup>1</sup> of *C. gariepinus* female broodstocks during the period of study.

	Dietary Treatments				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
Initial Weight (g)	207±10 <sup>ab</sup>	182 ± 18 <sup>b</sup>	192 ± 28 <sup>ab</sup>	198 ± 9 <sup>a</sup>	206 ± 23 <sup>ab</sup>
Final Weight (g)	277±54 <sup>b</sup>	272 ± 65 <sup>ab</sup>	359 ± 73 <sup>a</sup>	379 ± 63 <sup>a</sup>	304 ± 82 <sup>b</sup>
Percentage body weight gained (%)	34.05 <sup>a</sup>	49.56 <sup>b</sup>	86.90 <sup>c</sup>	91.38 <sup>c</sup>	47.20 <sup>b</sup>
Specific Growth Rate	0.22±0.05 <sup>a</sup>	0.26 ± 0.05 <sup>ab</sup>	0.32 ± 0.16 <sup>b</sup>	0.34 ± 0.03 <sup>b</sup>	0.25 ± 0.05 <sup>a</sup>
Fulton's Condition Factor (kf)	0.82±0.25 <sup>a</sup>	1.02 ± 0.32 <sup>a</sup>	1.02 ± 0.26 <sup>a</sup>	1.11 ± 0.34 <sup>a</sup>	0.89 ± 0.22 <sup>a</sup>
Survival Rate (%)	100.0	98.0	100.0	100.0	100.0
Gross Feed Input (g)	15,649 <sup>a</sup>	13,759 <sup>b</sup>	14,515 <sup>c</sup>	14,968 <sup>c</sup>	15,573 <sup>a</sup>
Feed Input/g body weight gained	1.88 <sup>a</sup>	1.69 <sup>b</sup>	1.35 <sup>c</sup>	1.31 <sup>c</sup>	1.70 <sup>b</sup>

<sup>1</sup>Values with the same superscript in each row are not significantly different ( $P > 0.05$ ) from each other.

**Table 3.** Water quality parameters<sup>1</sup> of the culture media receiving different soybean-based feed input

Parameters	Dietary Treatments				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
<b>Physico chemistry</b>					
Temperature (°C)	28.75±0.97	28.88±0.83	28.50±0.87	28.38±0.92	28.88±0.85
pH	7.96±0.27	8.20±0.33	8.10±0.18	8.10±0.17	8.11±0.22
Total Alkalinity (mg/l)	141.89±15.45	179.25±1.30	172.52±15.27	173.63±16.08	174.75±14.13
Phenophatein Acidity (mg/l)	21.67±4.67	28.33±7.26	28.23±6.01	25.33±9.84	19.17±3.00
Conductivity (µs/cm)	32.30±1.02	36.43±1.00	33.63±0.56	34.88±2.14	34.25±0.33
Dissolved oxygen (mg/l)	3.76±1.02	5.20±2.31	3.93±1.65	2.70±1.82	4.43±1.65
<b>Ions (mg/l)</b>					
Na <sup>+</sup>	7.98±1.82	8.42±2.33	8.24±2.04	8.95±2.26	8.28±2.47
K <sup>+</sup>	48.56±18.11	50.60±12.96	47.40±13.92	51.10±19.39	49.27±18.05
Ca <sup>2+</sup>	29.05±6.37	31.35±4.44	31.03±7.43	43.05±17.53	42.97±20.27
Mg <sup>2+</sup>	0.67±0.12	0.70±0.21	0.68±0.14	0.66±0.10	0.59±0.13
Fe <sup>2+</sup>	3.45±1.81	5.91±1.09	2.25±0.85	2.91±1.09	1.60±0.65
PO <sub>4</sub> <sup>2-</sup>	13.06 ±1.00	18.36 ±0.78	12.79±3.28	18.37±3.44	18.65±1.43
NO <sub>3</sub> <sup>-</sup>	21.85 ±0.93	20.17±4.55	20.53±9.08	17.53±1.76	24.84±5.18

<sup>1</sup>Parameters not significantly different from each other ( $P > 0.05$ ).

(SB<sub>10</sub>) and  $0.34 \pm 0.03$  (SB<sub>25</sub>). Although variations occurred in the condition factor of the broodfish in the various culture media, the recorded differences were not significantly different ( $P > 0.05$ ) from each other (Table 2).

### Water quality

Analyses of the quality of water receiving the different dietary feed inputs shown in Table 3 revealed that the levels of the assayed parameters varied within a narrow amplitude of concentrations which were not statistically different from each other ( $P > 0.05$ ). Irrespective of the quality of the dietary nutrient input, the culture water was alkaline with a mean pH value ranging between 7.96 (SB<sub>10</sub>) and 8.20 (SB<sub>15</sub>) with a mean total alkalinity value of between 141.89 mg/l (SB<sub>10</sub>) and 179.25 mg/l (SB<sub>15</sub>).

Potassium (K<sup>+</sup>) was the dominant ion in the culture me-

dia, with concentration ranging between 47.40 mg/l (SB<sub>20</sub>) and 50.60 mg/l (SB<sub>15</sub>). Ca<sup>2+</sup> with concentrations varying between 29.05 mg/l (SB<sub>10</sub>) and 43.05 mg/l (SB<sub>25</sub>) also had relatively high concentration in the water of the various culture media. The recorded levels of NO<sub>3</sub><sup>-</sup> in the culture media ranged between 17.53 mg/l (SB<sub>25</sub>) and 24.84 mg/l (SB<sub>30</sub>) while the concentrations of PO<sub>4</sub><sup>2-</sup> ranged between 12.79 mg/l (SB<sub>20</sub>) and 18.65 mg/l (SB<sub>30</sub>).

### Phytoplanktonic composition

Four (4) phytoplanktonic phyla containing 7 orders with 30 species were recorded in the culture media receiving the differently toasted soybean-based feed inputs (Table 4). Irrespective of the type of feed input, the dominant cyanophyte in the culture media was *Anacystis incerta* (Order: Chroococcales). *Gomphosphaeria wichurae*

**Table 4.** Phytoplanktonic and abundance composition (individuals /m<sup>3</sup>) of the culture media receiving different soybean-based feed inputs.

	Dietary Treatments				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
<b>CYANOPHYTA</b>					
Order: Chroococcales					
<i>Anabacena variabilis</i> Kütz	-	-	-	17	-
<i>Anacystis incerta</i> Lemm.z	17275	78999	31133	44583	24216
<i>Gomphosphaeria wichurae</i> (Hilse)	6990	31125	66	8	3400
<i>Merismopedia</i> sp.	6900	8	10375	27500	-
<b>CHLOROPHYTA</b>					
Order: Volvocales					
<i>Eudorina elegans</i> (Ehren)	-	-	-	66	-
<i>Palmella miniata</i> (Leibl)	83	91	182	-	91
<i>Sphaerocystis schroeteri</i> (Chodat.)	51550	33	44724	27616	55,000
Order: Chlorococcales					
<i>Chlosteriopsis longissima</i> (Lemm.)	58	6974	157	6957	83
<i>Pediastrum biradiatum</i> (Meyen)	281	338	124	17300	248
<i>Scenedesmus denticulatus</i> (Lager.)	437	10466	198	361	17721
<i>Schroederia setigera</i> (Lemm.)	34508	-	-	17	-
<i>Tetraedron regulare</i> (Kütz)	8	-	33	66	-
<i>Treubaria crassispina</i> Smith	-	-	33	-	-
Order: Ulotrichales					
<i>Draparnaldia pulmosa</i> vaucher	8	-	-	-	-
<i>Stigeoclonium lubricum</i> (Dillw.)	8	-	-	-	-
<i>Ulothrix zonata</i> (Web et Mohr)	-	17	-	42	-
<i>Closterium acerolum</i> (Shrank)	-	-	-	8	-
<i>Closterium leibleini</i> (Kütz)	-	-	-	8	-
<i>Micrasterias radiata</i> (Hass)	17	25	17	50	8
<i>Mougeotia transedui</i> (Collias)	-	-	33	-	-
<i>Staurastrum arbutum</i> (Ralfs)	10449	8	8	6950	83
<b>EUGLENOPHYTA</b>					
Order: Euglenales					
<i>Euglena acus</i> (Ehren).	-	-	-	-	8
<i>Trachelomonas volvocina</i> Ehren.	107	99	99	8	33
<b>CHRYSOPHYTA</b>					
Order: Peridinales					
<i>Ceratium hirundinella</i> Schrank	-	10	-	-	6
Order: Bacillariales					
<i>Asterionella Formosa</i> Hass.	-	17	-	-	-
<i>Biddulphia laevis</i> Ehren.	-	8	-	-	-
<i>Navicula radiosa</i> Kütz	41	83	190	74	13,800
<i>Nitzschia sigmoidea</i> (Nitz.)	8	25	16	17	215
<i>Synedra ulna</i> (Nitz.) Ehren.	31125	34550	34548	17416	20816
<i>Species Richness</i> (R)	19	18	19	21	15

(Order: Chroococcales) was abundant in the culture media that received SB<sub>10</sub>, SB<sub>15</sub> and SB<sub>30</sub> experimental feeds while *Merismopedia* sp. was an important blue green alga in the culture media receiving SB<sub>10</sub>, SB<sub>20</sub> and SB<sub>25</sub> feed inputs.

*Sphaerocystis schroeteri* (Order: Volvocales) dominated the chlorophyte in most culture media. Other important green algae recorded in the different culture media belonging to the Order Chlorococcales include: *Pediastrum biradiatum* (SB<sub>25</sub>); *Scenedesmus denticulatus* (SB<sub>15</sub> and

**Table 5.** Microinvertebrate composition and abundance (individuals /m<sup>3</sup>) of the culture media receiving different soybean-based feed inputs.

	Dietary Treatments				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
<b>ROTIFERA</b>					
Order: Ploima					
<i>Albertia typhalina</i> Harr. and Myers	25	83	-	74	25
<i>Anuraeopsis racenesis</i> Lauter.	272	91	91	41	25
<i>Argonotholca foliacea</i> Ehren.	-	-	-	8	-
<i>Asplanchna priodonta</i> Goose	511	41	74	108	41
<i>Brachionus angularis</i> Goose	58	25	10565	7271	7372
<i>Brachionus calyciflorus</i> Pallas	25	223	231	314	33
<i>Brachionus falcatus</i> Zach.	10375	107	-	8	8
<i>Brachionus rubens</i> Hud. and Goose	305	1746	173	132	10573
<i>Euchlanis dilatata</i> Ehren.	-	-	-	-	-
<i>Keratella cochlearis</i> Goose	8	-	58	75	33
<i>Lecane (Lecane) luna</i> Muller	272	303	8311	7065	800
<i>Lecane (Monostyla) bulla</i> Goose	-	124	116	99	124
<i>Lepadella patella</i> Muller	8	-	17	-	17
<i>Polyarthra vulgaris</i> Ehren.	388	17	363	140	-
<i>Trichocerca bicristata</i> Goose	173	50	3945	248	545
<i>Trichocerca rutterni</i> Donner	17	8	1020	107	8
<b>ARTHROPODA</b>					
Order: Copepoda					
<i>Cyclops scutifer</i> Sars	421	17	157	215	124
Order: Cladocera					
<i>Daphnia magna</i> Straus	10466	116	25	17	421
<i>Nauplius</i> larvae	4011	91	710	436	347
Order: Diptera					
Chironomid larvae	148	24	35	49	60
Species Richness (R)	17	16	16	19	17

SB<sub>30</sub>); *Schroederia setigera* (SB<sub>10</sub>) and *Staurastrum arbuticulare* (Order: Ulotrichales) (SB<sub>10</sub> and SB<sub>25</sub>). The dominant diatom species in the various culture media was *Synedra ulna* (Order: Bacillariales), while *Navicula radio-sa* was abundant in the culture medium receiving the SB<sub>30</sub> feed input.

The culture medium receiving SB<sub>25</sub> feed input had the richest floristic composition with 21 species and a Margalef's Richness index (R<sup>1</sup>) of 1.66. The poorest phytoplanktonic assemblages (15 species) occurred in the culture medium receiving SB<sub>30</sub> feed input. The poor species richness recorded also reflected in the calculated Margalef's Index (1.17) obtained (Table 5). The Simpson's Index of diversity ( $\lambda$ ) for the phytoplanktonic assemblages in the culture media ranged between 0.16 (SB<sub>25</sub>) and 0.20 (SB<sub>15</sub>). The values obtained in the calculation of the Simpson's Index (known to be inversely proportional to diversity) revealed the dominance of a few species in the culture media receiving SB<sub>15</sub> and SB<sub>30</sub> feed inputs respectively.

The highest number of phytoplanktonic individuals per

cubic meter of water (180,628 ind/m<sup>3</sup>) was recorded in the experimental culture medium receiving SB<sub>10</sub> feed input closely followed by the culture medium that received SB<sub>15</sub> feed inputs (173,365 ind/m<sup>3</sup>). The least number of phytoplanktonic individuals (129,084 ind/m<sup>3</sup>) was recorded in the culture medium receiving SB<sub>20</sub> feed inputs (Table 4). Deductions from Hill's second diversity number (N<sub>2</sub>) showed that the number of phytoplanktonic species which accounted for the abundance in the various experimental culture media vary (Table 6). In the culture media receiving SB<sub>10</sub> and SB<sub>15</sub> feed inputs, 6 and 4 species accounted for 92 and 90% of the recorded abundance respectively. For the culture medium receiving SB<sub>25</sub> feed inputs, only 7 algal species accounted for 91% of the abundance compared to 5 species being responsible for 86% of the abundance in the culture medium receiving SB<sub>30</sub> feed inputs. The results also showed that 6 phytoplanktonic species accounted for 94% of the abundance in the culture medium that received SB<sub>20</sub> feed input. The Hill's Evenness indices (E4 and E5) (Table 6) confirmed the dominance of few abundant phytoplanktonic species

**Table 6.** The planktonic richness, diversity and evenness of the culture media receiving different soybased nutrient inputs.

Indices	Dietary Treatments				
	SB <sub>10</sub>	SB <sub>15</sub>	SB <sub>20</sub>	SB <sub>25</sub>	SB <sub>30</sub>
<b>Phytoplankton richness</b>					
Margalef ( $R^1$ )	1.49	1.41	1.53	1.66	1.17
<b>Diversity Indices</b>					
Simpson ( $\lambda$ )	0.18	0.29	0.26	0.16	0.20
Hill's Second Diversity Number ( $N_2$ )	6.67	3.47	3.86	6.22	4.91
Abundance (%)	92.0	90.0	94.0	91.0	86.0
<b>Evenness Indices</b>					
Hill ( $E_4$ )	0.83	0.78	0.88	0.88	0.84
Modified Hill ( $E_5$ )	0.79	0.72	0.84	0.86	0.80
<b>Zoo invertebrates richness</b>					
Margalef ( $R^1$ )	1.57	1.47	1.48	1.86	1.61
<b>Diversity Indices</b>					
Simpson ( $\lambda$ )	0.31	0.35	0.30	0.38	0.40
Hill's Second Diversity Number ( $N_2$ )	3.23	2.88	3.39	2.61	2.52
Abundance (%)	90.4	74.1	92.1	89.9	91.2
<b>Evenness Indices</b>					
Hill ( $E_4$ )	0.76	0.53	0.75	0.72	0.74
Modified Hill ( $E_5$ )	0.68	0.42	0.68	0.61	0.64

in the different culture media.

### Microinvertebrate composition

Twenty (20) microinvertebrate species belonging to 4 orders and 2 phyla were recorded in the various media receiving soybean-based nutrient inputs (Table 5). Analyses showed that the lowest number of species (16 species) was recorded in the culture media receiving SB<sub>15</sub> and SB<sub>20</sub> feeds while the culture medium receiving SB<sub>25</sub> feed had the highest microinvertebrate faunae of the culture media were dominated by the rotifers (Order: Ploima). In all, 16 rotifer species were recorded compared to 4 species of arthropods. The microinvertebrate population of the culture medium receiving SB<sub>10</sub> feed input was dominated by *Brachionus falcatus* (Rotifera) and *Daphnia magna* (Arthropoda). *B. rubens* (Rotifera) dominated the culture medium receiving SB<sub>15</sub> feed input while *B. angularis* (Rotifera) dominated the culture media receiving SB<sub>20</sub> and SB<sub>25</sub> feed inputs respectively. In the culture medium receiving SB<sub>30</sub> feed input, *B. angularis* and *B. rubens* were the dominant rotifers (Table 5).

Margalef index of diversity ( $R^1$ ) (Table 6) which ranged between 1.47 (SB<sub>15</sub>) and 1.86 (SB<sub>25</sub>) also reflected the micro-invertebrates faunistic richness in the media receiv-

ing the different soybean-based feeds. The Simpson's diversity index ( $\lambda$ ) for the microinvertebrates in the different culture media which ranged between 0.30 (SB<sub>20</sub>) and 0.40 (SB<sub>30</sub>) probably indicated the dominance of a few species. The culture medium receiving SB<sub>10</sub> feed had the highest number of microinvertebrate individuals (27, 483 ind/m<sup>3</sup>) closely followed by the culture media receiving SB<sub>20</sub> feed (25,891 ind/m<sup>3</sup>) and SB<sub>30</sub> feed (20,556 ind/m<sup>3</sup>) (Table 5). The calculated Hill's second diversity number ( $N_2$ ) showed that 3 species accounted for 90.4% of the microinvertebrate faunal abundance in the culture medium receiving SB<sub>10</sub> feed input compared to 4 species which accounted for 91.2% of the faunistic abundance in the culture medium receiving SB<sub>30</sub> feed input. Analyses further showed that 3 species accounted for 89.9% and 74.1% of the microinvertebrates in the culture media receiving SB<sub>25</sub> and SB<sub>15</sub> soybean-based feed inputs respectively (Table 6). The analyses also showed that the abundance of the species in the various culture media receiving feed inputs relatively diverge away from evenness.

### DISCUSSION

Significantly higher growth performances exhibited by female *C. gariepinus* broodfish fed SB<sub>25</sub> and SB<sub>20</sub> diets (compared to other experimental diets) indicated obvious differences in the nutrient composition of the experimental diet (Keembiyehetty and Gatlin, 1997; Adewumi, 2005). Heat treatment is known to deactivate trypsin inhibitory activity to varying degrees in soybean meal (Fournier et al., 2004). Anti-nutritional factors which were not deactivated completely in diets compounded from low-heated soybean-meals probably impaired the absorption of the needed essential amino acids needed for new protein formation in the fish (De Francesco et al., 2004). Conversely, adequate toasting of soybean meal component of the diets probably lowered the trypsin inhibitory activities (Savage, 1989) resulting in higher protein utilization in the fish (Peres et al. 2003) which subsequently translated significantly to the weight gained recorded (Adewumi, 2005). Fish fed the long-heated soybean-based diet (SB<sub>30</sub>) showed relatively low growth, as indicated by low weights of individuals fed the diet. Longer heating of soybean meal probably resulted in denaturation of heat labile essential amino acids and formation of indigestible protein-carbohydrate complexes with consequential reduction in the availability of the essential lysine and arginine (Renitz, 1984). The high condition factor recorded in the *C. gariepinus* broodfish culture in all the treatments might have resulted from the weight of maturing ovary (De Silva and Anderson, 1998) which was high during the period of study.

The characteristics of water in the different culture media showed positive chemical and biological sensitivities to impactment from the different soybean-based diets (Elliot et al., 2006). The soybean toasting time which

influenced the nutrient composition of the various diets probably directly dictate the type and concentration of nutrients leached from uneaten feeds and faecal matter into the culture medium. Secondly, the bacteria mineralization in the uneaten food and faecal matters is expected to add to the nutrient load of the various culture media (Van De Bund et al., 2004). Relatively high amount of  $PO_4$  and  $NO_3$  recorded in the culture media during the period of study probably disproportionately affected phytoplankton composition and productivity (Steiner et al., 2005; Van Ruijven and Berendse, 2005). Such increment in algal biomass and productivity with nutrient enrichment has been reported by Smith (2003). Variability of the nutrient types in the different culture media also affected the phytoplankton biodiversity (Dodson et al., 2000) and the phenology of the dominant taxa (Elliot et al., 2006). The phytoplankton biomass was largely dominated by many blooming species which include: *A. incerta*, *S. schroeteri*, *Spirogyra* varians, *S. ulna*, *P. biradiatum* and *S. setigera* (Loreau, 2000).

Variations which occur in the phytoplankton community structure of the media receiving the different diets was probably be due to species specific differences in nutrient use (Loreau, 2000; Cardinale et al., 2002). Complementary nutrient use which occurs when species do not exhibit complete overlap in nutrient use (Cardinal et al., 2004) probably accounted for the dominance of different phytoplankton species in the various media receiving the differently treated soybean-based diets. The zooplankton community structure was probably regulated by the phytoplankton quantity and quality (Qin and Culver, 1996). Zooplankton species composition and diversity is known to be dependent on the efficiency of zooplanktonic herbivory on the phytoplanktons (Sommer et al., 2001; Van Ruijven and Berendse, 2005; Romanuk et al., 2006). The study showed that compounding fish diets with adequately heat-treated soybean meal (20 – 25 min) produced female broodfish with higher mean weight gained. Conversely, low heat treatment of the soybean component of the fish diet produced fish with relatively low weights. Higher levels of trypsin inhibitory activities in the low-heated soybean-based diets caused growth depression in the fish due to reduced nutrient availability (Peres et al., 2003). Overheating (30 min) of the soybean component of the female fish brooder's diet which resulted in denaturation of some essential amino acids as well as complexing of others with the carbohydrate components (Renitz, 1984) also led to low weight gained in the fish. Desirable water quality parameters during the period of study were probably responsible for the low fish mortality recorded. The planktonic quantity and quality were dependent on the nutrient availability in the treatment tanks because other environmental parameters were homogenous during the period of study.

## REFERENCES

Ackermann T, Robert L, Bouch L (1969). Biochemical Microcalorime-

- try (Ed. Brown HD), Academic Press, New York. pp. 23-80.
- Adeniyi IF (1978). Studies on the physico-chemical factors and the planktonic algae of Lake Kainji, Nigeria. Ph.D. Thesis, University of Ife, Ile-Ife, Nigeria p. 597.
- Adewumi AA (2005). The effects of the heating time of soybean for broodstock nutrition on the reproductive performance of *Clarias gariepinus* (Burchell, 1822). Ph.D. Thesis, Obafemi Awolowo University, Ile-Ife. p. 162.
- Adewumi AA, Olaleye VF, Adesulu EA (2005). Egg and Sperm quality of the African catfish, *Clarias gariepinus* (Burchell) broodstock fed differently heated soybean-based diets. Res. J. Agric Biol. Sci 1 (1) : 17-22.
- Adewumi AA, Olaleye VF, Adesulu EA (2006). Swim-up fry production in the African catfish *Clarias gariepinus* (Burchell) broodstocks fed with differently heated soybean-based diets. Aquac. Nutr. 37 : 543 – 549.
- AOAC (Association of Official Analytical Chemists) (1990). Official Methods of Analysis (14<sup>th</sup> Edition) (Ed. Williams, S.), Arlington, V.A. p. 1102.
- APHA, AWWA, WEF (American Public Health Association, American Water Works Association and Water Environment Federation) (1992). Standard Methods for Examination of Water and Waster Waters. 18<sup>th</sup> Edition. Prepared and published by American Public Health Association, American Water Works Association and Water Environment Federation, New York, pp.39-134.
- Barreto MG, Uieda VS (2000). Influence of abiotic factors on the ichthyofauna composition in different orders stretches of Copivata River, Sao Paulo State, Brazil. Verh. Int. Verein. Limnol. 26(5): 2180-2183.
- Bergheim A, Asgard T (1996). Waste production from Aquaculture. In: Baird DJ, Beveridge MCM, Kolly LA, Muir JF (Eds) Aquaculture and Water Resource Management, Blackwell, New York, pp.50-76.
- Beyruth Z, Tanaka FM (2000). Biovolume of phytoplankton in aquaculture tropical ponds. Verh. Internat. Verein. Limnol. 27(2): 689-695.
- Brown ME (1957). Experimental studies on growth. In: Physiology of Fishes, Vol. 1, Academic Press Inc., New York, U.S.A. Chapter IX, pp.361 - 400
- Cardinale BJ, Palmer MA, Collins SL (2002). Species diversity enhances ecosystem function through interspecific facilitation. Nature, 415:426-429.
- Cardinale BJ, Ives AR, Inchausti P (2004). Effects of species diversity on the primary productivity of ecosystems: extending our spatial and temporal scales of inference. Oikos, 104: 437-450.
- Charrier C, Brunet C, Delcambre H, Touzeau C (2000). Effects of rapidly varying nutritional conditions on the dynamics of an algal population. Verh. Internat. Verein. Limnol. 27(5): 3086-3091.
- De Francesco MF, Medale F, Lupi P, Kaushik J, Poli BM (2004). Effect of long-term feeding with a plant protein mixture based diet on growth and body-fillet quality traits of large rainbow trout (*Oncorhynchus mykiss*). Aquaculture 235 (1-4): 413-429.
- De Silva SS, Anderson TA (1998). Fish Nutrition in Aquaculture. 308pp.
- Diks DM, Allen HE (1983). Correlation of copper distribution in a freshwater sediment system to bioavailability. Bull. Environ. Conta. Toxicol. 30: 37-43.
- Dodson SI, Arnott SE, Cottingham KL (2000). The relationship in lake communities between primary productivity and species richness. Ecology 81: 2662-2679.
- Edmondson WT (1959). Freshwater Biology. 2<sup>nd</sup> Edition, John Wiley and Sons, New York, p. 1248.
- Elliot JA, Jones ID, Thackeray SJ (2006). Testing the sensitivity of phytoplankton communities to changes in water temperature and nutrient load, in a temperate lake. Hydrobiology. 559: 401-411.
- Ezenwa BIO (1994). Aquaculture Development and Research in Nigeria. In: Aquaculture Development and Research in Sub-Saharan Africa (ed. AGcoche). CIFA Tech paper 23 suppl. FAO, Rome. pp.41 – 80.
- FAO (2000). Aquaculture Development Beyond 2000: The Bangkok Declaration and Strategy. Conference on Aquaculture Development in the Third Millenium. 20 -25<sup>th</sup> Feb., Bangkok, Thailand. FAO Aquaculture Newsletter. pp.11 – 18.
- Fernando CH (2002). A Guide to Tropical Freshwater Zooplankton



- Identification, Ecology and Impact on Fisheries. Backhuys Publishers Leiden, The Netherlands.
- Fournier V, Huelvan C, Desbruyeres E (2004). Incorporation of a mixture of plant feedstuffs as a substitute for fish meal in diets of juvenile turbot (*Psetta maxima*). *Aquac.* 236(1-4): 451-465.
- Golterman HL, Oymo RS, Onhastad MAM (1978). Method for Physical and Chemical Analysis of Freshwater. IBP Handbook No. 8, Oxford: Blackwell Scientific Publications pp.38-56.
- Keembiyehetty CN, Gatlin II DM (1997). Performance of sunshine bass fed soybean-meal based diets supplemented with different methionine compounds. *Prog. Fish. Cult.* 59(1): 25-30.
- Khalifa F, Belleville V, Sarda J, Prost J (1992). Short term effects of feeding raw or heated soya flour and casein meals on lipid intestinal digestion and absorption in rat. *J. Nut. Biochem.* 3: 224 – 231.
- Le Cren ED (1951). The length-weight relationship and seasonal cycle in gonad Weight condition in the perch (*Perca fluviatilis*). *J. Anim. Ecol.* 20: 201 – 219.
- Lim C, Akiyama DM (1992). Full-fat soyabean meal utilization by fish. *Asian Fisheries Sci.* 5: 181 – 197.
- Loreau M (2000). Biodiversity and ecosystem functioning: recent theoretical advances. *Oikos* 91: 3-17.
- Ludwig JA, Reynolds JF (1988). *Statistical Ecology: A Primer on Methods and Computing*. John Wiley and Sons, New York, pp.85-109.
- Page JW, Andrews JW (1973). Interactions of dietary levels of protein and energy on channel catfish (*I. punctatus*) *J. Nutr.* 103:1339 – 1346.
- Pitcher TJ, PJB Hart (1982). *Fisheries Ecology*. The AVI publication company Inc. West Port, Connecticut. pp.107 – 147.
- Peres H, Lim C, Klesus PH (2003). Nutritional value of heat-treated soybean meal for channel catfish (*Ictalurus punctatus*). *Aquaculture* 225(1-4): 67-82.
- \*Qin J, Culver DA (1996). Effect of larval fish and nutrient enrichment on plankton dynamics in experimental ponds. *Hydrobiol.* 321(2): pp??
- Reinitz I (1984). Protein dispersibility index (P.D.I.) as a quality control measure for soyflour used in brown trout starter feed. *Prog. Fish Cult.* 46(3): 161-164.
- Romanuk TN, Vogt RJ, Kolasa J (2006). Nutrient enrichment weakens the stabilizing effect of species richness. *Oikos*, 114: 291-302.
- Savage GP (1989). Antinutritional factors in peas. In: *Recent Advances in Research on Antinutritional Factors in Legume Seeds*. Proceedings of the First International Workshop on ANF in Legumes. Wageningen, Netherlands.
- Smith VH (2003). Eutrophication of freshwater and coastal marine ecosystems: a global problem. *Environ. Sci. Pollu. Res.* 10: 126-139.
- Sommer U, Sommer F, Santer B, Jamieson C, M, Becker C, Hansen T (2001). Complementary impact of copepods and cladocerans on phytoplankton. *Ecol. Lett.* 4: 545-550.
- Steiner CF, Darcy-Hall TL, Dorn NJ, Garcia EA, Mittelbach GG, Wojdak JM (2005). The influence of consumer diversity and indirect facilitation on trophic level biomass and stability. *Oikos* 110: 556-566.
- Takeuchi TW, Tanabe T, Ogino C (1978). Studies on the nutritive value of dietary Lipids in fish. XI. Supplementary effects of lipid in a high protein diet of Rainbow trout. *Bull. Jap. Soc. Sc. Fish.* 44: 677 – 686.
- Temporetti P, Lopez W, Gonzalez LP, Pedrozo F (2000). Limnological effects of intensive aquaculture on lakes and reservoirs and reduction of nutrient inputs. *Verh. Int. Verein. Limnol.* 27(4): 1812-1815.
- Van De Bund WJ, Romo S, Villena MJ, Valentin M, Van Donk E, Vicente E, Vakkilainen K, Svensson M, Stephen D, Stahl-Delbanco A, Rueda J, Moss B, Miracle MR, Kairesalo T, Hansson LA, Heitala J, Gyllstrom M, Goma J, Garcia P, Fernandez-Alaez M, Fernandez-Alaez C, Ferrinol C, Collings SE, B ecares E, Balayla DM, Alfouso T (2004). Responses of phytoplankton to fish predation and nutrient loading in shallow lakes: a pan-European meocosm experiment. *Freshwater Biol.* 49(12): 1608-1618.
- Van Ruijven J, Berendse F (2005). Diversity-productivity relationships: initial effects, long-term patterns, and underlying mechanisms. *Proc. Nat. Acad. Sci.* 102 .695-700.