

Review

Water hyacinth as a biosorbent: A review

Mahamadi C.

Department of Chemistry, Bindura University of Science Education, P. Bag 1020, Bindura, Zimbabwe. E-mail: cmahamadi@buse.ac.zw, courtiema@yahoo.com. Tel: +263 (0)777 350 801.

Accepted 8 December, 2011

Water hyacinth (*Eichhornia crassipes*), has attracted significant attention as the world's worst invasive aquatic plant due to its extremely rapid proliferation and congest growth, presenting serious challenges in navigation, irrigation, and power generation. Attempts to control the weed have proved to be costly with minimum results. However, the same plant has demonstrated an amazing ability to absorb and concentrate many toxic metals from aquatic environments. Consequently, research activity on utilization of the plant has been registered over the last few decades. This article reviews literature related to the utilization of *E. crassipes* in the biosorption and recovery of metals from aquatic environments. Availability in large quantities, high biosorption capacity, renewability and low cost determine the extent to which biosorbents can be adapted technologically. Sorption dynamics through classical and competitive models, effect of physical and chemical treatment, pH, temperature, initial metal concentration and biosorbent dose on metal removal by water hyacinth is discussed.

Key words: Biosorption, heavy metals, precious metals, recovery, water hyacinth.

INTRODUCTION

The toxicity and health hazards associated with heavy metals have been established beyond any doubts. Considerable attention has been paid to methods for metal removal and those commonly used include; electrochemical methods, reverse osmosis, chemical precipitation, ion exchange, biological processes, flotation and membrane processes (Park et al., 2010). Unfortunately, limitations synonymous with these methods have been understood to include incomplete metal removal, high energy or reagent requirements or the generation of toxic sludge and other waste products which require disposal. In addition, these methods are also limited when used for the decontamination of aquatic systems with adsorbates in a low concentration range (Vieira and Volesky, 2000; Volesky, 2007). As a result of

these challenges, there has been considerable interest in the use of other sorbent materials, particularly biosorbents (Volesky and Holan, 1995; Vasudevan et al., 2001; Vijayaraghavan and Yun, 2008).

'Biosorption' has been defined by Volesky (2007), a pioneer in the field, as the property of certain biomolecules (or types of biomass) to bind and concentrate selected ions or other molecules from aqueous solutions. As opposed to a much more complex phenomenon of bioaccumulation based on active metabolic transport, biosorption by dead biomass (or by some molecules and/or their active groups) is passive and based mainly on the "affinity" between the (bio) sorbent and sorbate (Volesky, 2007). The biosorption process involves a solid phase (sorbent or biosorbent)

and a liquid phase (solvent, normally water) containing a dissolved species to be sorbed (sorbate, metal ion). Due to higher affinity of the sorbent for the sorbate species, the latter is attracted and bound with different mechanisms. The process continues till equilibrium is established between the amount of solid-bound sorbate species and its portion remaining in the solution (Hussain et al., 2010). The biosorbent materials possess metal-sequestering property and can be used to decrease the concentration of heavy metal ions in solution from ppm to ppb level. Advantages of this technology over conventional ones include its low cost, high efficiency, minimisation of chemical and/or biological sludge, no additional nutrient requirement, regeneration of biosorbent and possibility of metal recovery following adsorption (Gadd, 1993).

In their recent review paper, Park et al. (2010), observed that biosorbents that have been sought for the removal of metal ions primarily fall into the following categories: bacteria, fungi, algae, industrial wastes, agricultural wastes, natural residues and other biomaterials, the former two perhaps giving broader choices. Some mycelia available in large quantities as waste materials from large scale fermentation processes have been used as a source of biosorbents conveniently (Kapoor and Virraghavan, 1995). Seaweeds, which can be obtained in large quantities from ocean and inexpensive, have also attracted attention as possible biosorbents (Suzuki et al., 2005). Other abundant natural materials which have been investigated for their heavy metal removal potential are the aquatic macrophytes such as *Pistia stratiotes* and *Lemna minor*, which are present at high growth rate (Wang, 1990; Klumpp et al., 2002). Recently, water hyacinth (*E. crassipes*) has attracted significant attention as a possible biosorbent for the removal of heavy metals (Mahamadi and Nharingo, 2007; 2010a; 2010b).

Water hyacinth is a wild fern belonging to the family pontederiaceae found abundantly in various tropical and sub-tropical countries of Latin America and the Caribbean, Africa, Southeast Asia and the Pacific (Mailu, 2001; Thomson et al., 2002; Mirona, 2004; Plummer, 2005). Generally, the level of reproduction of water hyacinth has been very high in countries where the plant has recently been introduced and this has been attributed, to a large extent, to the eutrophication in water bodies and the absence of natural enemies of the plant. The weed has attracted significant attention as the world's worst invasive aquatic plant due to its extremely

rapid proliferation and congests growth. In many places, the weed presents serious challenges in navigation, irrigation, and power generation (Malik, 2007). The explosive nature of water hyacinth growth and its related problems have called for serious efforts to control its spread. Management strategies have taken a physical, chemical or biological approach (Westerdahl and Getsinger, 1988; Ram and Moolani, 2000; Julien and Griffiths, 1998; Julien and Orapa, 1999; Mbatia and Neuenschwander, 2005; Kathiresan, 2000; Zhang et al., 2005; Shabana and Mohamed, 2005; Babu et al., 2004; Ogwang and Molo, 2004).

UNLOCKING VALUE OF WATER HYACINTH

As attempts to control the weed have caused high costs and labour requirements, leading to nothing but temporary removal of the water hyacinths, considerable research effort has been directed towards exploration of various ways of unlocking value in the otherwise troublesome weed. A number of possible uses of the plant, some of which have been developed and others are still in their infancy have been suggested. These include possible utilization of the weed as a phytoremediation agent (Sajn et al., 2005), as a biosorbent for toxic metals (Malik, 2007), for power alcohol and biogas production (Mshandete et al., 2004; Ali et al., 2004), as a compost (Szczeczek, 1999), and for animal fodder/ fish feed (Aboud et al., 2005). Other potential uses of the weed include its use as pulp material for producing greaseproof paper (Goswami and Saikia, 1994) and the production of fibreboards for a variety of end uses such as bituminized board for low-cost roofing material, indoor partitioning, etc. Formulations of water hyacinth-based medicines to treat certain diseases have been reported in India (Oudhia, 1999a, b).

Applications of water hyacinth in metal removal

Water hyacinth, among other aquatic macrophytes, has been shown to possess a great potential to remove pollutants when being used as a biological filtration system (Muramoto et al., 1991). It contains many polyfunctional metal-binding sites for both cationic and anionic metal complexes. Potential metal cation-binding sites of algal cell components include carboxyl, amine,

imidazole, phosphate, sulphate, sulfhydryl, hydroxyl and chemical functional groups contained in cell proteins and sugars (Mahmood et al., 2010c). The weed could remove several heavy metals and other pollutants (Pinto et al., 1987; Delgado et al., 1993; Zaranyika et al., 1994; Mahamadi and Nharingo, 2007, 2010a, 2010b). Recorded achievements triggered efforts directed towards the utilization of water hyacinth in phytoremediation.

Application of raw, dried water hyacinth roots in metal removal

Dried roots from water hyacinth have been used to study its removal behaviour from monocomponent aqueous metal solutions (Table 1). Schneider et al. (1995) studied the feasibility of dried water hyacinth roots for the removal of Pb^{2+} , Cu^{2+} , Cd^{2+} and Zn^{2+} ions from aqueous solution. The authors showed that in terms of lead and copper uptake per dried mass of biosorbent, dried roots and aerial parts of water hyacinth are better biosorbents than the biomass of the bacterium *Mycobacterium phlei*, the yeast *Candida parapsilosis*, fungal *Rhizopus oryzae* strains, and acacia bark. Sorption data for the metal ions fitted a Langmuir isotherm with the accumulation following the order $Pb^{2+} > Cd^{2+} \approx Zn^{2+}$ on a weight basis. It was deduced that the mechanism of adsorption involved the carboxylate group on the surface of the biomass. The authors further suggested that dried biomass of the plant might be placed in simple bags and used in a very low-cost metal ion removal system for decontamination of chemical and mining industrial wastewater. In an oncoming publication, Schneider et al. (2001), it was shown that the biomass of macrophytes *Potamogeton luscens*, *Salvinia herzogii* and *E. crassipes* can be subjected to a number of elution cycles without losing their adsorption capacity. Al Rmalli et al. (2005) showed that a biomaterial produced from dried water hyacinth roots, can provide a simple, effective, yet cheap method for removing arsenic from contaminated water. Elangovan et al., (2008) reported a study for the removal of Cr^{6+} and Cr^{3+} using dried aquatic weeds; reed mat (*Cannomois virgata*), water lettuce (*Pistia stratiotes*), arrow-leaved tear thumb (*Polygonum sagittatum*), lotus flower (*Nelumbo nucifera*), green taro (*Colocasia esculenta*), water lily flower (*Nymphaea* sp.) mangrove leaves (*Rhizophora mangle* L.) and water hyacinth. It was found that the removal of Cr^{3+} followed the order: reed

mat > water hyacinth > mangrove leaves > water lily > green taro, whereas for the removal of Cr^{6+} , mangrove leaves showed the greatest uptake.

On the removal of Cu^{2+} from aqueous solution by dried water roots, Zheng et al. (2010) showed that the biomass have a high affinity and large sorption capacity for the removal of the metal ion. Sorption equilibrium data obtained fitted the Langmuir isotherm and magnitude of the activation energy of the biosorption pointed towards a chemisorption process. The authors concluded that the water hyacinth roots are efficient and inexpensive biosorbent for Cu^{2+} removal. In a more recent publication, Hasan et al. (2010) utilized dried water hyacinth powder of the aerial part of the weed (stem and leaves) for the removal of Cr^{6+} and found that the sorption was negatively affected by pH whereas temperature had a positive effect.

Application of activated carbon and ash derived from water hyacinth in metal sorption

A growing number of researchers have explored the possibility of removing metal ions using activated carbon and ash derived from water hyacinth (Table 2). Shekinah et al. (2002) showed that removal of Pb^{2+} from aqueous solution using activated carbon prepared from water hyacinth increased with increase in pH. In a similar study utilizing activated carbon derived from water hyacinth, Kadirvelu et al. (2004) reported that the sorption of Hg^{2+} depended on contact time, pH, Hg^{2+} and carbon concentration. The sorption data obtained fitted the Langmuir and Freundlich models.

Mahmood et al. (2010c) reported that ash of water hyacinth satisfied the criteria for a hyperaccumulator of Pb^{2+} , Zn^{2+} , Cr^{6+} , Zn^{2+} and Ni^{2+} . Furthermore, it was possible to recover the metals from industrial waste water through desorption experiments. The authors concluded that water hyacinth-derived ash offers several advantages including cost effectiveness, high efficiency, minimization of chemical/biological sludge, and regeneration of biosorbent with possibility of metal recovery.

Application of acid/alkali treated water hyacinth in metal sorption

Biosorption usually contains organic functional groups

Table 1. Batch sorption of metal ions using raw dried roots derived from water hyacinth.

Experimental parameter	Removal/maximum sorption capacity	Reference	Comment
2 mg ml ⁻¹ biosorbent, initial pH 5.5, with agitation, initial metal concentration 6.3-20 mg l ⁻¹ .	Pb (47 mg g ⁻¹) Cu (23.0 mg g ⁻¹) Cd (27.0 mg g ⁻¹) Zn (20 mg g ⁻¹)	Schneider et al. (1995)	Biosorption capacity of water hyacinth found to be greater than that of yeast, bacterial and fungal strains.
30 mg l ⁻¹ biosorbent dose, 200 µg l ⁻¹ As(III) or As(v), pH 6, 120 min shaking time.	93% of As(III) 95% of As(v)	Al Rmalli et al. (2005)	As concentration in treated solution < 10 µg l ⁻¹ (WHO guideline limit value).
10 mg ml ⁻¹ absorbent, initial pH 5, 150 rpm agitation rate, 303 K temperature, initial concentration 10-150 mg l ⁻¹ , 6 hr sorption time.	Cr(III) (6.6 mg g ⁻¹), Cr(VI) (0.3 mg g ⁻¹).	Elangovan et al. (2008)	Biosorption capacity of water hyacinth found to be comparable to that of water lily, green taro, mangrove and reed mat.
0.4 mg ml ⁻¹ biosorbent, initial pH 5.5, 150 rpm agitation rate, 298 K temperature.	Cu (22.7 mg g ⁻¹)	Zheng et al. (2009)	Sorption of Cu appeared to involve ion exchange and complex formation.
2 mg ml ⁻¹ biosorbent, pH 2, 200 rpm agitation rate, 140 min sorption time 40°C.	Cr(VI) (91.6 mg g ⁻¹)	Hasan et al. (2010)	Chromium uptake decreased with an increase in pH, but increased with temperature and initial metal concentration.

such as alcohols, aldehydes, ketones, carboxylic, phenolic, and other groups on their surface. Ionization of these groups in aqueous solution enables them to participate in cation binding (Ofomaja and Ho, 2008). The ionization of functional groups on the biomass surface depends on pH of the solution: at pH > pK_a, most of these functional groups are mainly in ionized form and can exchange H⁺ with metal ions ion solution, at pH < pK_a, complexation phenomenon occurs (Fourest and Volesky, 1996). Several researchers have therefore studied biosorption of metal ions by acid/alkali-treated biomass (Table 3). Yao and Ramelow (1997) showed that up to 100% metal uptake was possible for Pb²⁺, Cu²⁺, Cd²⁺, Cr³⁺,

Mn²⁺, Ni²⁺, Co²⁺ and Ag¹⁺, with acidic and basic treatment enhancing the metal uptake. Acidic treatment was shown to increase the Cr⁶⁺ uptake capacity of water hyacinth roots whereas the removal of Cr³⁺ was lowered (Elangovan et al., 2008). Ibrahim et al. (2009) combined various spectroscopic techniques and quantum *ab initio* calculations to study the potential of water hyacinth dry matter for the removal of heavy metals from wastewater. It was shown that water hyacinth which is subjected to acetic acid is able to absorb acetate which finds its way to the cellulose of the plant. The dried and treated system acts like carboxylated cellulose and in particular, it can accelerate the removal of Cd

from the aquatic environment. It was further suggested that treated water hyacinth has the additional advantage that it reduces or even eliminates the diverse impact of the weed on the environment.

In a more recent publication, Mahamadi and Nharingo (2010b) carried out a competitive adsorption study using acid treated water hyacinth. The sorption dynamics the Pb²⁺, Cd²⁺, Zn²⁺ mixture were probed using q_e'/q_e ratios, where the prime denotes the presence of other metal ions. The equilibrium data showed that the uptake of ions from binary and ternary systems followed the order Pb >> Cd > Zn. The multielement effect was found to be antagonistic and the Pb-Zn,

Table 2. Batch sorption of metal ions using activated carbon and ash derived from water hyacinth.

Source/form of water hyacinth biosorbent	Experimental parameter	Removal/maximum sorption capacity	Reference	Comment
Water hyacinth-derived activated carbon	20 to 25 mg l ⁻¹ Pb(II), pH 3, 100-125 min sorption time.	16.6 mg g ⁻¹	Shekina et al. (2002)	Increasing pH from 2 to 4 resulted in increased Pb removal
Water hyacinth-derived ash	100 mg/ml adsorbent dose, pH 1.6-6.5, 5-30 mg l ⁻¹ Ni(II), Zn(II), Cd(II), Pb(II), Cr(VI) and Cu(II), 24 hr sorption time.	29.8 mg g ⁻¹ Pb(II), 24.0 mg g ⁻¹ Zn(II), 28.4 mg g ⁻¹ Cd(II), 29.8 mg g ⁻¹ Cu(II), 29.8 mg g ⁻¹ Ni (II).	Mahmood et al. (2010c)	Adsorption-desorption of the metals showed potential for their recovery.
Water hyacinth-driven activated carbon	10-40 mg l ⁻¹ Hg(II) concentration, pH 5, biosorbents particle size of 125-180 µm, 200 to 360 min sorption time.	28.4 mg g ⁻¹	Kadirvelu et al. (2004)	Sorption dependent on contact time, pH, Hg(II) and adsorbent concentration. Equilibrium data fitted both the Langmuir and Freundlich isotherm models.
Water hyacinth-derived ash	5-30 µg ml ⁻¹ Ni ²⁺ , pH 6.54, 25 °C temperature.	1.978 µg/g	Hussain et al. (2010)	Desorption capacity found to be 3.71 µg/g

Pb-Cd and Pb-Cd-Zn systems showed good fitting to the Langmuir competitive model. The authors make important conclusions that the removal of Pb²⁺ ions in the presence of comparable amounts of either Cd²⁺ or Zn²⁺ or both, showed greater potential than the removal of Cd²⁺ and Zn²⁺ in the presence of Pb²⁺.

METAL DESORPTION AND RECOVERY

An important consideration for the industrial application of water hyacinth is the question of metal recovery and biomass regeneration. Metal recovery is important especially when it involves valuable metals and desorption also becomes very much necessary when the biomass preparation/generation is costly. The economic application of a given biosorbent therefore

depends not only on its biosorptive capacity, but also on the ease of its generation and reuse. It has also been noted that a successful desorption process requires the proper selection of the eluants, which are a function of the biosorbent type and mechanism of biosorption (Das et al., 2010). Some workers have reported significant studies on the recovery of precious metals as well as those of industrial significance (Table 4).

Using *Sargassum* as the biosorbent, Sathishkumar et al. (2010), observed that the biosorbed ionic gold in metallic form through pyro-crystallization technique showed the possibility of 91.44% recovery of the metal in pure form. In conclusion, the authors noted that gold from dilute solutions could be recovered through green processes in ionic, nanocrystalline, or metallic form as described by the end-consumer using *Sargassum* as biomaterial. On the desorption of

gold from egg shell membrane, Ishikawa et al. (2002) showed that NaOH solution (0.1 mol/l) was capable of desorbing more than 95% of the gold (I) and NaCN (0.1 mol/l) was capable of desorbing more than 92% of gold(III). On application of an egg shell membrane column, 98% recovery of gold was reported. Kuyucak and Volesky (1989) observed that elution of gold adsorbed on brown marine alga, *Sargassum natans* could be achieved using a solution of 0.1 mol/l thiourea with 0.02 mol/l ferric ammonium sulphate. A limitation noted in the approach was the long desorption time (17 h). Recovery of ionic gold from exhausted decarboxylated biosorbent through elution (93%) was more efficient than incineration (61.89%) (Kwak and Yun, 2010). It was also reported that the combined method of biosorption and incineration could be effective for concentration and recovery of gold from the cyanide solutions.

Table 3. Batch sorption of metal ions using treated water hyacinth roots.

Experimental parameter	Removal (%)/maximum sorption capacity for single, binary and ternary component systems	Reference	Comment
5 mg l ⁻¹ initial metal concentration, initial pH 2-6. Acidic and basic treatment.	Up to 100% metal uptake was achieved for Pb, Cu, Cd, Cr, Mn, Ni, Co, Au, Ag, Cd –(single component)	Yao and Ramelow (1997)	Acidic and basic treatments increased the metal uptake for the metals.
2 mg ml ⁻¹ adsorbent, initial pH 4.8, 150 rpm agitation rate, 303 K temperature, initial concentration 10-60 mg l ⁻¹ , 3 h sorption time. Acidic treatment.	Pb (26.3 mg g ⁻¹)-single Zn (12.6 mg g ⁻¹)-single Cd (12.6 mg g ⁻¹)-single Pb (25.4 mg g ⁻¹)-binary (Pb-Cd) Pb (22.1 mg g ⁻¹)-binary (Pb-Zn) Cd (4.1 mg g ⁻¹)-binary (Cd-Zn) Zn (3.0 mg g ⁻¹)-binary (Zn-Pb) Pb (14.3 mg g ⁻¹)-ternary (Pb-Cd-Zn) Cd (3.0 mg g ⁻¹)-ternary (Cd-Zn-Pb) Zn (3.7 mg g ⁻¹)-ternary (Zn-Cd-Pb)	Mahamadi and Nharingo (2010a,b)	Pb ions were effectively removed from aqueous solution in the presence of both Cd and Zn, but removal of Cd ²⁺ and Zn ²⁺ ions was suppressed in the presence of Pb ²⁺ .
10 mg ml ⁻¹ adsorbent, initial pH 5, 150 rpm agitation rate, 303 K temperature, initial concentration 10-150 mg l ⁻¹ , 6 h sorption time. Acidic treatment	Cr(III) (6.2 mg g ⁻¹)-single Cr(VI) (6.4 mg g ⁻¹)-single	Elangovan et al. (2008)	Acid treatment significantly increased the Cr(VI) removal capacity whereas the removal of Cr(III) was lowered.
0.2; 1.2 mg ml ⁻¹ initial Cd concentration, 2 h sorption time.	55% uptake (low initial metal conc.) 80% uptake (high initial metal conc.)	Ibrahim et al. (2009)	Coordination of acetate to the plant found to occur via the cellulose of the plant where also Cd uptake takes place.

Silver recovery from AgNO₃ and Ag(S₂O₃)₂³⁻ showed that the second strain of microalgae *Chlorella vulgaris* gave a higher uptake percentage (Cordery et al., 1994). Ishikawa et al. (2002) reported no loss biosorbent capacity when egg shell membrane was taken through five sorption/desorption cycles during the sorption of Au (I) and Au (III). Ma et al. (2006) showed that acidic thiourea was effective for the desorption and recovery of platinum and palladium from bayberry tannin immobilized collagen fiber. Reuse

of the biosorbent was not possible due to lost biosorbent capacity when Cu (II) and Zn (II) were eluted from immobilized *Mentha arvensis* distillation waste using HCl (Hanif et al., 2009). Zhao et al. (1999) showed that biosorption using *Azolla filiculoides* could be used to treat wastewater for recycling and the recovered metal fed back to the planting bath. Ishikawa et al. (2002) identified Cu (II) and Pb (II) as metal ions which were sorbed for a time and later desorbed from egg shell membrane. On investigating the

desorption of heavy metals from water hyacinth ash, Mahmood et al. (2010) reported that desorption was useful for the recovery of cadmium, copper and lead. However, a survey of the literature shows that limited work has been done so far on the biosorptive recovery of precious metals using water hyacinth.

Conclusion

The use of water hyacinth for metal biosorption

Table 4. Desorption and recovery of selected metals.

Adsorbent	Eluant	Desorption	Recovery	Reference	Comment
Egg shell membrane	NaOH NaCN	>95% Au(I), >92% Au(III)	98%	Ishikawa et al. (2002)	No loss of biosorbent capacity was observed through five consecutive sorption/desorption cycles.
<i>Sargassum</i>	NaOH	97.8% Au(III)	91.4%	Sathishkumar et al. (2010)	Gold from dilute solutions could be recovered through green processes in ionic, nanocrystalline, or metallic form using <i>Sargassum</i> biomass.
Bayberry tannin immobilized collagen fiber	Acidic thiourea	90.7% Pt(IV) 58.4% Pt(II)		Ma et al. (2006)	Acidic thiourea was found to be more effective for the desorption and recovery of platinum and palladium.
Immobilized <i>Mentha arvensis</i> distillation waste	HCl	90% Cu(II) and Zn(II)		Hanif et al. (2009)	Re-use of the biomass was not possible due to lost biosorbent capacity
<i>Myriophyllum spicatum</i> L.	HCl	55% Cu(II), 60% Ni(II), 71 to 72% Co(II) and Zn(II)		Lesage et al. (2007)	Desorption by HCl was not suggested as a useful strategy in recovering the metals and regenerating the biomass
<i>Azolla filiculoides</i>	HCl H ₂ SO ₄		86 to 88% Zn(II) 96 to 99% Zn(II)	Zhao et al. (1999)	The technique showed that treated waste water could be recycled and the recovered zinc fed back to the planting bath.
Egg shell membrane	NaOH		17% Cu(II) 15% Pb(II) 4% Ni(II) 3% Zn(II)		Cu(II) and Pb(II) identified as metal ions which are sorbed for a time, and later desorbed.
Water hyacinth ash	HNO ₃	61 to 98% Pb(II) 70 to 97% Cd(II) 71 to 79% Cu(II) 25 to 42% Cr(III) 24 to 40% Zn(II) 12 to 21% Ni(II)		Mahmood et al. (2010)	Desorption was found to be useful for the recovery of cadmium, copper and lead.

has been reviewed. Excellent removal capabilities of the biomass for several metals of environmental concern were apparent. Factors, such as pH, temperature, adsorbent dose, etc significantly affect the biosorption capacities. On considering the invasive nature of the weed and environmental problems associated with its control, biosorption offers an attractive application of the plant. However, utilization of the biomass in industrial-scale applications is still far from being realized. More investigation is still required on structural properties of the biosorbent, multi-metal studies, mechanistic modeling, possible metal recovery, biosorbent modification and adaptation to continuous flow utilization.

ACKNOWLEDGEMENT

The International Foundation for Science, Stockholm, Sweden supported this research, through Grant Number W/4266-2 to Courtie Mahamadi.

REFERENCES

- About AAO, Kidunda RS, Osarya J (2005). Potential of water hyacinth (*Eichhornia crassipes*) in ruminant nutrition in Tanzania. *Livest. Res. Rural Dev.* 5: 17. [<http://www.log1.net/search/lrrd/lrrd17/8/abou17096.htm> (Published 5 August 2005)].
- Ali N, Chaudhary BL, Khandelwal SK (2004). Better use of water hyacinth for fuel, manure and pollution free environment. *Indian J. Environ. Prot.*, 24: 297–303.
- Al Rmalli SW, Harrington CF, Ayub M, Haris PI (2005). A biomaterial based approach for arsenic removal from water. *J. Environ. Monit.*, 7: 279–282.
- Babu RM, Sajeena A, Seetharaman K (2004). Solid substrate for production of *Alternaria alternata* conidia: A potential mycoherbicide for the control of *Eichhornia crassipes* (water hyacinth). *Weed Res.*, 44: 298–304.
- Cordery J, Will AJ, Atkinson K, Willis BA (1994). Extraction and recovery of silver from low grade liquors using microalgae. *Min. Eng.*, 7(8): 1003-1015.
- Das D, Charumathi D, Das N (2010). Combined effects of sugarcane bagasse extract and synthetic dyes on the growth and bioaccumulation properties of *Pichia fermentans* MTCC 189. *J. Hazard. Mater.*, 183: 497-505.
- Delgado M, Bigeriego M, Guardiola E (1993). Uptake of Zn, Cr and Cd by water hyacinths. *Water Res.*, 27: 269–272.
- Elangovan R, Philip L, Chandraraj K (2008). Biosorption of chromium species by aquatic weeds: Kinetics and mechanism studies, *J. Hazard. Mater.*, 152: 100–112.
- Fourest E, Volesky B (1996). Contribution of sulfonate groups and alginate to heavy metal biosorption by the dry biomass of *Sargassum fluitans*. *Environ. Sci. Technol.*, 30: 277–282.
- Gadd GM (1993). Interactions of fungi with toxic metals. *Phytologist.*, 124: 25-60.
- Goswami T, Saikia CN (1994). Water hyacinth—a potential source of raw material for greaseproof paper. *Bioresour. Technol.*, 50: 235–238.
- Hanif A, Bhatti HN, Hanif MA (2009). Removal and recovery of Cu(II) and Zn(II) using immobilized *Mentha arvensis* distillation waste biomass. *Ecol. Eng.*, 35: 1427-1434.
- Hasan SH, Ranjan D, Talat M (2010). Water hyacinth biomass (WHB) for the biosorption of hexavalent chromium: Optimization of process parameters. *Biol. Res.*, 5(2): 563-575.
- Hussain ST, Mahmood T, Malik SA (2010). Phytoremediation technologies for Ni⁺⁺ by water hyacinth. *Afr. J. Biotechnol.*, 9(50): 8648-8660.
- Ibrahim M, Kühn O, Scheytt T (2009). Molecular Spectroscopic study of water hyacinth dry matter. *Open Chem. Phys. J.*, 2: 1-6.
- Ishikawa SI, Suyama K, Arihara K, Itoh M (2002). Uptake and recovery of gold ions from electroplating wastes using eggshell membrane. *Bioresour. Technol.*, 81: 201–206.
- Julien MH, Griffiths MW (1998). Biological control of weeds. A world catalogue of agents and their target weeds, 4th edition. Wallingford: CABI Publishing; p. 223.
- Julien MH, Orapa W (1999). Successful biological control of water hyacinth (*Eichhornia crassipes*) in Papua New Guinea by the weevils *Neochetina bruchi* and *Neochetina eichhorniae* (Coleoptera: Curculionidae). In: Spencer NR, editor. Proceedings of the X international symposium on biological control of weeds. Bozeman, Montana, USA: Montana State University; p. 1027.
- Kadirvelu K, Kanmani P, Senthilkumar P, Subburam V (2004). Separation of mercury (II) from aqueous solution by adsorption onto an activated carbon prepared from *Eichhornia crassipes*. *Adsorp. Sci. Technol.*, 22, 207–222.
- Kapoor A, Viraraghavan T (1995). Fungal biosorption. An alternative treatment option for heavy metal bearing wastewaters: A review. *Bioresour. Technol.*, 53: 195–206.
- Kathiresan RM (2000). Allelopathic potential of native plants against water hyacinth. *Crop Prot.*, 19: 705–708.
- Klumpp A, Bauer K, Franz-Gerstein C, de Menezes M (2002). Variation of nutrient and metal concentrations in aquatic macrophytes along the Rio Cachoeira in Bahia (Brazil). *Environ. Int.*, 28: 165–171.
- Kwak IS, Yun YS (2010). Recovery of zero-valent gold from cyanide solution by a combined method of biosorption and incineration. *Bioresour. Technol.*, 101(22): 8587-92.
- Lesage E, Mundia C, Rousseau DPL, Van der Moortel AMK, Du Laing G, Meers E, Tack FMG, De Pauw N, Verloo MG (2007). Sorption of Co, Cu, Ni and Zn from industrial effluents by the submerged aquatic macrophytes *Myriophyllum spicatum* L. *Ecol. Eng.*, 30: 320-325.
- Ma HW, Liao XP, Liu X, Shi B (2006). Recovery of platinum (IV) and palladium(II) by bayberry tannin immobilized collagen fiber membrane from water solution. *J. Membrane Sci.* 278: 373–380.
- Mahamadi C, Nharingo T (2010a). Utilization of water hyacinth weed (*Eichhornia crassipes*) for the removal of Pb(II), Cd(II) and Zn(II) from aquatic environments: an adsorption isotherm study. *Environ. Technol.*, 31(11): 1221-1228.
- Mahamadi C, Nharingo T (2010b). Competitive adsorption of Pb(II), Cd(II) and Zn (II) ions onto *Eichhornia Crassipes* in Binary and Ternary systems. *Bioresour. Technol.*, 101(3): 859-864.
- Mahamadi C, Nharingo T (2007). Modelling the kinetics and equilibrium properties of cadmium biosorption by river green alga and water hyacinth weed. *Toxicol. Environ. Chem.*, 89(2): 297-305.
- Mahmood T, Malik SA, Hussain ST (2010c). Biosorption and recovery of heavy metals from aqueous solutions by *Eichhornia crassipes* (water hyacinth) ash. *Bioresour.*, 5(2): 1244-1256.

- Malik A (2007). Environmental challenge *vis a vis* opportunity: The case of water hyacinth. *Environ. Int.*, 33: 122-138.
- Mailu AM (2001). Preliminary assessment of the social, economic and environmental impacts of water hyacinth in the Lake Victoria basin and the status of control, biological and integrated control of water hyacinth, *Eichhornia crassipes*, ACIAR Proceedings No. 102.
- Mbati G, Neuenschwander P (2005). Biological control of three floating water weeds, *Eichhornia crassipes*, *Pistia stratiotes*, and *Salvinia molesta* in the Republic of Congo. *Biocontr.*, 50: 635-645.
- Mironga JM (2004). Geographic information systems (GIS) and remote sensing in the management of shallow tropical lakes. *Appl. Ecol. Environ. Res.*, 2: 83-103.
- Mshandete A, Kivaisi A, Rubindamayugi M, Mattiasson BO (2004). Anaerobic batch codigestion of sisal pulp and fish wastes. *Bioresour. Technol.*, 95: 19-24.
- Ogwan JA, Molo R (2004). Threat of water hyacinth resurgence after a successful biological control program. *Biocontrol Sci. Technol.*, 14: 623-626.
- Ofomaja AE, Ho YS (2008). Kinetic biosorption study of cadmium onto coconut copra meal as biosorbent. *Int. J. Environ. Pollut.*, 34: 1-4.
- Oudhia P (1999a). Medicinal weeds in rice fields of Chhattisgarh (India). *Int. Rice Res. Notes* 24: 40.
- Oudhia P (1999b). Studies on allelopathy and medicinal weeds in chickpea fields. *Int Chickpea Pigeonpea Newsl.* 6: 29-33.
- Park D, Yun YS, Park JM (2010). The past, present, and future trends of biosorption. *Biotechnol. Bioprocess Eng.*, 15: 86-102.
- Pinto CLR, Caconia A, Souza MM (1987). Utilization of water hyacinth for removal and recovery of silver from industrial wastewater. *Water Sci. Technol.*, 19: 89-101.
- Plummer ML (2005). Impact of invasive water hyacinth (*Eichhornia crassipes*) on snail hosts of schistosomiasis in Lake Victoria, East Africa. *Eco. Health*, 2: 81-86.
- Ram S, Moolani MK (2000). Herbicidal weed control of water hyacinth under semi-arid conditions. *Pestol.*, 24: 69-71.
- Sajn SA, Bulc TG, Vrhovsek D (2005). Comparison of nutrient cycling in a surfaceflow constructed wetland and in a facultative pond treating secondary effluent. *Water Sci. Technol.*, 51: 291-298.
- Sathishkumar M, Mahadevan A, Vijayaraghavan K (2010). Green Recovery of Gold through biosorption, Biocrystallization, and Pyro-Crystallization. *Ind. Eng. Chem. Res.*, 49(16): 7129-7135.
- Schneider IAH, Rubio J, Misra M, Smith RW (1995). *Eichhornia crassipes* as biosorbent for heavy metal ions. *Min. Eng.*, 8: 979-988.
- Schneider IAH, Rubio J, Smith R (2001). Biosorption of heavy metals onto plant biomass: exchange adsorption or surface precipitation. *Int. J. Min. Process.*, 62: 111-120.
- Shabana YM, Mohamed ZA (2005). Integrated control of water hyacinth with a ycoherbicide and a phenylpropanoid pathway inhibitor. *Biocontrol Sci. Technol.*, 15: 659-669.
- Shekinah P, Kadirvelu K, Kanmani P, Senthilkumar P, Subburam V (2002). Adsorption of lead(II) from aqueous solution by activated carbon prepared from *Eichhornia*. *J. Chem. Technol. Biotechnol.*, 77: 458-464.
- Szczeczek MM (1999). Suppressiveness of vermicompost against fusarium wilt of tomato. *J. Phytopathol. Phytopathologische Zeitschrift* 47: 155-161.
- Suzuki Y, Kametani T, Maruyama T (2005). Removal of heavy metals from aqueous solution by nonliving *Ulva* seaweed as biosorbent. *Water Res.*, 39(9): 1803-1808.
- Thomson S, Enala TM, Mick M (2002). Control of aquatic weeds through pollutant reduction and weed utilization: a weed management approach in the lower Kafue River of Zambia. *Physics Chem Earth, Parts A/B/C*, 27: 983-991.
- Vieira RHSF, Volesky B (2000). Biosorption: a solution to pollution? *Int. Microbiol.*, 3: 17-24.
- Volesky B (2007). Biosorption and me. *Water Res.*, 41: 4017-4029.
- Wang W (1990). Literature review on duckweed toxicity testing. *Environ. Res.*, 52: 7-22.
- Westerdahl HE, Getsinger KD (1988). Aquatic plant identification and herbicide use guide, volume II: Aquatic plants and susceptibility to herbicides. Technical report, A-88-9. Department of the Army, Waterways Experiment Station, Corps of Engineers, Vicksburg, MS.
- Yao H, Ramelow GJ (1997). Evaluation of treatment techniques for increasing the uptake of metal ions from solution by nonliving biomass derived from several strains of lichen, *Sphagnum* (peat) moss, and *Eichhornia crassipes* (water hyacinth) root. *Microbios.*, 90: 97-109.
- Zaranyika MF, Mutoko F, Murahwa H (1994). Uptake of Zn, Co, Fe and Cr by water hyacinth (*Eichhornia crassipes*) in Lake Chivero, Zimbabwe. *Sci. Total Environ.*, 153: 117-121.
- Zhang M, Ling B, Kong C, Liang G, Dong Y (2005). Allelopathic effects of lantana (*Lantana camera* L.) on water hyacinth (*Eichhornia crassipes* (Mart.) Solms). *Allelopathy, J.*, 15:125-129.
- Zhao M, Duncan JR, Van Hille RP (1999). Removal and recovery of zinc from solution and electroplating effluent using *Azolla Filiculoides*. *Wat. Res.*, 33(6): 1516-1522.
- Zheng JC, Feng HM, Lam MHW, Lam PKS, Yan-Wei Ding YW, Yu HQ (2009). Removal of Cu(II) in aqueous media by biosorption using water hyacinth roots as a biosorbent material. *J. Hazard. Mater.*, 171: 780-785.
- Zheng L, Dang Z, Yi X, Zhang H (2010). Equilibrium and kinetic studies of adsorption of Cd(II) from aqueous solution using modified corn stalk. *J. Hazard. Mater.*, 176: 650-656.