

Full Length Research Paper

Regional interpretation of river Indus water quality data using regression model

Aziz-ur-Rahman and Muhammad Owais Chughtai

Department of Energy and Environment, Hamdard University, Karachi, Pakistan.

Accepted 11 December, 2013

Water samples were collected from river Indus over 591 km² from Kashmor to Keti Bandar/Shah Bandar in the province of Sindh, Pakistan, during 2008 and 2009 on seasonal bases. These samples were analyzed for 12 water quality variables including physical and chemical parameters. Then correlation study was carried out and correlation co-efficient “r” was determined using correlation matrix to identify the highly correlated and interrelated parameters. Regression model were developed to test the significant ‘r’ and P-value test was carried out to test the significance of the pair of parameters. F-test was also used to examine the joint-effects of several independent variables without taking into account the separate effects of each variable. The comparison of the observed and predicted values of different parameters using regression equation suggested that the regression model provide useful means of rapid and easier monitoring of water quality of a river in a region.

Key words: Regression model, correlation coefficient, river indus, monitoring, physico-chemical parameters.

INTRODUCTION

Water is undoubtedly the most precious natural resource that exists on our planet. Although we recognize this fact, we disregarded it by polluting our rivers, lakes and oceans. It is the basic duty of every individual to conserve water resources (Jothivenkatachalam et al., 2010; Vega et al., 1998). The riverine system is most important resources of water supply in different countries of the world. At the source of a river, the water is relatively pure as it flows towards downstream. In Pakistan riverine system are getting polluted day by day. A primary reason for this is that all three major sources of pollution (industry, agriculture and domestic) are concentrated along the river belt and work together to reduce water quality which is a cause of alarm.

According to world wide fund for nature (WWF), five rivers in Asia serving 870 million people are among the most threatened in the world, as dams, water extraction and climate change all take their toll. The Indus River is among the top five most threatened river basins. Various

studies were carried out to investigate the quality of Indus River water (Sadia et al., 2013; Ali et al., 2004; Tassaduqe et al., 2003; Khuhawar et al., 2000).

The environment, economic growth and development of Pakistan especially in province of Sindh are highly influence by water, its regional and seasonal availability, and quality of surface water and microbiological parameters. A few number of research articles are available regarding the analysis of water quality data using regression techniques (Mulla et al., 2007; Patil and Patil, 2010, 2011; Kumar and Sinha, 2010; Obiefuna and Orazulike, 2010; Mehta, 2010; Sami et al., 2011; Jain and Sharma, 2000; Joarder et al., 2008). These parameters are closely interlinked. All the research so far reported on river water quality of Pakistan based on physico-chemical analysis, no attempt has been made to predict the river water quality by using regression analysis.

In the present study, river water quality was determined by measuring the concentration of some physico-




Figure 1. Location map. 1. Guddu Barrage; 2. Sukkur barrage; 3. Dadu-Moro Bridge; 4. Indus link 1 km downstream from Latifabad; 5. River Indus after Indus link outfall 6. Kotri Barrage at main bridge; 7. Sajawal bridge.

chemical parameters and comparing them with drinking water standards (WHO). Regression equations for the parameters having significant correlation were used to estimate the concentration of other constituents. Regression model were developed to test the significant correlation coefficient to overall water quality relative concentration of various pollutants in water to provide necessary clue for implementation of rapid water quality management.

MATERIALS AND METHODS

Water quality parameters of Indus River were monitored at seven different sampling points: Guddu Barrage; Sukkur barrage; Dadu-Moro Bridge; Indus link 1 km downstream from Latifabad; River Indus after Indus link outfall ; Kotri Barrage at main bridge and

Sajawal bridge, considering sampling locations (receiving bulk quantity of effluent either by municipal/industrial/agriculture return flow) (Figure 1).

Chemical analysis

The samples were collected using boat from the middle of the flow within 3-9 inches from the surface of water. Two to four sub samples of equal volume were also collected from vertical section. The samples were mixed well and saved in a 2.5 L glass bottle. The samples were analyzed for physicochemical parameters such as alkalinity, bi-carbonate, calcium, pH, conductivity, total dissolved solid, and total suspended solid, chloride, chemical oxygen demand, biological oxygen demand, sulfate, sodium, potassium and magnesium using standard methods (APHA, 2005). Results obtained were subjected to regression statistical analysis using Statistical Package for social Scientist (SPSS) 14th version.

Table 1a. Estimated values of water quality parameters and WHO prescribed limits (pre-monsoon).

| Parameter | ST-1 | ST-2 | ST-3 | ST-4 | ST-5 | ST-6 | ST-7 | St.Dev | WHO limit |
|--------------------------------|-------|-------|-------|-------|-------|-------|-------|---------|-----------|
| Conductivity | 0.381 | 0.323 | 0.323 | 0.417 | 0.380 | 0.396 | 0.494 | 0.059 | 0.3 |
| pH | 7.7 | 7.5 | 7.6 | 7.8 | 7.6 | 7.7 | 7.7 | 0.098 | 7 to 8.5 |
| Alkalinity | 71.5 | 61.1 | 77 | 89 | 91.2 | 45 | 76 | 16.021 | 100 |
| BOD | 28 | 24 | 22 | 33 | 57 | 21.3 | 39 | 12.711 | 06 |
| COD | 67 | 56 | 53.3 | 41.7 | 131 | 50 | 93 | 31.449 | 10 |
| TSS | 324 | 321 | 187 | 63.3 | 167 | 60.3 | 24 | 123.342 | 500 |
| TDS | 220.3 | 200.5 | 224.7 | 233.7 | 237.1 | 207.1 | 287 | 28.372 | 500 |
| Cl ⁻¹ | 40 | 38.7 | 38.4 | 39.3 | 39.3 | 37 | 50.3 | 4.454 | 200 |
| HCO ₃ ⁻¹ | 87.3 | 74.7 | 93.6 | 93.3 | 92.7 | 73.0 | 97.7 | 9.798 | - |
| SO ₄ ⁻² | 30.3 | 28 | 29.3 | 33.3 | 35 | 37.3 | 53 | 8.520 | - |
| Ca ⁺² | 35.3 | 29.3 | 35.6 | 37 | 37.7 | 37.5 | 40 | 3.358 | 100 |
| Mg ⁺² | 5.2 | 6 | 5.9 | 6.1 | 6 | 2.9 | 8.5 | 1.645 | - |
| Na ⁺¹ | 21.8 | 22 | 21.6 | 22.1 | 23 | 24.6 | 31.3 | 3.476 | - |
| K ⁺¹ | 1.9 | 1.8 | 1.8 | 2.2 | 2.2 | 3 | 6 | 1.513 | - |

Statistical screening

The regression analysis was used to estimate two water quality parameters to describe realistic ground water situations. In addition, this analysis attempts to establish the nature of the relationship between the variables and thereby provides a mechanism for prediction or forecasting (Kumar and Sinha, 2010; Agarwal and Sexena, 2011; Mulla et al., 2007). To find the relationship between two physicochemical parameters X and Y, the Karl Pearson's correlation coefficient, r is used and it is determined as follows:

$$r = \frac{n \sum xy - (\sum x \cdot \sum y)}{\sqrt{[n \sum x^2 - (\sum x)^2][n \sum y^2 - (\sum y)^2]}} \quad (1)$$

where, n = number of data points; x = values of X-variable; y = values of Y-variable.

To evaluate the straight-line by linear regression, the following equation of straight line can be used:

$$Y = aX + b \quad (2)$$

where, Y = dependent variable ; X = independent variable ; a = slope of line; b = intercept on y-axis

$$a = \frac{n \sum xy - (\sum x \cdot \sum y)}{n \sum x^2 - (\sum x)^2} \quad (3)$$

and

$$b = \bar{y} - a \bar{x} \quad (4)$$

Where, \bar{x} = arithmetic mean of all values of x; \bar{y} = arithmetic mean of all values of y.

For good correlation, value of r should be between $-1 < r < 1$.

RESULTS AND DISCUSSION

The values of physicochemical parameters, standard deviation and WHO prescribed limits are presented in Table 1a and b during pre and post-monsoon, respectively. The relationship between alkalinity with HCO₃⁻¹ is highly correlated and with BOD and Mg⁺² are moderately correlated where as alkalinity is weakly correlated with COD and TDS during pre-monsoon (Table 2a).

The relationship between alkalinity with Na⁺¹, Ca⁺² and TDS during post-monsoon which give correlation coefficient are equal to 0.448, 0.546 and 0.531, respectively (Table 2b) showing less moderate to moderate as values of 'r' lies between -1 and +1.

The terms alkalinity and hardness are often used interchangeably when discussing water quality. These aggregate properties of water share some similarities but are distinctly different. Alkalinity is a measure of the acid-neutralizing capacity of water.

Alkalinity in most natural waters is due to the presence of carbonate (CO₃⁻²), bicarbonate (HCO₃⁻¹), and hydroxyl (OH⁻¹) anions. However, borates, phosphates, silicates and other bases also contribute to alkalinity if present. This property is important when determining the suitability of water for irrigation and/or mixing some pesticides and when interpreting and controlling wastewater treatment processes.

In the present study, the alkalinity values were maximum during pre-monsoon. This may be attributed to increase in the rate of organic decomposition during which CO₂ is liberated, which reacts with water to form HCO₃⁻¹, thereby increasing total alkalinity in pre-monsoon. The increased alkalinity during pre-monsoon was due to the concentration of nutrients in water. The decrease was due to dilution caused by the rainwater during monsoon. Similar observation was reported by

Table 1b. Estimated values of water quality parameters and WHO prescribed limits (post-monsoon).

| Parameter | ST-1 | ST-2 | ST-3 | ST-4 | ST-5 | ST-6 | ST-7 | St.Dev | WHO limit |
|--------------------------------|-------|-------|-------|-------|-------|-------|-------|---------|-----------|
| Conductivity | 0.377 | 0.408 | 0.342 | 0.378 | 0.415 | 0.330 | 0.119 | 0.102 | 0.3 |
| pH | 7.8 | 7.8 | 7.8 | 7.4 | 7.6 | 7.3 | 7.4 | 0.219 | 7 to 8.5 |
| Alkalinity | 63.5 | 67.5 | 70.8 | 62.4 | 49.3 | 53 | 75.4 | 9.336 | 100 |
| BOD | 14.3 | 25.2 | 18 | 35.6 | 36 | 15.9 | 9.7 | 10.447 | 06 |
| COD | 38 | 67.3 | 40.3 | 98.3 | 67.7 | 46 | 21 | 25.572 | 10 |
| TSS | 239.7 | 260.3 | 160.3 | 153.3 | 321.3 | 210.3 | 3.3 | 101.735 | 500 |
| TDS | 229.2 | 228.6 | 234.5 | 399.2 | 276.7 | 227.8 | 805.3 | 213.055 | 500 |
| Cl ⁻¹ | 42.3 | 40.7 | 42.3 | 35 | 81.4 | 35.7 | 35.7 | 16.484 | 200 |
| HCO ₃ ⁻¹ | 75.7 | 77 | 80.3 | 82.5 | 97.1 | 80 | 101.7 | 10.239 | - |
| SO ₄ ⁻² | 40 | 43 | 39 | 38.7 | 89.4 | 34 | 87.3 | 24.262 | - |
| Ca ⁺² | 38 | 37.1 | 38.7 | 36.8 | 37.5 | 42 | 80 | 15.838 | 100 |
| Mg ⁺² | 8.3 | 7.6 | 8.4 | 6.3 | 71.6 | 5.2 | 22.7 | 24.11 | - |
| Na ⁺¹ | 20.6 | 20.3 | 22 | 19.7 | 45.3 | 18.7 | 129.3 | 40.731 | - |
| K ⁺¹ | 4 | 2.7 | 3.7 | 18.3 | 15.1 | 3.3 | 6.3 | 6.365 | - |

Table 2a. Statistical regression analysis data (pre-monsoon).

| R -Equation | I-Var. | D-Var. | n | r | F value | S.E |
|---|--------------------------------|------------|---|--------|---------|--------|
| Alkalinity = 30.650 x pH - 161.720 | pH | Alkalinity | 7 | 0.187 | 0.181 | 17.241 |
| Alkalinity = 0.234x COD+56.499 | COD | DO | 7 | 0.460 | 1.342 | 15.583 |
| Alkalinity = 0.880x BOD + 44.785 | BOD | DO | 7 | 0.698 | 4.748 | 12.569 |
| Alkalinity= - 0.012 x TSS + 74.926 | TSS | DO | 7 | -0.092 | 0.043 | 17.476 |
| Alkalinity = 0.278 x TDS +8.905 | TDS | DO | 7 | 0.493 | 1.607 | 15.267 |
| Alkalinity = 45.23 x Conductivity+55.44 | Conductivity | DO | 7 | 0.166 | 0.141 | 17.307 |
| Alkalinity = 0.841x Cl ⁻¹ + 38.97 | Cl ⁻¹ | DO | 7 | 0.234 | 0.289 | 17.063 |
| Alkalinity = 1.391 x HCO ₃ ⁻¹ - 48.66 | HCO ₃ ⁻¹ | DO | 7 | 0.850 | 13.070 | 9.232 |
| Alkalinity = 0.069 x SO ₄ ⁻² + 70.55 | SO ₄ ⁻² | DO | 7 | 0.037 | 0.007 | 17.538 |
| Alkalinity = 1.421 x Ca ⁺² + 21.74 | Ca ⁺² | DO | 7 | 0.298 | 0.487 | 16.754 |
| Alkalinity = 5.952 x Mg ⁺² + 38.450 | Mg ⁺² | DO | 7 | 0.611 | 2.982 | 13.890 |
| Alkalinity = -0.359 x Na ⁺¹ + 81.512 | Na ⁺¹ | DO | 7 | -0.078 | 0.031 | 17.497 |
| Alkalinity = -0.531x K ⁺¹ + 74.404 | K ⁺¹ | DO | 7 | -0.050 | 0.013 | 17.528 |

Table 2b. Statistical regression analysis data (Post-monsoon).

| R-Equation | I-Var. | D-Var. | n | r | F value | S.E |
|--|--------------------------------|------------|---|--------|---------|--------|
| Alkalinity = 11.134 xpH-21.328 | pH | Alkalinity | 7 | 0.262 | .367 | 9.871 |
| Alkalinity = -0.153 x COD +71.379 | COD | DO | 7 | -0.418 | 1.057 | 9.292 |
| Alkalinity=-0.477 x BOD +73.664 | BOD | DO | 7 | -0.533 | 1.989 | 8.650 |
| Alkalinity = -0.067 x TSS + 76.119 | TSS | DO | 7 | -0.735 | 5.868 | 6.937 |
| Alkalinity = 0.023 x TDS +55.142 | TDS | DO | 7 | 0.531 | 1.966 | 8.664 |
| Alkalinity = -54.03xConductivity+81.4 | Conductivity | DO | 7 | -0.588 | 2.645 | 8.271 |
| Alkalinity = -0.341 x Cl ⁻¹ + 78.395 | Cl ⁻¹ | DO | 7 | -0.603 | 2.851 | 8.161 |
| Alkalinity = 0.25 x HCO ₃ ⁻¹ + 61.024 | HCO ₃ ⁻¹ | DO | 7 | 0.027 | 0.004 | 10.224 |
| Alkalinity = -0.010 x SO ₄ ⁻² + 63.662 | SO ₄ ⁻² | DO | 7 | -0.026 | 0.003 | 10.224 |
| Alkalinity = 0.322 x Ca ⁺² + 48.880 | Ca ⁺² | DO | 7 | 0.546 | 2.119 | 8.571 |
| Alkalinity= -0.192 x Mg ⁺² + 66.689 | Mg ⁺² | DO | 7 | -0.495 | 1.620 | 8.888 |
| Alkalinity = 0.103 x Na ⁺¹ + 59.083 | Na ⁺¹ | DO | 7 | 0.448 | 1.254 | 9.145 |
| Alkalinity = -0.562 x K ⁺¹ + 67.415 | K ⁺¹ | DO | 7 | -0.383 | 0.860 | 9.447 |

(Brion, 1973; Wetzel, 1983).

Alkalinity is important because it buffers the pH of water within the system. Without this buffering capacity, small additions of acids or bases would result in significant changes of pH, which could be deleterious for aquatic life. Alkalinity also influences the distribution of some organisms within aquatic systems. The pH range was 7.5 to 7.8 during pre-monsoon and 7.3 to 7.8 during post-monsoon. The pH of most natural waters falls in the 6 to 9 range because of the bicarbonate buffering.

In the present study, the TDS and TSS values were maximum during post-monsoon and minimum during pre-monsoon. High values of TSS during post-monsoon may be due to siltation, deterioration, heavy precipitation and mixing runoff rain water which carried mud, sand, etc. Similar observations were made by Jawale and Patil (2009) and Salve and Hiware (2006).

COD is the amount of chemical oxidant required for the oxidation of the organic matter present in the waste. River Indus receives high amount of organic matter which generally originate from domestic and industrial effluents on the bank of Indus River. In the present study, the COD value vary from 41.7 to 131 during pre-monsoon and 21 to 98.3 during post-monsoon. For biodegradation, this organic waste requires oxygen, causing significant depletion of dissolved oxygen in river Indus water. The oxygen depletion not only effect biotic community of the river but also affects the purification capacity. High value of COD indicates that river received high amount of organic matter during pre monsoon. Hence the low value of alkalinity indicates that the compound responsible for decrease alkalinity is working as chemical oxidant for COD and hence significant increase in the value of COD. As regression equation, $Y = 0.477 X + 73.664$ and $Y = -0.153 X + 71.379$ was used to estimate the values of BOD and COD, respectively. It also helps to find the value of BOD/COD ratio to analyze the extent of pollution and biodegradability of water. The high value of BOD suggest that oxygen present in water is consumed by aerobic bacteria which makes fish, blind dolphins and other aquatic species to find it difficult to survive during pre monsoon.

REFERENCES

- Agarwal A, Sexena M (2011). Assessment of pollution by physico-chemical parameters using regression analysis: A case study of Gagan River at Moradabad-India. *Adv. Appl. Sci. Res.* 2 (2):185-189.
- Ali M, Salam A, Ahmed N, Bakhtywar AK, Khokhar MY (2004). Monthly Variation in Physico-Chemical Characteristics and Metal Contents of Indus River at Ghazi Ghat Muzaffargarh Pakistan. *Pak. J. Zool.* 36(4):295-300.
- APHA (2005). Standard methods for the examination of water and waste water. American Public Health Association, Washington D.C. (21st edn).
- Brion M (1973). The influence of environmental factors of the distribution of fresh water algae on experimental Study. *The role of pH, Carbon dioxide and biocarbonate system.* *J. Ecol.* 6(1):157.
- Jain CK, Sharma MK (2000). Regression analysis of groundwater quality data of Sagar District Madhya Pradesh. *Indian J. Environ. Health* 42(4):159-168.
- Jawale AK, Patil SA (2009). Physico-chemical characteristics and Phytoplanktons abundance of Mangrul Dam Dist-Jalgaon Maharashtra. *J. Aqua. Biol.* 24(1):7-12.
- Joarder M A, Raihan F, Alam J B, Hasanuzzaman S (2008). Regression analysis of Groundwater quality data of Sunamjang District, Bangladesh. *Int. J. Environ. Res.* 2(3):291-296.
- Jothivenkatachalam K, Nithya A, Chandra Mohan S (2010). Correlation analysis of drinking water quality in and around Perur block of Coimbatore District, Tamil Nadu, India. *Rasayan J. Chem.* 3(4):649-654.
- Khuhawar MY, Leghari SM, Bozdar RB, Mastoi GM, Jahanger TM (2000). Variation in water quality Assessment of River Indus from Kotri Barrage to Arabian Sea. *Proc. National Seminar on Drainage in Pakistan, Mehran University of Eng. Tech. Jamshoro August 16-18.*
- Kumar N, Sinha DK (2010). Drinking water quality management through correlation studies among various physicochemical parameters: a case study. *Int. J. Environ. Sci.* 1(2):253-259.
- Mehta KV (2010). Physicochemical characteristics and statistical study of groundwater of some places of Vadgam taluka in Banaskantha district of Gujarat state (India). *J. Chem. Pharm. Res.* 2(4):663-670.
- Mulla JG, Farooqui M, Zaheer A (2007). A correlation and regression equations among water quality parameters. *Int. J. Chem. Sci.* 5(2):943-952.
- Obiefuna GI, Orazulike DM (2010). Physicochemical characteristics of groundwater quality from Yola Area, Northeastern Nigeria. *J. Appl. Sci. Environ. Manage.* 14(1):5 - 11.
- Patil VT, Patil PR (2010). Physicochemical analysis of selected groundwater samples of Amalner town in Jalgaon district, Maharashtra, India. *Electron. J. Chem.* 7(1):111-116.
- Patil VT, Patil PR (2011). Groundwater quality of open wells and tube wells around Amalner town of Jalgaon, district, Maharashtra, India. *Electron. J. Chem.* 8(1):53-78.
- Sadia A, Feroza HW, Imran Q, Muhammad Hamid SW, Tirmizi SA, Muhammad AQ (2013). Monitoring of anthropogenic influences on underground and surface water quality of Indus River at district Mianwali-Pakistan. *Turk. J. Biochem.* 38(1):25-30.
- Salve BS, Haware CJ (2006). Studies on water quality of Wanparakalpa Reservoir Nagapur near Parli Vaijath Dist Beed Marathwada region. *J. Aquat. Biol.* 21(2):113-117.
- Sami GD, Ahmed SW, Ahmed SB, Manal HO (2011). Linear correlation analysis study of drinking water quality data for AlMukalla City, Hadrhamout, Yemen. *Int. J. Environ. Sci.* 1(7):692-1701.
- Tassaduq K, Ali M, Salam A, Latif M, Zahra T (2003). Study of the Seasonal Variations in the Physico Chemical and Biological aspects of Indus River Pakistan. *Pak. J. Biol. Sci.* 6(21):1795-1801.
- Vega M, Prado R, Barrado E, Deban L (1998). Assessment of seasonal and polluting Effects on the quality of river water by exploratory data analysis. *Water Res.* 32 (2):3581- 3592.
- Wetzel RG (1983). *Limnology*. Second Edition. Edited by Wetzel LG. Michigan State University. CRS College Publishing Philadelphia New York Chicago. p. 789.