

**Research Paper**

*Afr. J. Traditional,
Complementary and
Alternative Medicines*
www.africanethnomedicines.net

ISSN 0189-6016©2007

PHYTOCHEMICAL AND *IN VITRO* ANTIMICROBIAL ASSAY OF THE LEAF EXTRACT OF
NEWBOULDIA LAEVIS

***Usman, H. and Osuji, J. C.**

Department of Chemistry, University of Maiduguri, Maiduguri – Nigeria.

E-mail: husman321@yahoo.com

Abstract

The methanolic leaf extract of *Newbouldia laevis* was subjected to preliminary phytochemical screening and *in-vitro* antimicrobial tests. The extract revealed the presence of flavonoids, tannins, terpenes, steroidal and cardiac glycosides. The antimicrobial activity of the plant extract was assayed by the agar plate disc diffusion and nutrient broth dilution techniques. Test microorganisms were *Pseudomonas aeruginosa*, *Escherichia coli*, *Staphylococcus aureus*, *Salmonella typhi*, *Klebsiella spp.* and *Candida albicans*; all the organisms were laboratory isolates. The extract inhibited the growth of all the test organisms especially against *Klebsiella spp.* and *S. aureus* which had mean inhibition zone of 42.3 ± 1.5 and 32.3 ± 1.5 mm respectively. The results showed minimum inhibitory concentration (MIC) of 1.563 mg/ml against *Escherichia coli* and *Klebsiella spp.* and 3.125 mg/ml against *Pseudomonas aeruginosa*, *Staphylococcus aureus* and *Salmonella typhi*. The minimal bactericidal concentration (MBC) against *Escherichia coli* and *Staphylococcus aureus* was 0.39 mg/ml. This study has justified the traditional use of this plant for the treatment of stomach discomfort, diarrhea, dysentery and as a remedy for wound healing whose causative agents are some of the organisms used in this study.

Key words: Antimicrobial, Leaf Extracts, *In vitro*, Phytochemical, *Newbouldia laevis*

Introduction

The use of plants as medicine is an ancient practice common to all societies especially the African society. This practice continues to exist in the developing nations. It is on this basis that researchers keep on working on medicinal plants in order to produce/develop the best medicines for physiological uses.

Newbouldia laevis (P. Beauv) seem or boundary tree called variously as; 'Aduruku' in Hausa, 'Ogirisi' in Igbo and 'Akoko' in Yoruba languages (Hutchinson and Dalziel, 1963) is a medium sized angiosperm which belongs to the Bignoniaceae family. It grows to a height of about 7 – 8 (up to 15) metres, more usually a shrub of 2 – 3 metres, many-stemmed forming clumps of gnarled branches (Arbonnier, 2004). *Newbouldia laevis* is native to tropical Africa and grows from Guinea Savannas to dense forests, on moist and well-drained soils. It inhabits the secondary forest extending from Senegal to Cameroon, Gabon, Democratic Republic of Congo, Angola (Arbonnier, 2004).

In Nigeria, the bark is chewed and swallowed for stomach pains, diarrhea and toothache (Lewis and Manony, 1977). The plant has been found to be effective in the treatment of elephantiasis, dysentery, rheumatic swellings, syphilis, constipation, pile and as a vermifuge to round worms. It has also been found useful for earache, sore feet, chest pain, epilepsy and children's convulsion (Akunyili, 2000). The leaf, stem and fruits have been used for febrifuge; wound dressing and stomach ache (Iwu, 2000).

Earlier studies on the leaves and bark of Congolese *Newbouldia laevis* revealed the absence of flavonoids; saponins, quinones, terpenes or steroids (Oliver-Bever, 1986). Recent phytochemical studies on the root, root bark and stem of this plant revealed the presence of alkaloids, quinoid and phenylpropanoid amongst others (Gafner *et*

al., 1997; Aladesanmi *et al.*, 1998; Germann *et al.*, 2006). There was no extensive report on the presence of compounds from the leaves of this species. In this investigation, the *in vitro* antimicrobial effects of the crude methanolic leaf extract of this plant against the organisms found commonly responsible for the ailments aforementioned; including *Staphylococcus aureus* whose related infections are one of the most common cause of nosocomial (hospital acquired) infections were investigated.

Materials and Methods

The leaf of *Newbouldia laevis* was collected from the premises of City Girls Secondary School Enugu, Enugu State, Nigeria in the early morning of 11th February, 2006. The herbarium specimen was identified by Mr. A. O. Ozioko of the International Centre for Ethnomedicine and Drug Discovery (INTERCEDD), Nsukka, Enugu State, Nigeria where a voucher (No. BDCP/INTERCEDD 033) specimen was deposited. The leaves of *N. laevis* was dried under shade and pulverized into fine powder. About 300 g of the powdered form was extracted with 95% (v/v) methanol in H₂O employing the reflux method. The extract was concentrated under reduced pressure to yield a dark green mass weighing 21.1 g (7.03 % w/w). The crude extract was then coded "CME" and stored aseptically in a desiccator until required.

Phytochemical Screening

The phytochemical analyses of the crude methanolic extract was carried out in order to ascertain the presence of its constituents such as flavonoids, alkaloids, saponins, steroidal nucleus, tannins, cardio-active glycosides utilizing standard methods of analyses (Sofowora, 1993; Trease and Evans, 2002).

Organisms

Staphylococcus aureus, *Escherichia coli*, *Klebsiella spp.*, *Pseudomonas aeruginosa* and *Salmonella typhi*. The only fungus utilized was *Candida albicans*. Most of the isolates were obtained from the Department of Veterinary Microbiology & Parasitology, University of Maiduguri; *Staphylococcus aureus* and *Candida albicans* were obtained from the Department of Microbiology, University of Maiduguri Teaching Hospital, Maiduguri, Nigeria.

Antimicrobial Tests

The antimicrobial susceptibility test was conducted using the method earlier described by Sidney *et al.*, (1978); Vollekova *et al.*, (2001) with little modification by Usman *et al.*, (2005). The tests were carried out using a stock concentration of 100 mg/ml prepared by dissolving 1 g of the crude extract into 10 ml of sterile distilled water. The dilution ratio for Gram-positive bacteria and Gram-negative bacteria was 1:1000 and 1:5000 respectively using peptone water (Usman *et al.*, 2005). About 0.5 ml of the dilute cultures was aseptically inoculated on the surface of sterile Petri-dishes containing sterile solid nutrient agar. Discs impregnated with the crude extract at the concentration of 5 mg/disc were aseptically mounted on agar and thereafter incubated at 37 °C for 24 h, the inhibition zone was observed and then recorded in millimeters using a transparent metre rule. The tests were conducted in triplicate and results presented as mean ± SEM. Standard antimicrobial disc used were Amoxiclave (30 µg), Ceflunat (30 µg), Levoxine (5 µg), Ofloxacin (5 µg) and Peflotab (5 µg).

Minimum Inhibitory Concentration

MIC was defined as the lowest concentration where no visible turbidity was observed in the test tubes. The concentrations were determined as earlier described by Vollekova *et al.*, (2001) with some modification by Usman *et al.*, (2005). The MIC was determined for the micro-organisms that showed reasonable sensitivity to the test extracts. In this test, the micro-organisms were prepared using the broth dilution technique. The stock extract concentration of 100 mg/ml was made by dissolving 1 g of the extract in 10 ml of sterile distilled water and the working concentrations prepared by two-fold serial dilution technique that ranged from 0.195 mg/ml to 50 mg/ml using nutrient broth and later inoculated with 0.2 ml suspension of the test organisms. After 24 h. incubation at 37 °C, the tubes were observed for turbidity. The lowest concentrations where no turbidity were observed was determined and noted (Usman *et al.*, 2005).

Minimum Bactericidal Concentration

The minimal bactericidal concentration was determined from broth dilution test resulting from the MIC tubes as described previously (Vollekovà et al., 2001; Usman et al., 2005) by inoculating the content of each test tube on a nutrient agar plate. The plates were then incubated at 37 °C for 24 h.. The lowest concentration of the extract that showed no growth was noted and recorded as the minimum bactericidal concentration

Results and Discussions

The phytochemical screening of the crude methanolic leaf extracts of *N. laevis* revealed the presence of flavonoids, tannins, terpenes, steroidal and cardiac glycosides; alkaloids and saponins were found to be absent. These classes of compound are known to show curative activity against several pathogens and therefore could explain its use traditionally for the treatment of wide array of illnesses (Hassan et al., 2004; Usman, et al; 2005). Cardiac glycosides, flavonoids, tannins and terpenoids were detected

The *in vitro* antimicrobial screening presented in Table 1 showed the susceptibility test against Grams–positive and negative organisms and a fungal species. The extract exhibited considerable level of inhibition against all the test organisms with the highest activity on *Klebsiella* (42.3 ± 1.5 mm) and the lowest against *Candida albicans* (14.3 ± 2.0 mm). The zone of inhibition produced by most antibiotic discs against some of the organisms were found to be appreciable in relation to those activities produced by most organisms under study though not statistically comparatively to that presented by the extract. However, it is suggested that diameters of zones of inhibition ≥ 10 mm were considered active (Zwadyk et al., 1972; Usman et al., 2005). From the results of the minimum inhibitory concentration (MIC) and minimum bactericidal concentration (MBC) presented in Tables 2 and 3 respectively; it was observed that the broadest activity of the extract was against most Gram–negative organisms studied (*E. coli*, *Klebsiella spp.* and *P. aeruginosa*) while lower activity was noticed against *Candida albicans*. The extract exhibited considerable activity against *S. aureus*, a pyogenic Gram–positive bacterium known to play a significant role in invasive skin diseases including superficial and deep follicular lesion (Usman et al., 2005). It also showed appreciable activity against *E.coli*. The high antibacterial activities of this extracts could not be unrelated to the presence of the plant secondary metabolites detected. In line with these findings, it has been reported that tannins had been widely used as an application to sprains, bruises and superficial wounds (Usman et al., 2005).

Table 1: Antimicrobial susceptibility tests of Methanolic leaf extracts of *Newbouldia laevis*.

Organisms zones of inhibition (mm)							
Extract/ Standards	Conc/ disc	P. e	E.c	S.a	S.t	Kb	C.a
CME	5 mg	25.3 ± 1.5	24.0 ± 1.0	32.3 ± 1.5	24.3 ± 1.5	42.3 ± 1.5	14.3 ± 2.0
AMX	30 µg	–	*	–	10.0 ± 0.0	–	NT
CEF	30 µg	*	22.0 ± 0.0	*	*	*	NT
LEV	5 µg	23.0 ± 0.0	*	20.0 ± 0.0	30.0 ± 0.0	*	NT
OFL	5 µg	21.0 ± 0.0	35.0 ± 0.0	*	*	42.0 ± 0.0	NT
PEF	5 µg	*	31.0 ± 0.0	26.0 ± 0.0	38.0 ± 0.0	28.0 ± 0.0	NT
Distilled H ₂ O	100%	–	–	–	–	–	–

Key: CME = Crude methanolic extract; P.e = *Pseudomonas aeruginosa*; E.c = *Escherichia coli*; S.a = *Staphylococcus aureus*; S.t = *Salmonella typhi*; Kb = *Klebsiella*; C.a = *Candida albicans*; – = No activity; * = No disc; NT = not tested; AMX = Amoxiclives; CEF = Cefunat; LEV = Levoxine; OFL = Ofloxacin; PEF = Peflotab.

Table 2: Minimum inhibitory Concentration (MIC) of the Methanolic leaf extracts of *Newbouldia leavis*.

Organisms	Concentration (mg/ml)									
	0.098	0.195	0.390	0.780	1.563	3.125	6.250	12.500	25.00	50.000
P.e	+	+	+	+	+	σ	–	–	–	–
E.c	+	+	+	+	σ	–	–	–	–	–
S.a	+	+	+	+	+	σ	–	–	–	–
S.t	+	+	+	+	+	σ	–	–	–	–
Kb	+	+	+	+	σ	–	–	–	–	–
C.a	+	+	+	+	+	+	σ	–	–	–

Key: P.e = *Pseudomonas aeruginosa*; E.c = *Escherichia coli*; S.a = *Staphylococcus aureus*; S.t = *Salmonella typhi*; Kb = *Klebsiella*; C.a = *Candida albicans*; + = Turbidity observed; – = No turbidity observed; σ = MIC value.

Table 3: Minimum Bactericidal Concentrations (MBC) of the Methanolic leaf extracts of *Newbouldia leavis*.

Organisms	Concentration (mg/ml)									
	0.098	0.195	0.390	0.780	1.563	3.125	6.250	12.500	25.00	50.000
P.e	+	+	+	α	–	–	–	–	–	–
E.c	+	+	α	–	–	–	–	–	–	–
S.a	+	+	α	–	–	–	–	–	–	–
S.t	+	+	+	α	–	–	–	–	–	–
Kb	+	+	+	α	–	–	–	–	–	–
C.a	+	+	+	+	α	–	–	–	–	–

Key: P.e = *Pseudomonas aeruginosa*; E.c = *Escherichia coli*; S.a = *Staphylococcus aureus*; S.t = *Salmonella typhi*; Kb = *Klebsiella*; C.a = *Candida albicans*; + = Growth = No growth; α = MBC value.

In conclusion, the fact that the extracts produced inhibitory activities against almost all test organisms particularly Gram – negative organisms and marked higher activities (though not of comparative concentrations) than most of the reference drugs could provides some scientific basis for some of the folkloric claims. We, therefore, suggest the isolation and possible characterization of the bioactive constituent(s) from the extracts of this plant species as a possible antibacterial agent especially now that numbers of reports of methicilin-resistant *S. aureus* (MRSA) and vancomycine-resistant *S. aureus* (VRSA) is on the increase (Hiramatsu et al., 1997; Fridkin, 2001).

Acknowledgements

The authors are grateful to the Department of Chemistry, University of Maiduguri for providing some of the reagents and apparatus for the work and Messrs Samson Amali and Samson Gamache of the Department of Veterinary Microbiology University of Maiduguri and Department of Microbiology, University of Maiduguri Teaching Hospital respectively for their technical assistance.

References

- Aladesanmi, A.J, Nia, R. and Nahrstedt, A. (1998). New Pyrazole Alkaloids from the Root bark of *Newbouldia leavis*. *Planta Medica*. **64**: 90 – 91.
- Akunyili, D.N. (2000). Anticonvulsant Activity of the Ethanolic extract of *Newbouldia leavis*. 2nd NAAP Scientific conference. 155 – 8.
- Arbonnier, M. (2004). Trees, Shrubs and Lianas of West African Dry Zones. CIRAD, Margraf Publishers GMBH MNHN, Cote d'Ivoire. p. 194..
- Fridkin, S. K. (2001). Vancomycine-intermediate and Resistant *Staphylococcus aureus*: What the Infection Disease Specialist Need to Know. *Clin. Infect. Dis.* **32**: 108-15.
- Gafner, S., Wolfender, J.L., Nianga, M. and Hostettmann, K. (1997). Phenylpropanoid Glycosides from *Newbouldia laevis* Roots. *Phytochemistry* **44** (4): 687 – 690.

6. Germann, K., Kaloga, M., Ferreira, D., Marais, J.P. and Kolodziej, H. (2006). Newbouldioside A–C Phenylethanoid Glycosides from the Stembark of *Newbouldia leavis*. *Phytochem* **67** (8): 805 – 811.
7. Hassan, M.M., Oyewale, A.O., Amupitan, J.O., Abdullahi, M.S. and Okonkwo, E.M. (2004). Preliminary Phytochemical and antibacterial investigation of crude extracts of the root bark of *Detarium microcarpum*. *J. Chem. Soc. Nigeria* **29** (1): 26 – 29.
8. Hiramatsu, K.; Hanaki, H.; Ino, T.; Yabuta, K.; Oguri, T., and Tenover, F.C. (1997). MRSA Clinical Strain with reduced Vancomycine Susceptibility. *J. Antimicrochemother.* **40** (1): 135 – 136.
9. Hutchinson, J. and Dalziel, J.M. (1963). Flora of West Tropical Africa Vol. II. Crown Agents for Oversea Government and Administration 4, Millbank, London, S.W. 1. pp. 435 – 436
10. Iwu, M.M. (2000). Handbook of African Medicinal Plants. CRC Press, Inc. London p. 19.
11. Lewis, W.H. and Manony, P.F.E. (1977). Medical Botany: Plants Affecting Man's Health. John Wiley and Sons. New York, USA p. 240.
12. Oliver-Bever, B. (1986). Medicinal Plants in Tropical West Africa. Cambridge University Press, London. pp. 117-8, 168.
13. Sidney, M.F., William, J.M. and Elvyn, G.S. (1978). Bailey and Scott's Diagnostic Microbiology. C.V. Mosby: St. Louis, USA. pp. 385 – 403.
14. Sofowora, A. (1993). Screening Plants for Bioactive Agents. In: *Medicinal plants and Traditional Medicinal in Africa*. 2nd Ed. Spectrum Books Ltd, Sunshine House, Ibadan, Nigeria, pp. 134 – 156.
15. Trease, G.E. and Evans, W.C. (2002). Pharmacognosy. 15th Ed. Saunders Publishers, London. pp. 42 – 44, 221 – 229, 246 – 249, 304 – 306, 331 – 332, 391 – 393.
16. Usman, H., Haruna, A.K., Akpulu, I.N., Ilyas, M., Ahmadu, A.A. and Musa, Y.M. (2005). Phytochemical and Antimicrobial Screenings of the Leaf Extracts of *Celtis integrifolia* Lam. *J. Trop. Biosci.* **5** (2): pp. 72 – 76.
17. Vollekova, A., Kost'alova, D. and Sochorova, R. (2001). Isoquinoline alkaloids from *Mahonia aquifolium* Stembark is Active against *Malassezia spp.* *Folia Microbiol.* **46**: 107 – 111.
18. Zwadyk, P. (1972). Enterobacteriaceae in Zinsser Microbiology. 20th Ed. George Thiene Verlag, Stuttgart. pp. 20 – 32