

FLAVONOIDS FROM SUDANESE *ALBIZIA ZYGIA* (LEGUMINOSAE, SUBFAMILY MIMOSOIDEAE), A PLANT WITH ANTIMALARIAL POTENCY

Muna Ali Abdalla and Hartmut Laatsch\*

Institute of Organic and Biomolecular Chemistry, Georg-August University of Göttingen, Tammannstrasse 2, D-37077 Göttingen, Germany, <sup>2</sup>Department of Food Science and Technology, Faculty of Agriculture, University of Khartoum, Khartoum North, P.O Box 13314 Shambat, Sudan

\*E-mail: [hlaatsc@gwdg.de](mailto:hlaatsc@gwdg.de)**Abstract**

Three flavonoids were isolated for the first time from the Sudanese medicinal plants *Albizia zygia*. Compounds **1-3** were identified by interpretation of ESI mass data, <sup>1</sup>H, <sup>13</sup>C and 2D NMR as well as by comparison with published data as 4',7-dihydroxyflavanone (**1**), 3',4',7-trihydroxyflavone (**2**), 3-O-methylfisetin (3',4',7-trihydroxy-3-methoxyflavone, **3**). All flavonoids were tested against *Plasmodium falciparum*, and only compound **2** showed high antimalarial activity (IC<sub>50</sub> 0.078 µg/ml).

**Keywords:** *Albizia zygia*; Mimosoideae; flavonoids; antimalarial activity

**Introduction**

*Albizia zygia* (DC.) J.F. Macbr. (*Leguminosae* subfamily *Mimosoideae*) is a gum producing tree widely found in West Africa (Hutchinson et al., 1972). In previous phytochemical studies of *A. zygia*, lupen-20(30)-en-3β-ol and its glycoside, stigmast-5-en-3β-ol, and 5α-stigmasta-7,22-dien-3β-ol were isolated (Schoppa and Pachaly, 1981) as well as albiziaprenol and phytol were reported (Pachaly et al., 1983). Also, the gum of the plant has been widely investigated for its chemical and physical properties (e.g. as thickening agent) in comparison with other mucilages (Ashton et al., 1975; Mital et al., 1979). In traditional medicine, the powdered bark of *A. zygia* is used alone or as a decoction in southern Sudan as an antimalarial and antiparasitic drug. The methanolic extract of the stem bark exhibited antiprotozoal activity (IC<sub>50</sub> 1.0 µg/ml) against *Plasmodium falciparum* strain K1, the protozoa responsible for malaria, and *Trypanosoma brucei rhodesiense* (IC<sub>50</sub> 0.2 µg/ml), which causes African trypanosomiasis (Ndjakou et al., 2007). It is interesting to note that lupeol, which was isolated previously from this plant has been found to inhibit the growth of *P. falciparum* by 45% at 25 µg/ml (De Almeida Alves et al., 1997), which could account at least in part for the antiprotozoal activity observed for the methanol extract of *A. zygia* (Ndjakou et al., 2007).

**Material and Methods**

The bark of *A. zygia* was collected from Shambat, Sudan, in June 2006. The plant material was authenticated by Mr. Wail El Saddig (National Research Centre, Sudan) based on comparison with voucher specimen deposited in the herbarium of the Institute of the Medicinal and Aromatic Plants, Khartoum, Sudan.


An air-dried sample (1.0 kg) of the bark of *A. zygia* was powdered and extracted (3×) with 50% CH<sub>2</sub>Cl<sub>2</sub>/MeOH. The extracts were pooled and evaporated under reduced pressure to give a dark brown residue (148.6 g). This extract was partitioned between ethyl acetate and water, which delivered 7.3 g of crude product after evaporation of the organic phase, and 139.5 g from the water phase. Fractionation of the organic extract was carried out using silica gel flash chromatography (230-400 mesh) with a cyclohexane/EtOAc gradient. According to the TLC profiles three major fractions were obtained. Fraction FI was further purified on Sephadex LH-20 (Lipophilic Sephadex, Amersham Biosciences Ltd.; purchased from Sigma-Aldrich Chemie, Steinheim, Germany) using CH<sub>2</sub>Cl<sub>2</sub>/MeOH (6:4) followed by gradient chromatography on LiChroprep RP-18 (Merck KGaA, Darmstadt, Germany) eluted with MeOH/H<sub>2</sub>O (10:90 to 50:50), which afforded compound **1** (1.0 mg). Fraction FII was also separated on LiChroprep RP-18 eluted with MeOH/H<sub>2</sub>O (10:90 to 50:50) to give compounds **2** (2.7 mg) and **3** (1.7 mg).

The gum from the water phase was triturated with MeOH and the soluble part, after evaporation to dryness, was subjected to gradient chromatography on silica gel eluted with EtOAc/MeOH. This afforded compound **4** (2.9 mg).

...

**Results and Discussion**

Compounds **1-4** were unambiguously identified by interpretation of ESI mass data, <sup>1</sup>H, <sup>13</sup>C and 2D NMR spectra as well as by comparison with published data as 4',7-dihydroxyflavanone (**1**) (Umehara et al., 2009), 3',4',7-trihydroxyflavone (**2**) (Wu et al., 2008), 3-O-methylfisetin (3',4',7-trihydroxy-3-methoxyflavone, **3**) (Wu et al., 2008), and lup-20(29)-en-3-ol (**4**) (Blair et al., 1970).



The genus *Albizia* consists of over 150 species. The occurrence of albiziasaponins A, B, C (Pal, et al., 1995), macrocyclic alkaloids budmunchiamines L4, L5, and L6 (Ajay and Laxmi, 1997), and albizinin (Ma, 1997) have been reported from *Albizia lebbek*. Kaempferol and quercetin were isolated from *A. julibrissin* (Lau et al., 2007) and *A. lebbek* (El-Mousallamy, 1998). Prenylated flavonoids such as sophoflavescenol and kurarinone were also known from *A. lebbek* (Jung et al., 2004). The unusual biflavonoids, eucaediflavone and albiproflavone were delivered from *A. procera* (Yadav and Bhardoria, 2004). Other flavonoids were reported from *A. amara* and *A. adianthifolia* (Arthur et al., 1978); however, this is the first report of flavonoids from the species *A. zygia*.

In our study compounds 1-3 were tested against *P. falciparum* K1: compound 2 exhibited high antimalarial activity (IC<sub>50</sub> 0.078 µg/ml), but displayed unfortunately also cytotoxic effects (IC<sub>50</sub> 0.405 µg/ml) against a cell line L6.

The results of this phytochemical study of the bark of *A. zygia* are in agreement with previous data reported for other *Albizia* species Arthur et al., 1978).

## Acknowledgements

The authors would like to thank Dr. H. Frauendorf and Mr. R. Machinek for the mass and NMR spectra, F. Lissy and A. Kohl for technical assistance and Marcel Kaiser in Swiss Tropical Institute, Parasite Chemotherapy for determining the antimalarial activity. MAA thanks the German Academic Exchange Service (DAAD) for a Ph.D. grant.

## References

1. Ajay, K.D., Laxmi, N.M. (1997). Macrocyclic budmunchiamine alkaloids from *Albizia lebbek*. J. Nat. Prod. 60: 1036-1037.
2. Arthur, C.H., Brookes, K.B., Bull, J.R., McGarry, E.J., McGarry, J.M. (1978). Flavonoids of *Albizia adianthifolia*. Phytochemistry 17: 1681-1682.
3. Ashton, W.R., Jefferies, M., Morley, R.G., Pass, G., Phillips, G.O., Power, D.M. (1975). Physical properties and applications of aqueous solutions of African *Albizia zygia* gum. J. Sci. Food Agric. 26: 697-704.
4. Blair, J.A., Ongley, P.A., Chiswell, J., Griffiths, M.H.G. (1970). Isolation of lupeol from the bark of *Heritiera utilis* (*Tarrietia utilis*). Phytochemistry 9: 671.
5. De Almeida Alves, T.M., Nagem, T.J., de Carvalho, L.Z., Krettli, A.U., Zani, C.L. (1997). Anti-plasmodial triterpene from *Vernonia brasiliensis*. Planta Med. 63: 554-555.
6. El-Mousallamy, A.M.D. (1998). Leaf flavonoids of *Albizia lebbek*. Phytochemistry 48: 759-761.
7. Hutchinson, J., Dalziel, J.M., Keay, R.W.J. (1972). Flora of West Tropical Africa. University of Virginia Press, Virginia.
8. Jung, M.J., Kang, S.S., Jung, H.A., Kim, G.J., Choi, J.S. (2004). Isolation of flavonoids and a cerebroside from the stem bark of *Albizia julibrissin*. Arch. Pharm. Res. 27: 593-599.
9. Lau, C.S., Carrier, D.J., Beitle, R.R., Bransby, D.I., Howard, L.R., Lay Jr., J.O., Liyanage, R., Clausen, E.C. (2007). Identification and quantification of glycoside flavonoids in the energy crop *Albizia julibrissin*. Bioresource Technol. 98: 429-435.
10. Ma, Y.T. (1997). Tannins from *Albizia lobbek*. Phytochemistry 46: 1451-1452.
11. Mital, H.C., Ocran, J., Fokuo, Y.D. (1979). Studies on *Albizia zygia* gum. Acta Pharm. Technol. 25: 93-100.

- 12.Ndjakou, L.B., Vonthron-Senecheau, C., Fongang Soh, R., Tantangmo, F., Ngouela, S., Kaiser, M., Tsamo, E., Anton, R., Weniger, B. (2007). *In vitro* antiprotozoal activities and cytotoxicity of some selected Cameroonian medicinal plants. J. Ethnopharmacol. 111: 8-12.
- 13.Pachaly, P., Redeker, F., Schoppa, T. (1983). Inhaltsstoffe von *Albizzia zygia*, 2. Arch. Pharm. 316: 651-652.
- 14.Pal, B.C., Achari, B., Yoshikawa, K., Arihara, S. (1995). Saponins from *Albizia lebbbeck*. Phytochemistry 38: 1287-1291.
- 15.Schoppa, T., Pachaly, P. (1981). Inhaltsstoffe von *Albizzia zygia*. Archiv der Pharmazie 314: 18-25.
- 16.Umehara, K., Nemoto, K., Matsushita, A., Terada, E., Monthakantirat, O., De-Eknamkul, W., Miyase, T., Warashina, T., Degawa, M., Noguchi, H. (2009). Flavonoids from the heartwood of the Thai medicinal plant *Dalbergia parviflora* and their effects on estrogenic-responsive human breast cancer cells. J. Nat. Prod. 72: 2163-2168.
- 17.Wu, J-H., Tung, Y-T., Chien, S-C., Wang, S-Y., Kuo, Y-H., Shyur, L-F., Chang, S-T. (2008). Effect of phytocompounds from the heartwood of *Acacia confusa* on inflammatory mediator production. J. Agric. Food Chem. 56: 1567-1573.
- 18.Yadav, S., Bhadoria, B.K. (2004). Novel biflavonoids from the leaves of *Leucaena diversifolia* and *Albizia procera* and their protein binding efficiency. J. Indian Chem. Soc. 81: 392-394.