

EVALUATION OF *IN-VITRO* ANTIMICROBIAL ACTIVITIES AND PHYTOCHEMICAL CONSTITUENTS OF *Cassia occidentalis*

SAGANUWAN, Alhaji Saganuwan and GULUMBE, Mohammed Lawal

Department of Veterinary Physiology, Pharmacology and Biochemistry, College of Veterinary Medicine, University of Agriculture, PMB 2373, Makurdi, Benue State, Nigeria

Department of Veterinary Public Health and Animal Production, Faculty of Veterinary Medicine, Usmanu Danfodiyo University, PMB 2346, Sokoto State, Nigeria

Corresponding Author: Saganuwan, A. S. Department of Veterinary Physiology, Pharmacology and Biochemistry, College of Veterinary Medicine, University of Agriculture, PMB 2373, Makurdi, Benue State, Nigeria. Email: pharn_saga2006@yahoo.co Phone: 234-802 7444 269

ABSTRACT

The research was carried out to evaluate the in-vitro antimicrobial activity and phytochemical constituents of Cassia occidentalis. Cassia leaves were collected from Kacha town in Niger State and extracted using methanol, hexane, chloroform and water extraction methods. Serial concentrations: 50, 60, 70, 80, 90 and 100 % methanol, hexane, chloroform and aqueous extracts were prepared and sterilized. The bacterial isolates used; E. coli, P. multocida, S. typhi, S. typhimurium, S. pyogenes, S. pneumoniae and K. pneumoniae were authenticated using biochemical and serological methods. The suspension (0.5) of each bacterial isolate was prepared in isotonic sodium chloride. The disc agar diffusion method was performed on 70 Mueller-Hinton agar plates, 10 per microorganism, using serial diffusion concentration: 500, 600, 700, 800, 900 and 1000 mg of hexane, methanol, chloroform and water. The results showed that all the extracts of Cassia occidentalis have antimicrobial activity on E. coli at concentrations between 900 – 1000 mg. E. coli was most susceptible to hexane extract at concentration ranges between 500 – 1000 mg, there was no antimicrobial activity exhibited against the other tested microorganisms. Phytochemical analyses showed the presence of alkaloid, tannin, saponin, glycoside and flavonoid, steroid was absent.

Keywords: Evaluation, *In-vitro*, Antimicrobial activity, Phytochemical properties, *Cassia occidentalis*

INTRODUCTION

Cassia occidentalis (Caesalpinaceae) is a small tree growing 5 – 8 metres in height (Blumgarten, 1937). The leaves are compound, composite, paripinnate with 5 – 8 pairs of leaflets usually oval. Inflorescence occurs as auxiliary or terminal, yellow, short cluster of flower. Fruit is a pod, narrow, flat, slightly curved about 15 cm long with 10 – 12 seeds, brownish at maturity (Mann, 2003).

Traditionally, its roots, leaves, flowers and seeds are used as laxative and purgative (Todd, 1967). It is a vermifuge, anticonvulsant and used against chicken pox (Mann, 2003). Other uses include febrifuge, extrusion of guinea worm (Iwu, 1993) and black quarter (Ndi *et al.*, 2000). Previous studies have shown that its leaves exhibited *in-vitro* antibacterial, antimalarial and antihepatotoxic properties (Gasquet, 1993; Percez, 1994; Saraf, 1994). Seeds are brewed into a coffee like beverage for asthma and the flower infusion is used for bronchitis in the Peruvian Amazon (Akinloye *et al.*, 2003).

Phytochemically, the aqueous extract of *Cassia occidentalis* contained tannins, anthraquinones, sterol, cardiac glycosides, saponin and alkaloids (Muyibi *et al.*, 2000). The changing pattern of bacterial aetiology of infection and their altered sensitivities to antimicrobial agents employed in their treatment call for intensive regular exploration of indigenous plants. This will help us

identify plants with antimicrobial value that will not only serve as resource for our indigenous pharmaceutical industries but will also serve as an alternative complementary medicine (Saganuwan and Gulumbe, 2006). Orji *et al.* (2003) reported that a particular characteristic of a plant is that different chemical substances are obtained in members of even the same species in different areas.

The objectives of this study was to evaluate the *in-vitro* antimicrobial activity and photochemical constituents of *Cassia occidentalis* of Nupe land as soil nature and environmental factors like climate, weather and humidity may affect phytochemical properties of plant grown different soil textures and environments.

MATERIALS AND METHODS

Plant: The fresh leaves of *Cassia occidentalis* were collected from Katcha, the headquarter of Katcha Local Government Area of Niger State, but identified and authenticated in Herbarium of Biological Science Department of Ahmadu Bello University, Zaria, Kaduna State, Nigeria.

Extraction: The leaves of *Cassia occidentalis* were collected fresh, pounded with mortar and pestle then dried in the sun for 30 minutes to 1 hour and grounded into powder using mortar and pestle.

The extraction was carried out as described by Saganuwan and Gulumbe (2006). Serial concentrations; 50, 60, 70, 80, 90 and 100% of the hexane, methanol, chloroform and water extracts were prepared and sterilized through 0.45 µm membrane filter paper (Cheesebrough, 1985).

Collection of Bacterial Isolates: Bacterial isolates of *S. pyogenes* and *S. pneumoniae* were obtained from Microbiology Laboratory of Sokoto specialist hospital. *K. pneumoniae* and *E. coli* were supplied by Microbiology Laboratory, College of Health Sciences, Usmanu Danfodiyo University all in Sokoto. But *P. multocida* was isolated in Veterinary Public Health Laboratory, Usmanu Danfodiyo University Sokoto, whereas *S. typhi* and *S. typhimurium* were donated by Department of Veterinary Public Health and Preventive Medicine, University of Nigeria Nsukka, Nigeria. The isolates were authenticated by chemical and serological tests as described by Cheesebrough (1985), preserved on blood agar slant and stored at 4 °C until ready to use.

Disc Diffusion Method: The isolates of *K. multocida*, *S. typhi*, *S. typhimurium*, *K. pneumoniae*, *E. coli*, *S. pyogenes* and *S. pneumoniae* were subcultured overnight at 37 °C on nutrient agar plates, ten plates per microorganism. The suspensions of each bacterial isolate were prepared as described by John *et al.* (1999) in isotonic sodium chloride solution. Dried Petri dish, ten per each microorganism of Mueller-Hinton agar were flooded with the appropriate suspension of the bacterial isolates.

Sterile 6 mm diameter absorbent filter papers (punched out from No. 1 whatman paper) were impregnated with the appropriate concentrations; 500, 600, 700, 800, 900 and 1000 mg of the hexane, chloroform, methanol and water extracts and placed on the corresponding inoculated 70 plates. Ten each of *S. typhi*, *S. pyogenes*, *S. pneumoniae*, *K. pneumoniae*, *E. coli*, *P. multocida* and *S. typhimurium*. After the incubation at 37 °C for 24 hours, all the plates were observed for zones of growth inhibition and the diameters of the zones measured in millimeter (mm) using calibrated ruler.

RESULTS

E. coli showed diametric zones of inhibition at concentrations between 900 – 1000 mg of 10.0 and 17.0 mm with average mean of 4.5 mm as there was no any zone of inhibition shown on other microorganisms by the hexane extract of *Cassia occidentalis*. Diametric zones of inhibition were shown on *E. coli* by the chloroform extract at concentration ranges between 900 – 1000 mg of 10.0 and 16.0 mm with average mean of 4.3 mm. No zone of inhibition shown on other microorganisms by the chloroform extracts (Table 1).

E. coli showed diametric zones of inhibition at concentration ranges between 900 – 1000 mg of 10.0 and 15.0 mm with average mean of 4.16 mm as there was no any zone of inhibition shown on other microorganisms by the methanol extract. Diametric zones of inhibition of 8 mm and 10 mm were shown on *E. coli* at concentration ranges between 900 – 1000 mg with average mean of 3.67 mm as there was

Table 1: Serial concentrations of the hexane, chloroform, methanol and water extracts, and their corresponding diametric zones of inhibition

Conc. of the extracts (mg)	Diametric zone of inhibition (mm)														
	Hexane						Chloroform								
	S. ty	S. typhim	E. col	S. pneu	K. pneu	P. mul	S. pyo	S. ty	S. typhim	E. col	S. pneu	K. pneu	P. mul	S. pyo	
500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
600	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
700	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
800	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
900	0.0	0.0	10.0	0.0	0.0	0.0	0.0	0.0	0.0	10.0	0.0	0.0	0.0	0.0	
1000	0.0	0.0	17.0	0.0	0.0	0.0	0.0	0.0	0.0	16.0	0.0	0.0	0.0	0.0	
Mean: 750	0.0	0.0	4.5	0.0	0.0	0.0	0.0	0.0	0.0	4.3	0.0	0.0	0.0	0.0	
	Methanol						Water								
500	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
600	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
700	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
800	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	0.0	
900	0.0	0.0	10.0	0.0	0.0	0.0	0.0	0.0	0.0	8.0	0.0	0.0	0.0	0.0	
1000	0.0	0.0	15.0	0.0	0.0	0.0	0.0	0.0	0.0	14.0	0.0	0.0	0.0	0.0	
Mean: 750	0.0	0.0	4.16	0.0	0.0	0.0	0.0	0.0	0.0	3.67	0.0	0.0	0.0	0.0	

Keys: *S. ty* = *Salmonella typhi*, *S. typhim* = *Salmonella typhimurium*, *E. col* = *E. coli*, *S. pneu* = *Streptococcus pneumoniae*, *K. pneu* = *Klebsiella pneumoniae*, *P. mul* = *Pasteurella multocida*, *S. Pyo* = *Streptococcus pyogenes*

Table 2: Phytochemical constituents of *Cassia occidentalis* methanol, hexane, chloroform and aqueous leaf extracts

Extract	Alkaloid	Tannin	Glycoside	Flavonoid	Steroid	Saponin
Methanol	+	+	+	+	-	+
Hexane	+	+	+	+	-	+
Chloroform	+	+	+	+	-	+
Water	+	+	+	+	-	+

no zone of inhibition shown on other microorganisms by the aqueous extract of *Cassia occidentalis* (Table 1).

Phytochemical analyses revealed the presence of alkaloid, tannin, glycoside, flavonoid and saponin in *Cassia occidentalis* methanol, hexane, chloroform and aqueous leaf extracts as there was no steroid present in all the four extracts (Table 2).

DISCUSSION

The antimicrobial activity exhibited by hexane, chloroform, methanol and aqueous leaf extracts of *Cassia occidentalis* on *E. coli* at concentrations between 900 – 1000 mg agrees with the findings of (Gasquet, 1993; Percez, 1994; Saraf, 1994) that the leaves exhibited in-vitro antibacterial, antimalarial and antihepatotoxic properties. The plant may be used for the treatment of colibacillosis caused by *E. coli* which occurs in all species of newborn farm animals as major cause of death and economic loss in this age group (Radostits *et al.*, 2000). LeeFlang (1993) had earlier observed that indigenous knowledge and practices will be useful in the promotion of animal health and meat production in the near future in Nigeria. Furthermore, Tamboura *et al.* (2000) had also reported that ethnoveterinary medical health care will be the only alternative to western veterinary therapy. These ethnoveterinary remedies which rely on local plants or easily available materials are practical, effective and cheap (Tamboura *et al.*, 2000). However, lack of antimicrobial activities exhibited by all the extracts of *Cassia occidentalis* and generally at concentrations between 500 – 1000 mg on *P. multocida*, *S. typhi*, *S. typhimurium*, *S. pyogenes*, *S. pneumoniae* and *K. pneumoniae* is suggestive of limited antimicrobial activity of the plant. This was not pointed out by (Gasquet, 1993; Percez, 1994; Saraf, 1994). More so, uniformity of antimicrobial activity exhibited by hexane, chloroform, methanol and aqueous extracts of *Cassia occidentalis* leaf on only *E. coli* may confirm the limited antibacterial activity of the plant even *in-vivo*. Nonetheless, there is need to separate the chemical constituents of the plant leaf and then test each component on the microorganisms.

The results of qualitative analyses of all the four extracts confirmed the presence of alkaloid, tannin, glycoside and saponin as reported by Muiyibi *et al.* (2000) that the aqueous extract of *Cassia occidentalis* contained tannins, anthraquinone, sterol, cardiac glycoside, saponin and alkaloid. But there was no steroid as it also contained flavonoid.

REFERENCES

AKINLOYE, O. A., AHMED, A. S., AJAGBONA, O. P. and OLOREDE, B. R. (2003). Some biochemical effects of various doses of aqueous seed extracts of *Cassia occidentalis* in rabbits. *Bioscience Research Communication*, 15(1): 85 – 90.

- BLUMGARTEN, A. S. (1937). *Senna*. Pages 225 – 226. In: BLUMGARTEN, A. S. (Ed.) *Textbook of Material, Medical, pharmacology and Therapeutics*, 7th Edition, John Wiley and Sons. New York.
- CHEESEBROUGH, M. (1985). *Medical Laboratory Manual for Tropical Countries*, volume 17, Pages 203 – 305.
- GASQUET, M. (1993). Evaluation in-vitro and in-vivo of traditional antimalarial plants. *Fitoterapia*, 64(5): 423 – 424.
- IWU, M. M. (1993). *Catalogue of major medicinal plants. Handbook of African Medicinal plants* CRC Press, London.
- JOHN, C., KEREN, K. and BRADFORD, K. (1999). Generic protocol to estimate the burden of shigella diarrhea and dysenteric mortality. *WHO Bulletin*, 23: 28 – 30.
- LEEFLANG, P. (1993). Some observations on ethnoveterinary medicine in Northern Nigeria. *Veterinary Quarterly*, 15(2): 72 – 73.
- MANN, A., GBATE, M. and UMAR, A. N. (2003). *Senna occidentalis* (Linn.). *Medicinal and Economic Plants of Nupe land*, Jube Evan Books and Publication, Kaduna, 41 pp.
- MUYIBI, S. A., OLOREDE, B. R., AJAGBONNA, O. P. ONYEYILI, P. A., OSUNKWO, V. A. and MUHAMMED, B. Y. (2000). Acute toxicity and phytochemical studies of *cassia occidentalis* Linn extract in rats. *Sokoto Journal of Veterinary Sciences*, 2(2): 32 – 35.
- NDI, C., MUMAH, E. T., NDOKWO, K. J., MFI, A. N. NBENKUM, T., ASAH, H., EKUE, F. N. and TOYONG, J. N. (2000). Ethno veterinary plants used in the North West province of Cameroon. Pages 154 – 161. In: *Ethnoveterinary practices research and development. Proceeding of an International Workshop on Ethnoveterinary Practices*, held 14 – 18 August 2000, Kaduna Nigeria.
- ORJI, E. C., OGBWZUE, D. C. and ORJI, M. U. (2003). Evaluation of antimicrobial activities of leaf extracts of *Morinda lucida* and *Erythrina senegalensis* for medical uses. *International Journal of Medical and Health Studies*, 1(1): 79 – 83.
- PERCEZ, C. (1994). In-vitro anti-bacterial activity of argentine folk medicinal plants against *salmonella typhi*. *Journal of Ethnopharmacology*, 44(1): 41 – 46.
- RADOSTITS, M. O., GRAY, C. C., BLOOD, D. C. and HINCHLIFF, K. W. (2000). Diseases caused by bacteria. Pages 779 – 780. In: *Veterinary Medicine, A Textbook of the Diseases of Cattle, Sheep, Pigs, Goats and Horses*. W. B. Saunders, London.
- SAGANUWAN, S. A. and GULUMBE, M. L. (2006). Evaluation of *Sida acuta* Subspecies *acuta* leaf/flower combination for antimicrobial activity and phytochemical constituents. *African Journal of Clinical and Experimental Microbiology*, 7(2): 83 – 88.

- SARAF, S. (1994). Anti-hepatotoxic activity of *cassia occidentalis*. *International Journal of Pharmacology*, 322: 178 – 183.
- TAMBOURA, H. H. M., SAWADAGO, L. L., KABORE, H. and YAMEOGO, S. M. (2000). Ethnoveterinary medicine and indigenous pharmacopoeia of Pasore Province in Burkina Faso. *Annals of New York Academy of Science*, 916: 259 – 64.
- TODD, R. G. (1967). *Senna* and other purgatives. Pages 1266 – 1267. *In: Extra pharmacopoeia Martindale*, 20th Edition, W. B. Saunders, London.