

PARASITIC INCIDENCE IN CULTURED *Clarias gariepinus*

¹ADEYEMO, Abiodun Oluseye and ²FALAYE, Augustine Eyiunmi

¹Department of Fisheries Technology, Niger Delta University, Wilberforce Island, Bayelsa State

²Department of Wildlife and Fisheries Management, University of Ibadan, Nigeria

Corresponding Author: Adeyemo, A. O. Department of Fisheries Technology, Niger Delta University, Wilberforce Island, Bayelsa State. Email: text2abiiodun@yahoo.com Phone: 2348023040935

ABSTRACT

A sample of 27 fish was collected from the fish breeding centre of the Ministry of Agriculture and Natural Resources, Agodi, Ibadan. This sample was examined for parasitic incidence and the results showed 77.7 % parasitic prevalence while infection was not sex dependent (P > 0.05). Parasitic prevalence within the sample increases as length of fish increases. The prevalence of parasites encountered were Henneguya (3.7%), Dactylogyrus sp (11.1 %), Capillaria sp (7.4 %), unidentified larvae (7.4 %), unidentified cysts (40.7%), Dibothriocephalu sp (3.7 %), Argulus 7.4 %), Ichthophthirius (3.7 %) and Tricodina (3.7 %).

Keywords: Parasites, Prevalence, *Clarias gariepinus*, Infection

INTRODUCTION

Incidence of heavy parasitic infection in fish has been reported globally because fish serves as reservoir and intermediate host to most stages of parasites ranging from protozoan to metazoans (Hoffman and Meyer, 1967; Kabata, 1970; Pal and Ghosh, 1985). Several authors have worked on parasitic incidence of fish in Nigeria (Ukoli, 1963; Alfred-Ockiya, 1985; Awa, *et al*, 1988; Okaeme, 1991 and Adeyemo, *et al*, 2003) and discovered that in the natural environment healthy individuals co-exist with diseased ones and in most parasitic infections host may not be killed unless the parasitic burden is high. But usually, growth rate and market value of fish may be reduced while infection may also be of public health importance. For public health concerns, it is necessary to identify disease reservoirs in order to have adequate knowledge of the transmission mechanism. This will help to develop an effective method of preventing the access of pathogens and their reservoirs to healthy facilities or individuals.

One of the culturable fish species in Nigeria is *Clarias gariepinus*; it is widely accepted for consumption and rearing. The study of parasitic incidence on this species will further help to understand its adaptation for culture purpose. This study will also add to the current knowledge on parasitic infections of cultured fish in Nigeria.

MATERIALS AND METHODS

The parasitic investigation was carried out on *C. gariepinus* that were sampled from the Fish Breeding Centre of the Ministry of Agric and Natural Resources, Agodi, Ibadan. The centre is located between latitude 7°25' N and longitude 3° 33'W. Sampling was carried out using the method of Ossiander and Wedemeyer (1973). A total of 27 live fishes was collected during the raining season period between the months of May – July. They were examined for parasites using the identification outline of Amlacher (1966), and Pal and

Ghosh (1985). The individual fish were demobilized by pithing, while the organs were carefully examined using hand lens. Scrapping from the skin and gills were placed in Petri dishes with 2 drops of normal saline for examination, while the squash preparations from the organs were also examined under the low power (X10) magnification of a binocular microscope.

Prevalence and intensity were calculated using the indices of Margolis *et al*. (1982). Length range frequency in relation to prevalence within the sample was analyzed. The dependence of infection on sex was statistically determined using chi² analysis. The condition factor (K) was calculated and defined as $K = 100w/L^3$ where w is the weight and L as total length in cm (Bakare, 1970).

RESULTS AND DISCUSSION

Parasitic prevalence of 3.7 % was recorded for *Henneguya sp* a sporozoan parasite, which was found, encysted on the air sac of *C. gariepinus* (Table I). The parasite was discovered when the cyst was teased to release the content. Meanwhile, other unidentified cysts were also found at the prevalence rate of 40.7 % in the musculature, on the skin, ovary and kidney of the fish. These cysts could be explained to arise through connective tissue reaction to sporozoan infection and sometimes could be larvae of helminthes being encapsulated by connective tissue, which may eventually calcify. (Amlacher, 1966). The importance of these cysts lies in quarantine procedure, since after 3 – 4 weeks of quarantine, the cyst may burst to release the pathogenic exudates or helminthes. Prevalence rate of 11.1 % was recorded for *Dactylogyrus sp* a monogenetic trematode. This trematode was described by Hendrix (1994), to be cosmopolitan in nature and could be ecto or endoparasite. Awa, *et al*. (1988) also reported incidence of monogenetic worms on *Sarotherodon galileans* at the Ikoyi fish farm, Lagos.

Table 1: Parasitic incidence on cultured *Clarias gariepinus*

Parasites Found	Number of fish examined	Number of Infected fish	Parasite Burden	Prevalence (%)	Intensity of infection	Site infected
<i>Heneguya sp.</i>	27	*1	High	3.7	High	Air Sac
<i>Unidentified Cysts</i>	27	*11	50	40.7	4.7	Muscles, Skin, body cavity, Ovary, kidney
<i>Dactylogyrus sp</i>	27	*3	8	11.1	2.6	Intestine
<i>Dibothriocephalus sp</i>	27	1	1	3.7	1	Intestine
<i>Capillaria sp</i>	27	2	3	7.4	1.5	Stomach, body cavity
<i>Unidentified larvae</i>	27	2	5	7.4	2.5	Body surface
<i>Argulus sp</i>	27	2	2	7.4	1	Body surface
<i>Ichthophthirius sp</i>	27	1	Low	3.7	Low	Skin body surface
<i>Tricodina sp</i>	27	1	3	3.7	3	Stomach

* Parasite was found in more than one fish examined

Table 2: Sex ratio analysis of infected *Clarias gariepinus*

	Female	Male	Sex ratio
Number of fish examined	13	14	1:1.01
Number of fish infected	9	12	1:1.02
Prevalence	69	86	1:1.3

Table 3: Length-weight in relation to infection and condition factor (k)

Length Range (cm)	Weight average (gm)	No within the group	No of infected Fish	Prevalence Within the Group (%)	Condition Factor (k)
22-23.5	103.33	6	5	83	0.8
24-25.5	120.90	11	7	63	0.7
26-27.5	191.66	6	5	83	0.9
28-29.5	152.25	2	2	100	0.6
30-31.5	265.00	2	2	100	0.8
Total		27	21	77.7	

The dominant species among these worms are namely *Dactylogyrus sp* and *Gyrodactylus sp*. *Dactylogyrus vastator* had been known to cause heavy mortalities of fry and fingerlings (Sarig, 1971). The occurrence of fish mortality at Nagpur, India was found to be due to infection by cestode as reported by Pal and Ghosh (1985). This present study recorded the prevalence of *Dibothriocephalus sp* at 3.7 %, this tapeworm was described by Needham and Wooten (1978) as broad fish tapeworm of man, which is known to occur in a variety of fresh water fish species. It is also of public health importance, because tapeworm of *Diphylobothrium sp* has been found to be of zoonotic relevance. (USDHEW, 1973).

The unidentified larvae, *Capillaria sp* and *Argulus sp* were found at 7.4% prevalence rate respectively. These were known to cause skin irritation, which favours secondary infections and may lead to transmission of bacterial hemorrhagic septicemia (Moore *et al*, 1984). The two common protozoan, *Ichthophthirius sp* and *Tricodina sp* were found from the skin scrapings and stomach contents respectively, but there was no lesions observed on any of the samples. The parasitic infection on *Clarias sp* in this centre was found not sex dependent ($P > 0.05$) because dependence on sex was not statistically significant using χ^2 analysis. Sex ratio analysis of infected fish and the infection prevalence according to sex is shown in Table 2. Prevalence of infection according to length increases within the sample as length of fish increases while the overall

prevalence was 77.7 %, also the effect of infection on fish condition factor was low as shown in Table 3. The condition factor (K) measures the well being of the fish and is usually close to value 3 for a healthy fish, in this study the K values were below value 1 indicating that all specimens were not healthy considering the high (77.7 %) parasite prevalence at the center.

Conclusion: This report would not have succeeded in identifying all the parasites that may likely be found on cultured *Clarias species*. It however advocates for more work on diseases and parasites of fish especially culturable fish species. Since temperature regime in our ecological waters favours rapid multiplication and cycling of both fish and parasites. Meanwhile, aquaculture is developing progressively at a high rate in the country. Wild fish that are sometimes stocked or strayed into rearing ponds may bring about incidence of transferred infection, fish mortalities and consequent loss of production. It will also attract public health concern when infected fish are improperly prepared for consumption.

REFERENCES

- ADEYEMO, A. O., AGBEDE, S. A., TAIWO, V. O. and ADEDEJI, B. O. (2003). Prevalence, abundance and intensity of *Clinostomum*

- tilapiae* on cultured *Oreochromis niloticus*. *Tropical Veterinarian*, 21(3): 129 – 133.
- ALFRED-OCKIYA, J. F. (1985). Preliminary survey of parasites of *Clarias* sp. in the Delta area of Nigeria. Pages 210 – 215. In: ITA, E. O. (Ed). *Proceedings of the 4th Annual Conference of Fisheries Society of Nigeria (FISON)*.
- AMLACHER, E. (1966) *Textbook of Fish Diseases*. Oliver and Boyd Limited, Aberdeen. 310 pp.
- AWA, J. N., ANYANWU, P. and EZENWA, B. (1988). Incidence of parasites infection of pond raised *Tilapia* sp and some cultivable fish species from three ecological areas of Lagos State. Pages 20 – 21. In: Nigerian Institute for Oceanography and Marine Research (NIOMR) 1988 Annual Report.
- BAKARE, O. (1970). Bottom deposits as food of inland fresh water fish in Kainji a Nigeria Man-made lake. Volume 1, Pages 65 – 86. In: VISSER S. A. (Ed). *The Nigeria Institute of Social and Economic Research*.
- HENDRIX, S. S. (1994). *Marine Flora and Fauna of the Eastern United States*. National Oceanic and Atmospheric Administration (NOAA) Technical Report. National Marine Fisheries Services (NMFS) 121. US Department of Commerce, US. 106 pp.
- HOFFMAN, G. L and MEYER, F. P. (1967). *Parasites of freshwater fishes: A review of their control and treatment*. TFH, New Jersey. 224 pp.
- KABATA, Z. (1970). Crustacean as enemy of fishes. In: Sniezko, S. F and Axel, R. (Eds.). *Diseases of fishes*. TFH, New Jersey.
- MARGOLIS, L., ESCH, G. W., HOLMES, J. C., KURRIS, A. M. and SHAD, G. A. (1982). The use of ecological terms in parasitology. *Journal of Parasitology*, 68: 131 – 133.
- MOORE, B. R., MITCHELL, A. J., GRIFFIN, B. R. and HOFFMAN, G. L. (1984). Parasites and diseases of pond fishes. Pages 177 – 205. *3^d Annual Report of United States Department of Interior Fisheries and Wildlife Services*.
- NEEDHAM, T. and WOOTEN, R. (1978). Parasitology of Teleost. Pages 144 – 182. In: ROBERTS R.J (Ed). *Fish Pathology*. Cassell Limited. Great Britain.
- OKAEME, A. N. (1991). Helminthes fauna of tilapia of lake Kainji in pre and post impounded conditions. *Journal of Aquaculture*, 6(1): 1 – 8.
- OSSIANDER, F. and WEDEMEYER, G. (1973). Computer programme for sample sizes required to determine disease incidence in fish populations. *Journal of the Fishery Resources Board of Canada*, 30(9): 1383 – 1384.
- PAL, R. N. and GHOSH, A. K. (1985). *Identification and control of commonly occurring diseases in fresh water aquaculture*. Aquaculture Extension Manual Series, Volume 9, Central Inland Fish Institute, India. 23 pp.
- SARIG, S. (1971). *Prevention and treatment of warm water fish diseases under sub-tropical conditions with special emphasis on intensive fish farming*. TFH, New Jersey.
- UKOLI, F. M. A. (1963). Preliminary reports on the helminthes infection of fishes in River Niger at Shagumu. Pages 70 – 73. In: White E. (Ed). *1st Scientific Report of Kainji lake Biological Research Team*.
- USDHEW (1973). *Epidemiological aspects of some of the zoonoses*. Centre for Disease Control Public Health Service of United States Department of health, Education and Welfare (USDHEW). No CDC 75-8182.