

## REPLACEMENT VALUE OF GUINEA CORN FOR MAIZE IN PRACTICAL DIET FED TO QUAIL (*Coturnix coturnix japonica*) CHICKS

EDACHE, John Amedu, MUSA, Usman, HARUNA, Emmanuel Sheyin, ESILONU, John Obinna, JIBRIN, Ibrahim, OKPALA, Eugene Jideofor and DOGO, Iliya Goni

Livestock Investigation Department, National Veterinary Research Institute, PMB I, Vom, Plateau State

**Corresponding Author:** Edache, J. A., Livestock Investigation Department, National Veterinary Research Institute, PMB I, Vom, Plateau State

### ABSTRACT

A six week feeding trial was conducted to determine the replacement value of Guinea corn for maize in diet fed to 360, day-old quail chicks on deep litter. Four iso-nitrogenous (22 % Crude protein) diets incorporating graded levels (0, 15, 27 and 42 %) of guinea corn as replacement for maize were used in the trial. The ME levels of the diets ranged from 2700 – 2750 kcal/kg. Each treatment was replicated thrice. Feed intake, weight gain and feed/ weight gain ratio did not differ significantly ( $P>0.05$ ) among the treatments. Feed cost decreased across the treatments and was lowest for the diet in which 42 % maize was replaced by guinea corn. Feed cost/kg weight gain was lower for diet B(15 % guinea corn) than for other diet tested. Results of this study indicated that a dietary crude protein level of 22 % and ME of 2700 to 2750 kcal/kg feed, of 42 % guinea corn based diet was suitable for growth of Japanese quail chicks.

**Keywords:** Guinea corn, Quail chicks, Productive Performance

### INTRODUCTION

Much care is being exercised by many people on the quantity of animal fat consumed regularly because of the health implications. Japanese quail are small-bodied birds of the galliformiss family and are low in body fat and cholesterol (Garwood and Diehl, 1987, Schwartz and Allen, 1981). Therefore, much effort should be targeted to multiplying these birds for increased consumption of meat with low fat and high protein.

Livestock feed have become very expensive resulting in decrease in livestock production. There is increasing competition between man and livestock for available feedstuffs for food, feed and industrial raw materials. According to Bamgbose, *et al.* (2004), maize accounts for about 45 to 55 % of poultry feed. Therefore any effort to substitute maize in poultry feed will significantly reduce the cost of production. They successfully replaced 40 % maize with maize offal/cashew nut meal based diet and recorded no deleterious effect on carcass yield and nutrient digestibilities of broilers. The most relevant option to arrest the present feed crisis of the livestock industry is by-product utilization (Atteh, 1986). These deductions point clearly to alternative feed stuff for livestock feed productions in order to cut down feed prices and make them more affordable by livestock farmers. Olubamiwa, *et al.* (1999) had also successfully replaced 14 % maize with cocoa husk meal (CHM) with no depressive effect on the growth of quail chicks. Cullison, (1987), reported that sorghum can replace 50 % of corn with no adverse effect on animal performance but weight gain may reduce by 10 % or more with higher levels of replacement. This is contrary to the report of Spiridon, *et al.* (1979) who observed no depressive effect of sorghum on growth and feed efficiency even

at 100 % replacement of maize with sorghum in meat chickens. However, carcasses of birds fed most sorghum diets were lighter than control. Guinea corn contains 11 % crude protein, 3300 kcal/kg metabolizable energy and a crude fibre level of 3 % (Aduku, 1992). Reports of Lee *et al.* (1981) and Haruna, *et al.* (1997) had recommended crude protein levels of 24 % and 22 to 25 % respectively for quail chicks. This was contrary to the recommended level of 28 % crude protein (NRC, 1971). The work of Olubamiwa, *et al.* (1999) also recommended metabolizable energy levels of between 2,500 and 2,800-kcal/kg diet for growing quails.

This study investigated the effect of replacing maize with guinea corn in quail chick diet.

### MATERIALS AND METHODS

**Birds:** Three hundred and sixty (360) day old Japanese quail (*Coturnix coturnix japonica*) chicks hatched at National Veterinary Research Institute, Vom Poultry farm were selected on the basis of fitness and uniformity and bodyweight. They were housed in pens in a standard poultry brooding house and spaced 75 sq cm per bird as recommended (NVRI, 1996). Each pen housed 30 unsexed quail chicks. In all there were 12 experimental pens each fitted with 100-watt electric bulbs. Two kerosene stoves (modernized) heated the entire room. Cardboard sheets designed to keep the chicks from straying away from the heat source were put in each pen. These measures were to achieve the desired brooding conditions. All chicks were weighed together in groups before they were placed in the pens. Subsequently, 20 percent of the chicks were weighed from each pen at weekly interval (for six weeks).

**Table 1: Experimental diets (%)**

Ingredients	A	B	C	D
1. Maize	42	27	15	-
2. Guinea corn	15	27	42	
3. FF Soya	10	10	10	10
4. Wheat offal	20	20	20	20
5. Fish meal	1	1	1	1
6. Soya cake	23.8	23.8	23.8	23.8
7. Bone meal	2.0	2.0	2.0	2.0
8. Limestone	0.5	0.5	0.5	0.5
9. Common salt	0.25	0.25	0.25	0.25
10. Premix	0.25	0.25	0.25	0.25
11. Methionine	0.10	0.10	0.10	0.10
12. Lysine	0.10	0.10	0.10	0.10
	100	100	100	100
<b>Proximate Composition</b>				
CP	21.86	22.16	22.40	22.70
ME	2750	2730	2714	2695 (Kcal/kg)
Ca	1.07	1.51	1.51	1.09
P	0.63	0.67	0.70	0.72
<b>Feed cost/100 kg</b>	3327.75	3252.75	3192.75	3117.75
<b>Analysed Composition</b>				
CP	22.20	22.15	22.30	22.50
Ca	1.11	1.21	1.4	1.15
P	0.61	0.64	0.67	0.69

**Table 2: Effect of different guinea corn levels on mean feed consumption, weight gain, and feed efficiency of quail chicks at 6 weeks of age**

	A	B	C	D	SEM
<b>Feed Consumption</b>	382.6	383.16	365.66	373.84	± 14.19
<b>Weight gain</b>	128.52	140.43	20.52	129.97	± 4.62
<b>Feed/gain ratio</b>	3.38	3.14	3.38	4.84	± 1.43
<b>Feed cost (₦/kg)</b>	33.28	32.53	31.93	31.18	..
<b>Initial weight (g/bird)</b>	9.03	8.78	9.03	8.93	..
<b>Final weight (g/bird)</b>	137.56	149.11	130.89	138.89	..
<b>Cost/kg gain (₦)</b>	97.87	94.23	104.94	134.81	±24.47

Three groups each were randomly allocated the experimental feeds (B, C and D) while control group was fed with feed containing no guinea corn (0 level). Feed and water were given *ad. libitum*.

**Guinea Corn:** 100 kg was obtained from the open market in Vom, Plateau State and was verified to be in good condition, free from weevils.

**Experimental Diets:** Four experimental diets were formulated to contain graded levels of guinea corn (0, 15, 27 and 42) at the expense of maize. The diets represented by A, B, C, and D respectively were iso-nitrogenous containing 22 % crude protein. The energy levels of the diets ranged between 2700 to 2750 kcal/kg metabolizable energy. All experimental diets were analysed for proximate chemical compositions (Table 1) (AOAC, 1970).

**Data Collection:** The mean weekly body weight and feed consumption of birds were recorded throughout the experimental period. From the mean body weights and feed intake, feed conversion ratio was calculated. Feed cost /kg diet was calculated using

the prevailing market price of feed ingredients around Jos. Data collected were subjected to two way Analysis of variance (ANOVA) (Steel and Torrie, 1980).

## RESULTS AND DISCUSSION

Data on feed consumption, weight gain and feed/weight gain ratio are shown in Tables 2. All the groups on the different treatments started to lay by the 6<sup>th</sup> week of life which confirms the report of Martins (1987). Even though there were numerical differences between diets in terms of the different parameters measured, they were not significantly different ( $P > 0.05$ ). There were numerical differences in feed intake between treatment diets but the differences were not significant even though the energy levels were slightly decreasing across the diets from diet A. This may be attributed to the observed ability of quail to adjust feed intake over a wide range of dietary energy content (Olubamiwa, *et al.*, 1999). Even though there were seeming differences in weight gain of quail chicks (Table 2), the differences were not significant. The energy

contents of the diets decreased from 2,750 (diet A) to 2,695 (diet D) and apparently did not affect weight gain of quail chicks. Weber and Reid (1967) had reported that quail weight gain on 1760- 2400 kcal/kg ME diets was not significantly altered. Feed conversion efficiency of quail (table 2) did not differ significantly between treatments, though numerically it was highest for diet D and lowest for diet B. It is interesting that quail chicks did not register depressed weight gain, feed consumption or feed conversion even at 42 % guinea corn inclusion in the diet. The lack of depressive effect of the treatment diets on the productive parameters agrees with the report of Spiridon, *et al.* (1979). The tolerance to increasing guinea corn in the diets may be due to the ability of quails to adjust their eating habits. The feed conversion rate for quail was poor when compared to reports for chickens (Sobamiwa and Longe, 1998). This generally poor feed conversion had also been reported by Weber and Reid (1967) and Haruna, *et al.* (1997). This may be due to feed wastage characteristic of quail birds. In terms of feed cost/kg diet, diet D was the cheapest (Table 2). Feed cost/ kg weight gain was numerically lower for diet D than for other diets and compared favourably with those reported by Olubamiwa, *et al.* (1999). It is important to investigate the upper limit of guinea corn for quail chicks.

#### ACKNOWLEDGEMENT

The authors are grateful to the Director National Veterinary Research Institute, Vom for creating an enabling environment and for permission to publish this paper. We are also grateful to God for his grace.

#### REFERENCES

- AOAC (1970). *Official methods of analysis*. 10<sup>th</sup> Edition. Association of Official Analytical Chemist Inc. Association of Official Analytical Chemistry, Washington DC
- ADUKU, A. O (1992). Practical lives stock feeds production in the tropics. S. Asekome and Company, 49 pp.
- ATTEH, J. O. (1986). *Towards animal protein self-sufficiency. A case of increased production of monogastric animals*. The University of Ilorin Farmer. 2<sup>nd</sup> Edition. University of Ilorin, Nigeria. 12 pp.
- BAMGBOSE, A. M., OGUNGBENRO, S. D., OBASOHAN, E. E., ARUNA, M. B., OTEKU I. T., IGENE, U. F., OTOIKHIAN, C. S. O. and J. A. IMASUEN (2004). Replacement value of maize in ofal / cashew nut for maize in broiler diet. *Proceedings of the 29<sup>th</sup> Annual conference of the Nigerian society for Animal Production*. 29; 219 – 221.
- CULLISON E. A. (1987). *Air-dry energy feeds. Feeds and feeding*. 2<sup>nd</sup> Edition. Reston Publishing Company Incorporated. pp 161 – 163.
- GARWOOD, V. A. and DIEHL, R. C. J. (1987). Body volume and density of live coturnix quail and associated genetic relationships. *Poultry Science*, 66(8): 1269
- HARUNA, E. S., MUSA, U., OKEWOLE, P. A., SHAMAKI, D., LOMBIN, L. H., MOLOKWU, J. U., EDACHE, J. A. and KARSIN, P. D. (1997). Protein requirement of quail chicks in Plateau State, Nigeria. *Nigerian Veterinary Journal*, 18: 108 – 113.
- LEE, T. K., SHIN, K. F. and TAN, E. L., (1981). Protein requirement of breeding Japanese quail in the tropics. *Singapore Journal of Primary Industries*, 9(2): 101 – 110.
- MARTINS, B. (1987). Guinness world birds. Guinness Superlatives Limited. 158 pp.
- NRC (1971). *Nutrient requirement of domestic animals. No. 1. Nutrient requirements of poultry*. 6<sup>th</sup> Edition, Washington DC National Academy of Science, National Research Council. 233 pp
- NVRI (1996). Manual of quail production and Health management, National Veterinary Research Institute Publication, Vom. 43 pp.
- OLUBAMIWA, O., HARUNA E. S., MUSA, U., AKINWALE, T. O., LOMBIN, L. H., and LONGE, O. G. (1999). Effect of different energy levels of cocoa husk-based diets on productive performance of Japanese quails. *Nigerian Journal of Animal Production*, 26: 88 – 92.
- SCHWARTZ, R. W. and ALLEN, N. K. (1981). The effect of aging on the protein requirement of mature female Japanese quail for egg production. *Japanese Poultry Science*, 60(2): 342 - 348.
- SPIRIDON, G. DEXAMIR, A. INCULET, E., VISAN, I. POPESCU, A. and CIUPERCESCU, V. (1979). *Efficiency of mixed feeds with different amounts of maize and sorghum for meat chickens*. Institute of Agronomy, Nicolae Balcescu, Bucharest, Romania. pp 91 – 94.
- SOBAMIWA, O. and LONGE, O. G. (1998). An assessment of the optimal level of cocoa husk in broiler starter diets. Pages 161 – 162. In: ODUGUWA, O. O., FANIMO A. O and OSINOWO O. A. (Eds). *Proceedings of the Silver Anniversary Conference of the Nigerian Society of Animal Producers and Inaugural Conference of the West African Society of Animal Producers*. Abeokuta, Nigeria. March 1998.
- STEEL, R. G. D. and TORRIE, J. H. (1980). *Principles and procedures of statistics*. McGraw – Hill Book Company Incorporated, New York. 456 pp.
- WEBER, C .W. and REID, B. L. (1967). Protein requirements of coturnix quail chicks at 5 weeks of age. *Poultry Science*, 46: 1190 – 1194.