

GROWTH AND SEED YIELD RESPONSES OF SUNFLOWER (*Helianthus annuus* L.) TO ORGANIC AND/OR INORGANIC FERTILIZERS IN THE DERIVED SAVANNAH

*Ndukwu R.C. and Ogbonna P.E.

Department of Crop Science, University of Nigeria, Nsukka 410001, Nigeria

*Corresponding author's email: ruchysimeon@gmail.com; Mobile: +234 816 516 6534

ABSTRACT

*A field experiment was conducted at the Department of Crop Science Teaching & Research Farm, University of Nigeria, Nsukka to determine the effects of poultry manure and NPK 20:10:10 on the growth and seed yield of sunflower (*Helianthus annuus* L.) in Nsukka, southeastern Nigeria. The factors were poultry manure at the rates of 0, 5, 10, and 15 t ha⁻¹ and NPK 20:10:10 at 0, 100, 200, and 300 kg ha⁻¹. The experiment was laid out in a 4 × 4 factorial in a randomized complete block design with three replications. Parameters measured were plant height, stem girth, number of leaves, days to 50% and 100% flowering, head diameter, head weight and seed weight per plant. Stem girth, number of leaves, head diameter and head weight increased at the application of poultry manure. Plant height and seed weight were significantly ($p < 0.05$) affected by poultry manure application. Application of 15 t ha⁻¹ poultry manure gave similar seed yield per plant as 10 t ha⁻¹ but higher than the other rates. Days to 50% and 100% flowering decreased at the application of poultry manure. All parameters were significantly affected by NPK 20:10:10, increasing with increase in the rate of NPK 20:10:10. The application of 300 kg ha⁻¹ NPK 20:10:10 produced the highest seed yield compared to other rates. The 0 kg ha⁻¹ NPK 20:10:10 (control) gave the lowest value for all parameters measured and the highest days to 50% and 100% flowering. Poultry manure and NPK 20:10:10 interaction consistently showed non-significant effects; however, applying them at 15 t ha⁻¹ and 300 kg ha⁻¹, respectively gave the highest values for all parameters. Therefore, they should be combined at these rates, otherwise sole application of either poultry manure at 15 t ha⁻¹ or NPK 20:10:10 at 300 kg ha⁻¹ is recommended for optimum growth and production of sunflower in the derived savannah ecological zone.*

Key words: sunflower, fertilizer use, seed yield, southeastern Nigeria

INTRODUCTION

Sunflower (*Helianthus annuus* L.) stands as a key global oil crop. Originating from North and Central America, the sunflower belongs to the Helianthus genus and is notably recognized for its round head, which, when combined with its ligules, resembles the sun (Putt, 1997; Schilling and Edward, 2006). This and other varieties, particularly Jerusalem artichoke (*H. tuberosus*), are grown in temperate and certain tropical areas. They serve as food crops for humans, cattle, and poultry, and are also cultivated for ornamental purposes (Balogh, 2008). Sunflowers belong to the Asteraceae family, the largest plant family with nearly 1,550 genera and around 24,000 species. The Helianthus genus, within this family, consists of 51 species, comprising 14 annual and 37 perennial varieties.

The productivity of sunflowers is subject to the interplay of various factors, as highlighted by Kaleem *et al.* (2011). Cultural practices, including sowing timing and fertilizer application, along with environmental factors such as temperature and rainfall,

play pivotal roles in shaping the growth and development of sunflowers. Sunflower species can reach heights of 300 cm or more. Each "flower" is essentially a disc composed of tiny flowers, creating a larger false flower to enhance attraction to pollinators (Sommerlad, 2022). The cultivated sunflower holds a significant position alongside soybean, rapeseed, and peanut as one of the four major annual crops worldwide cultivated for edible oil. Its leaves serve as fodder, while the sunflower oil cake is utilized in livestock feed. The extracted oil finds applications in soap, paints, and as a lubricant. Sunflower seeds, whether dried, roasted, or ground into nut butter, are commonly included in birdseed mixes (Pilorgé, 2020).

The development of a sunflower is predominantly influenced by its genetic composition and origin. Moreover, the timing of its planting exerts notable effects on its growth. The species *H. annuus* generally thrives during the summer, reaching its peak growth phase in mid-summer (Dorling, 2008). The ideal sowing time for sunflowers can vary across

locations due to diverse climatic factors. Research conducted in Nigeria revealed that the timing of sowing (late July to mid-August) significantly impacted various yield parameters and the overall growth of sunflowers (Ali *et al.*, 2014). Sunflowers, known for their affinity to sunlight, thrive best in locations receiving six to eight hours of direct sun daily. Planting sunflower seeds is optimal when the soil temperature reaches at least 60 °C at a depth not exceeding one inch. These plants possess extensive tap roots that extend several feet into the ground, making them prefer loose, well-drained soil with a somewhat alkaline pH ranging from 6.5 to 7.5 (Vasyl, 2023). The growth of sunflowers undergoes distinct stages, categorized as vegetative and reproductive. Identification of heads or main branches, whether single or branched, helps determine these stages (Kandel *et al.*, 2020). Solar tracking in *H. annuus* is a noteworthy example of diurnal rhythm in plants. During the day, the shoot apex continuously reorients to follow the sun's position, causing the developing heads to track from East to West. Conversely, at night, the heads return to face east in anticipation of dawn. This cyclic movement diminishes and eventually ceases at anthesis, after which the sunflower head maintains an easterly orientation (Joshua *et al.*, 2014).

Sunflowers require fertile soils, with organic and inorganic sources of nitrogen, phosphorus, and potassium as soil amendments being beneficial. Granular fertilizers, ideal in spring for gradual nutrient release, or an organic approach with manure and other chemical options, can be used. Mokgolo *et al.* (2019) reported that application of poultry manure improved sunflower growth, grain yield, dry matter, and head dry matter in the second cropping season of their study. Animal manure is a valuable source of crop nutrient and organic matter which can improve soil biophysical conditions for sustainable crop production (Baiyeri and Tenkouano, 2007). Sustainable crop production systems, however, often combine the use of organic nutrient inputs with mineral fertilizers for optimum crop performance (Unagwu *et al.*, 2013; Nnadi *et al.*, 2019; Obalum *et al.*, 2020; Olawuyi *et al.*, 2022). This study was needed due to limited data on sunflower production in southeastern Nigeria and aimed to increase farmers' interest in this vital oil seed crop. Therefore, the study sought to evaluate the growth and seed yield of sunflowers (*H. annuus* L.) to co-application of organic and mineral fertilizer in Nsukka, agro-ecological zone of Nigeria.

MATERIALS AND METHODS

The experiment was conducted between September and December 2022 at the Department of Crop Science Teaching & Research Farm, University of Nigeria, Nsukka, which experiences distinct wet (April - October) and dry seasons (November -

March). Sunflower seeds were sourced from the seed collection of the Department of Crop Science, University of Nigeria, Nsukka, NPK 20:10:10 fertilizer was purchased from Ogige Market Nsukka, and decayed poultry manure sourced from the Department of Animal Science Farm, University of Nigeria Nsukka. The land area measuring 2,400 cm × 2,200 cm was cleared and ploughed. The seedbed was done manually which was partitioned into three blocks, each divided into 16 plots, each plots measuring 300 cm × 100 cm. The plots were separated from each other by 50 cm.

Soil characterization of the experimental site show that the soil is a sandy clay loam (70% sand, 7% silt and 23% clay), with a low pH in the range of 4.40-4.90, total nitrogen content of 0.15-0.42%, available phosphorus of 7.2-8.8 mg kg⁻¹, and exchangeable potassium of 0.12-0.25 cmol kg⁻¹ (Ogbonna and Ogbonna, 2010; Aba *et al.*, 2021). Also, the soil shows a low organic matter content in the range of 1.10-2.00%, indicating low fertility status. The sunflower seeds were directly sown into the seedbed by drilling method at a depth of 2.54 cm with a spacing of 50 cm × 50 cm in each plot.

The experimental design was a factorial in randomized complete block design (RCBD) with three replications. Treatments were four rates each of NPK 20:10:10 fertilizer (0, 100, 200, 300 kg ha⁻¹) and poultry manure (0, 5, 10, 15 t ha⁻¹). Data were collected on growth parameters viz plant height (from the soil level to the tip of the plant), stem girth (2 cm from soil level), and number of leaves plant⁻¹, as well as on days to 50% and 100% flowering, head diameter, head weight, and seed weight.

Data were subjected to analysis of variance following the procedure appropriate for factorial experiments in RCBD using GenStat 10.4 analytical software. The Fisher's least significant difference (F-LSD) test at a 5% alpha level, according to Obi (2006), was employed to separate treatment means for cases of significant difference.

Table 1: Monthly rainfall, temperature and humidity during 2022 collected from meteorological station at the experimental site

Months	Rainfall (mm)	Temperature (°C)		Relative humidity (%)	
		Min.	Max.	Morning	Evening
Jan	9.40	22.0	27.39	33.74	25.87
Feb	0.00	24.21	28.68	41.43	26.93
Mar	15.5	25.0	31.55	55.68	38.58
Apr	15.32	23.73	28.93	56.07	43.8
May	276.15	21.84	29.77	59.29	48
Jun	181.86	21.07	28.93	59.2	49.8
Jul	281.88	21.65	28.61	63.35	50.32
Aug	129.94	21.45	27.29	64.26	56
Sep	247.38	22.9	26.93	57.8	51
Oct	252.27	22.13	29.71	62.04	52.39
Nov	0.51	23.6	30.63	54.2	42.87
Dec	0.00	18.90	33.65	43.03	27.68
Total	1410.21	-	-	-	-

RESULTS

The weather record in the year of the experiment is shown (Table 1). Total annual rainfall was 1410.21 mm. There was an increase in rainfall from 247.38 mm in September to 252.27 mm in October, 2022. A total drop in rainfall was observed in November and December. Maximum temperature as recorded shows an increase from 26.93 °C in the month of September to 33.65 °C in the month of December. There were fluctuations in minimum temperature being 22.9, 22.13, 23.6 and 18.90 °C in September, October, November and December, respectively. Highest relative humidity was recorded in October.

Effect of Poultry Manure on the Growth and Seed Yield of Sunflower (*Helianthus annuus* L)

The data in Table 2 show that plant height increased with an increase in poultry manure rates within and across weeks 1, 3, 5 and 9 after fertilizer application with the tallest plants (122.1 cm) recorded at 9 weeks after the application of 15 t ha⁻¹ poultry manure. Stem girth initially decreased at 5 t ha⁻¹, followed by increases at 10 t ha⁻¹ and 15 t ha⁻¹ rates. The maximum stem girth (6.9 cm) occurred in week 5 with 15 t ha⁻¹ application. A decrease in the number of leaves was observed at 5 t ha⁻¹ during week one. Subsequently, there were fluctuations in the number of leaves across the following weeks. The highest leaf count of 23.5 was recorded 5 weeks after application of 10 t ha⁻¹ poultry manure. Differences were observed at 3 weeks after fertilizer application (WAF) between 15 and 10 t ha⁻¹ and between 5 and 0 t ha⁻¹.

Table 3 shows that days to 50% flowering decreased with increasing poultry manure rates.

Days to 100% flowering were lowest in the control (48.8), with the highest at 10 t ha⁻¹ (49.8). Head diameter values ranged from 12.2 cm (5 t ha⁻¹) to 16.7 cm (15 t ha⁻¹), with control at 13.6 cm. Poultry manure positively influenced head diameter. Both head weight and seed weight increased with poultry manure rate, reaching 157.6 and 55.4 g plant⁻¹, respectively at 15 t ha⁻¹. Notably, however, the differences between the application rates 15 and 10 t ha⁻¹ were nominal (Table 3).

Effect of NPK 20:10:10 on the Growth and Seed Yield of Sunflower (*Helianthus annuus* L)

Table 4 presents the effect of NPK 20:10:10 fertilizer rates on various growth parameters of sunflower. Plant height increased with fertilizer rate, reaching a peak of 129.6 cm at 300 kg NPK 20:10:10 in week 9. Stem girth progressively increased across weeks but decreased at week 9. The lowest stem girth (1.3 cm) was observed at week 1 with 0 kg NPK 20:10:10, and significant increases were noted with higher fertilizer rates. The number of leaves initially increased, peaking at week 5 with 300 kg NPK 20:10:10.

Table 5 shows that days to 50% and 100% flowering decreased with increasing fertilizer rates. Head diameter, head weight, and seed weight also increased with fertilizer rates, with the highest head weight (225.2 g plant⁻¹) recorded at 300 kg NPK 20:10:10. There were differences between fertilizer rates for such parameters as plant height, stem girth, number of leaves, days to flowering, head diameter, head weight, and seed weight (*p* > 0.05). Overall, the results highlighted the positive effects of NPK 20:10:10 fertilizer on sunflower growth and yield.

Table 2: Main effect of poultry manure rates on the growth parameters of sunflower (*Helianthus annuus*) measured at 1, 3, 5 and 9 weeks after fertilizer application

PM rates (t ha ⁻¹)	Plant height (cm)				Stem girth (cm)				Number of leaves			
	Weeks after fertilizer application				Weeks after fertilizer application				Weeks after fertilizer application			
	1	3	5	9	1	3	5	9	1	3	5	9
0	12.0	29.3	64.6	99.4	1.8	3.1	5.9	5.3	10.2	20.0	20.9	19.4
5	13.3	32.0	71.8	100.7	1.7	3.0	6.0	5.1	10.1	18.7	20.0	17.1
10	15.8	39.1	80.6	115.0	2.0	3.9	6.5	5.5	13.0	21.3	23.5	23.4
15	16.5	47.1	92.3	122.1	2.2	4.0	6.9	6.3	12.3	20.7	21.4	21.3
F-LSD (<i>p</i> ≤ 0.05)	2.6	10.2	15.4	16.3	NS	0.7	NS	NS	2.4	NS	NS	NS

Table 3: Main effect of poultry manure rates on the days to 50% and 100% flowering and on head diameter, head weight and seed weight of sunflower (*Helianthus annuus*)

Poultry manure rates (t ha ⁻¹)	DTF 50%	DTF 100%	HD (cm)	HWt (g plant ⁻¹)	SWt (g plant ⁻¹)
0	46.7	48.8	13.6	93.1	33.5
5	46.6	49.6	12.2	104.9	37.5
10	46.1	49.8	14.3	116	45.6
15	45.8	49.3	16.7	157.6	55.4
F-LSD (<i>p</i> ≤ 0.05)	NS	NS	NS	NS	13.8

DTF - Days to Flowering, HD - Head Diameter, HWt - Head Weight, SWt - Seed Weight

Table 4: Main effect of NPK 20:10:10 rates on the growth parameters of sunflower (*Helianthus annuus*) measured at 1, 3, 5 and 9 weeks after fertilizer application

NPK rates (kg ha ⁻¹)	Plant height (cm)				Stem girth (cm)				Number of leaves			
	Weeks after fertilizer application				Weeks after fertilizer application				Weeks after fertilizer application			
	1	3	5	9	1	3	5	9	1	3	5	9
0	10.7	30.7	50.9	95.6	1.3	2.4	5.1	5.0	8.7	15.9	18.3	18.3
100	13.7	30.7	68.2	95.5	1.7	3.1	5.7	5.0	10.8	18.6	20.9	19.2
200	16.0	39.2	85.4	116.5	2.0	3.8	6.8	5.7	12.8	21.7	21.7	22.8
300	17.2	47.0	104.9	129.6	2.2	4.7	7.8	6.5	13.5	24.6	24.9	21.0
F-LSD (<i>p</i> ≤ 0.05)	2.6	10.2	15.4	16.3	0.5	0.7	0.8	0.9	2.4	2.9	2.6	NS

Table 5: Main effect of NPK 20:10:10 rates on the days to 50% (DTF 50%) and 100% (DTF 100%) flowering and on head diameter (HD), head weight (HWt) and seed weight (SWt) of sunflower

NPK 20:10:10 rates (kg ha ⁻¹)	DTF		HD (cm)	HWt (g plant ⁻¹)	SWt (g plant ⁻¹)
	50%	100%			
0	49.9	52.7	9.3	39.2	16.2
100	46.8	50.3	12.2	81.5	32.2
200	44.8	48.4	14.3	125.7	45.9
300	43.6	46.2	16.7	225.2	77.6
F-LSD ($p \leq 0.05$)	2.6	2.1	1.8	48.1	13.8

Interaction Effect of Poultry Manure and NPK 20:10:10 on Sunflower Growth and Seed Yield

Table 6 shows that stem girth and number of leaves progressively increased during weeks 1, 3, and 5 but decreased in week 9. Table 7 shows that days to 50% flowering varied across the interactions, with the highest value (52.0) observed at 15 t 0-kg⁻¹ and the lowest (40.0) at 15 t 100-kg⁻¹. Days to 100% flowering decreased with increasing interaction rates. Head diameter, head weight, and seed weight exhibited an upward trend with increasing interaction rates. The highest head diameter (20.0 cm), head weight (327.7 g), and seed weight (109.8 g)

occurred at 15 t 300-kg⁻¹, while the lowest values are recorded at 0 t 0-kg⁻¹ (head diameter, 9.0 cm; head weight, 35.2 g; seed weight, 11.0 g).

DISCUSSION

The annual rainfall of 1410.21 mm during the study period slightly exceeded the FAO (2010) reported range (1000 mm) for sunflowers, indicating adaptability to regions with higher rainfall. The prevailing temperatures fell within the crop's tolerance range, aligning with Schneiter and Miller (1981), who stated that temperatures between 8 °C and 34 °C pose no significant yield reduction.

The positive effect of poultry manure on sunflower growth was evident in increased plant height, stem girth, and leaf number. This aligns with the reports of Mokgolo *et al.* (2019), affirming that poultry manure enhances plant height of sunflower at various growth stages. The slow release of nutrients from poultry manure, as observed in non-significant effects on stem girth across some weeks, resonates with findings by Kihanda *et al.* (2006), Shahzad *et al.* (2015), and

Table 6: Interaction effect of poultry manure rates and NPK 20:10:10 rates on the growth parameters of sunflower (*Helianthus annuus*) measured at 1, 3, 5 and 9 weeks after fertilizer application

Poultry manure rates (t ha ⁻¹)	NPK rates (kg ha ⁻¹)	Plant height (cm)				Stem girth (cm)				Number of leaves			
		Weeks after fertilizer application				Weeks after fertilizer application				Weeks after fertilizer application			
		1	3	5	9	1	3	5	9	1	3	5	9
0	0	10.0	24.0	45.8	87.5	1.4	2.2	5.0	4.8	7.3	15.8	19.5	15.7
	100	11.5	29.3	59.2	108.2	1.8	3.3	6.0	5.3	9.8	17.2	20.0	21.2
	200	12.8	30.3	64.5	103.8	1.8	3.5	6.5	5.8	11.0	23.5	21.5	20.5
	300	13.8	33.5	88.8	98.0	2.3	3.5	6.4	5.2	12.5	23.7	22.5	18.3
5	0	11.7	26.2	50.9	73.1	1.3	2.0	4.8	3.7	8.3	14.8	14.3	13.2
	100	14.2	30.3	62.8	84.0	1.8	2.9	5.5	4.5	10.2	19.0	21.2	16.7
	200	12.8	32.0	77.7	109.5	1.6	2.8	6.2	5.3	10.0	18.0	20.2	17.0
	300	14.7	39.5	95.8	136.2	2.0	4.5	7.8	6.8	12.0	22.8	24.5	21.7
10	0	11.0	26.5	47.0	102.7	1.3	2.8	5.5	5.2	11.2	18.5	21.3	18.8
	100	14.2	33.2	70.3	95.3	1.5	3.5	5.7	5.1	11.7	20.2	23.0	18.5
	200	19.0	48.0	98.3	126.5	2.6	4.5	7.3	5.6	16.0	23.3	23.5	33.0
	300	18.8	48.8	106.8	135.3	2.4	4.7	7.7	6.2	13.3	23.3	26.3	23.3
15	0	10.0	46.0	59.8	119.0	1.3	2.5	5.3	6.2	8.0	14.3	18.0	23.5
	100	15.0	29.8	80.3	94.3	1.5	2.8	5.8	4.8	11.3	18.0	19.5	20.5
	200	19.7	46.3	101.0	126.0	2.7	4.3	7.0	6.2	14.0	21.8	21.7	20.7
	300	21.3	66.2	128.2	149.0	3.1	6.3	9.3	8.0	16.0	28.5	26.3	20.5
<i>Grand mean</i>		<i>14.4</i>	<i>36.9</i>	<i>77.3</i>	<i>109.3</i>	<i>1.9</i>	<i>3.5</i>	<i>6.4</i>	<i>5.5</i>	<i>11.4</i>	<i>20.18</i>	<i>21.5</i>	<i>20.3</i>
F-LSD ($p \leq 0.05$)		NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS	NS

Table 7: Interaction effect of poultry manure and NPK 20:10:10 rates on the days to 50% (DTF 50%) and 100% (DTF 100%) flowering and on head diameter (HD), head weight (HWt) and seed weight (SWt) of sunflower

Manure rates (t ha ⁻¹)	NPK rates (kg ha ⁻¹)	DTF 50%	DTF 100%	HD (cm)	HWt (g plant ⁻¹)	SWt (g plant ⁻¹)
0	0	49.7	52.7	9.0	35.2	11.0
	100	47.7	50.7	12.3	36.2	22.0
	200	48.0	48.7	13.4	119.8	37.2
	300	41.3	43.3	15.8	181.2	63.8
5	0	49.7	51.7	8.0	27.8	12.2
	100	46.3	50.3	12.3	78.3	29.5
	200	44.7	48.7	14.5	109.1	40.8
	300	45.7	47.7	15.8	204.5	67.3
10	0	48.3	51.7	9.9	49.0	21.5
	100	47.0	51.3	11.5	81.3	32.0
	200	43.7	48.0	15.5	146.3	59.3
	300	45.3	48.3	15.3	187.5	69.7
15	0	52.0	54.7	10.2	45.0	20.0
	100	40.0	49.0	12.5	130.2	45.2
	200	43.0	48.3	13.6	127.4	46.4
	300	42.0	45.3	20.0	327.7	109.8
<i>Grand mean</i>		<i>46.3</i>	<i>49.4</i>	<i>13.1</i>	<i>117.9</i>	<i>43.0</i>
F-LSD ($p \leq 0.05$)		NS	NS	NS	NS	NS

Mahmood *et al.* (2017). Application of organic fertilizer, specifically poultry manure, not only boosts soil fertility but also aids in nutrient recycling and potentially mitigating environmental pollution. This corresponds to the observations by Abumere *et al.* (2019), emphasizing the role of organic fertilizers in enhancing soil microbes, nutrient availability, and uptake by sunflower plants.

Days to 50% and 100% flowering decreased with increasing poultry manure rates, aligning with Agu *et al.* (2015). The non-significant effect of poultry manure on head diameter and weight may be attributed to the gradual release of nutrients (Abumere *et al.*, 2019). The significant effect of poultry manure on seed weight implies that poultry manure enhances soil structure and soil fertility, thereby promoting enhanced nutrient uptake and leading to improved plant health and higher crop yields (Oshundiya *et al.*, 2014; Ogunzezi *et al.*, 2019). This highlights the role of organic fertilizer in increasing sunflower seed yield and attributes. The application rates 15 and 10 t ha⁻¹ produced similar seed yield; however, the former outyielded while the latter did not outyield the un-amended control. This observation makes the maximum application rate in this study to be the minimum rate for growing sunflower in the study area.

The significant effect of NPK 20:10:10 fertilizer on multiple facets of sunflower growth and yield, such as plant height, stem girth, number of leaves, days to flowering, head diameter, head weight, and seed weight can be attributed to the concentrated and balanced nutrient composition of NPK 20:10:10, minimizing nutrient wastage and ensuring efficient nutrient supply at different growth stages. This finding aligns with the work of Wahyu and Donald (2019), who observed the influence of NPK fertilizer on sunflower growth and seed yield. Among the four application rates of NPK 20:10:10, 300 kg ha⁻¹ gave the highest values for plant height, stem girth, number of leaves, head diameter, head weight, and seed weight. Additionally, this application rate gave the lowest days to 50% and 100% flowering. This suggests that a balanced supply of nutrients, as provided by the 300 kg ha⁻¹ rate, optimally supports sunflower growth, leading to increased plant growth and accelerated flowering. The positive relationship between nitrogen application and seed yield attributes is consistent with findings from Bakht *et al.* (2010), Ahmed *et al.* (2011), and Nasim *et al.* (2012). This study emphasizes the role of nitrogen in enhancing sunflower seed yield. The application of NPK 20:10:10, with its nitrogen content, likely contributes to improved seed yield by promoting favourable conditions for sunflower development.

The study shows positive growth and yield responses of sunflower to both organic and inorganic fertilizers, indicating an efficient conversion of applied nutrients for improved yield. The interaction effect between poultry manure and NPK 20:10:10 fertilizer further demonstrates the enhancement of growth and yield parameters in sunflower fields.

Abumere *et al.* (2019) observed that inorganic fertilizer facilitated accumulation of dry matter compared to organic fertilizer. This is attributed to the faster release of nutrients from mineral fertilizers, accelerating dry matter accumulation. The increased accumulation of dry matter in sunflower shoots, particularly in plots with higher rates of both organic and inorganic fertilizers, underscores the importance of nutrient availability for economic yield. This aligns with the findings of Paul *et al.* (2017), who highlighted the advantages of integrating organic and inorganic nutrient sources over the use of inorganic fertilizer alone.

Although the study found non-significant effects of the interaction between poultry manure and NPK 20:10:10 for all parameters, their combination at 15 t ha⁻¹ and 300 kg ha⁻¹, respectively outyielded the un-amended control by about 10 times. This is large compared to the corresponding number of times by which poultry manure at 15 t ha⁻¹ (4.8) and NPK 20:10:10 at 300 kg ha⁻¹ (1.7) outyielded the control in the main effects. Similar to these results, Esilaba *et al.* (2005) and Ogbonna (2008) reported non-significant effects of the interaction between manure and fertilizer in eggplant, but found their combination to produce the best yield.

CONCLUSION & RECOMMENDATION

Sunflower is not a common crop in the study area but it can thrive with appropriate cultural practices. The study has shown that though poultry manure and NPK 20:10:10 interaction consistently showed non-significant effects, their complementary use at 15 t ha⁻¹ and 300 kg ha⁻¹, respectively gave the highest performance. Therefore, if poultry manure and NPK 20:10:10 must be combined in sunflower production, they should be applied at these stated rates, otherwise our data support adoption of sole application of either poultry manure at 15 t ha⁻¹ or NPK 20:10:10 at 300 kg ha⁻¹ for optimum growth/productivity of sunflower in the derived savannah.

REFERENCES

- Aba S.C., Baiyeri K.P. and Ortiz R. (2021). Soil fertility dynamics and yield responses of four plantain genotypes to different nutrient management systems in Nsukka, southeastern Nigeria. *Fruits, The Int. J. Trop. Subtrop. Hort.*, **76** (3), 134-144. <https://doi.org/10.17660/th2021/76.3.4>
- Abumere V.I., Dada O.A., Adebayo A.G., Kutu F.R. and Togun A.O. (2019). Different rates of chicken manure and NPK 15-15-15 enhanced performance of sunflower (*Helianthus annuus L.*) on ferruginous soil. *Int. J. Agron.*, Vol. 2019, Article ID 3580562, 10 pp. <https://doi.org/10.1155/2019/3580562>
- Agu R.S., Ezema R.A., Udegbunam O.N. and Okoro A.C. (2015). Effect of different rates of poultry manure on growth and yield of cucumber (*Cucumis sativum*) in Iwollo, Southeastern Nigeria. *Agro-Sci.*, **14** (3), 41-44
- Ahmed M.E., Abdel-Gawad A.M., Shehata E.I., Tawfik S.A. (2011). Influence of using reed forage in different forms as fresh, silage and hay on blood profile and carcass quality of growing Rahmani sheep. *Egyptian J. Sheep Goat Sci.*, **6** (2), 25-35

- Ali B., Ridzwan A.H., Alireza N., Izham A. and Farzad A. (2014). Influence of plant spacing and sowing time on yield of sunflower (*Helianthus annulus* L.). *J. Food Agric. Environ.*, **12** (2), 688-691
- Bakht J., Sahfi M., Yousaf M., Raziuddin and Khan M.A. (2010). Effect of irrigation on physiology and yield of sunflower hybrids. *Pak. J. Bot.*, **42** (2), 1317-1326
- Balogh L. (2008). Sunflower species (*Helianthus* spp.) In: Botta-Dukát Z. and Balogh L. (eds.), *The Most Important Invasive Plants in Hungary* (pp. 227-255)
- Baiyeri K.P. and Tenkouano A. (2007). Manure placement influence growth and dry matter yield of a plantain hybrid. Presented at the 8th Afr. Crop Sci. Conf., Elminia, Egypt. *Afr. Crop Sci. Conf. Proc.*, **8**, 385-390
- Dorling K. (2008). *RHS A-Z Encyclopedia of Garden Plants* (p. 1136). United Kingdom
- Esilaba A.O., Byalebek J.B., Delve R.J., et al. (2005). Effect of organic and inorganic sources of fertilizer on growth, yield and fruit quality of eggplant (*Solanum melongena* L.). *Agric. Syst.*, **86**, 144-165
- FAO (2010). The State of Food Insecurity in the World: Addressing Food Insecurity in Retracted Crises. Food & Agriculture Organisation (FAO), Rome, Italy
- Joshua P.V., Robin I.B., Stacey L.H. and Benjamin K.B. (2014). Turning heads: the biology of solar tracking in sunflower. *Plant Sci.*, **224**, 20-26
- Kaleem S., Hassan F., Ahmad M., et al. (2009). Effect of growing degree days on autumn planted sunflower. *Afr. J. Biotechnol.*, **10**, 8840-8846. <https://doi.org/10.5897/AJB11.608>
- Kandel H., Endres G. and Buetow R. (2020). *Sunflower Production Guide*. North Dakota Agricultural Experimental guide and North Dakota State University Extension, A1995
- Kihanda F.M., Warren G.P. and Micheni A.N. (2006). Effect of manure application on crop yield and soil chemical properties in a long-term field trial of semi-arid Kenya. *Nutr. Cycl. Agroecosyst.*, **76** (2-3), 341-354
- Mahmood F., Khan I., Ashraf U., et al. (2017) Effects of organic and inorganic manures on maize and their residual impact on soil physico-chemical properties. *J. Soil Sci. Plant Nutr.*, **17** (1), 22-32
- Mokgolo M.J., Mzezewa J. and Odhiambo J.J.O. (2019). Poultry and cattle manure effects on sunflower performance, grain yield and selected soil properties in Limpopo province, South Africa. *S. Afr. J. Sci.*, **115** (11/12), Article No. 6410, 7 pp. <https://doi.org/10.17159/sajs.2019/6410>
- Nasim W., Ahmad A., Khaliq T., et al. (2012). Effect of organic and inorganic fertilizer on maize hybrids under agro-environmental conditions of Faisalabad-Pakistan. *Afr. J. Agr. Res.*, **7** (17), 2713-2719
- Nnadi A.L., Nnanna P.I., Onyia V.N., Obalum S.E. and Igwe C.A. (2020). Growth and yield responses of high-density coverage sweet potato to liming and fertilizer combinations for sandy-loam Ultisols at Nsukka, southeastern Nigeria. In: *Climate-Smart Soil Management, Soil Health/Quality and Land Management: Synergies for Sustainable Ecosystem Services* (pp. 263-269), Proc. 44th Annual Conf. Soil Sci. Soc. Nigeria, 16-20 March 2020 [Colloquia SSSN 44], Enugu State University of Science & Technology, Enugu, Enugu State, Nigeria
- Obalum S.E., Ogumba P.O. and Uzoh I.M. (2020). Influence of tillage-seedbed and manure-NPK-micronutrient management options on selected soil properties of sandy-loam Ultisols evaluated using sweet potato. *Nigerian J. Soil Sci.*, **30** (3), 117-125
- Obi I.U. (2006). *Statistical Methods of Detecting Differences Between Treatment Means and Research Methodology Issues in Laboratory and Field Experiments* (p. 117). AP Express Enugu, Nigeria
- Ogbonna P.E. (2008) Effect of combined application of organic and inorganic fertilizers on fruit yield of egg plant (*Solanum melongena*). *Proc. 42nd Annual Conf. Agric. Soc. Nigeria (ASN)*, pp. 236-250
- Ogbonna P.E. and Ogbonna C.S. (2010). Fruit yield responses of Eggplant (*Solanum melongena*) to single and combined applications of organic and inorganic fertilizers. *Bio-Research*, **8** (1), 633-637
- Ogunzei K.C., Okebalama C.B. and Obalum S.E. (2019). Optimum poultry droppings rate for coarse-loamy Ultisols based on soil macro-aggregation and fertility indices and evaluation using cucumber (*Cucumis sativus*). Presented under *Understanding Nigerian Soils for Sustainable Food & Nutrition Security and Healthy Environment*, the 43rd Annual Conference of Soil Sci. Soc. Nigeria (SSSN), 15-19 July 2019, University of Agriculture, Makurdi, Benue State, Nigeria
- Olawuyi O.J., Ezeanyi C.U. and Orkpeh U. (2022). Morphological characterization and response of red flower rag leaf (*Crassocephalum crepidioides* Benth S. Moore) to organic and inorganic fertilizers and arbuscular mycorrhizal fungus. *Agro-Sci.*, **21** (1), 51-60
- Oshundiya F.O., Olowe V.I.O., Sowemimo F.A. and Odedina J.N. (2014). Seed yield and quality of Sunflower (*Helianthus annuus* L.) as influenced by staggered sowing and organic fertilizer application in the humid. *Helia*, **37** (61), 237-255. <https://doi.org/10.1515/helia-2014-0012>
- Paul O.J., Frank O.O. and Ambrose A.E. (2017). Effect of poultry manure and NPK fertilizer on growth and yield of eggplant. *J. Agric. Rural Res.*, **1** (2), 112-117
- Pilorgé E. (2020). Sunflower in the global vegetable oil system: Situation, specificities and perspectives. *Oilseeds Fats Crops Lipids (OCL)*, **27**, 34 <https://doi.org/10.1051/ocl/2020028>
- Putt E.D. (1997). Early history of Sunflower, vol 35. Retrieved from <https://access.onlinelibrary.wiley.com/doi/abs/10.2134/agronmonogr35.c1>
- Schilling and Edward E. (2006). *Helianthus*. In: Flora of North America editorial committee (eds.). Vol 21
- Schneider A. and Miller J.F. (1981). Description of sunflower growth stages. *Crop Sci.*, **21**, 901-903. <https://doi.org/10.2135/cropsci1981.0011183X002100060024x>
- Shahzad T., Chenu C., Genet P., et al. (2015). Contribution of exudates, arbuscular mycorrhizal fungi and litter depositions to the rhizosphere priming effect induced by grassland species. *Soil Biol. Biochem.*, **80**, 146-155
- Sommerlad J. (2022). What is the national flower of Ukraine? <https://www.independent.co.uk/news/world/europe/ukraine>
- Steel R.G.D. and Torrie J.H. (1980). *Principles and Procedures of Statistics: A Biometrical Approach* (2nd edn.). McGraw-Hill Book Company, New York. <https://doi.org/10.2307/2287561>
- Unagwu B.O., Asadu C.L.A. and Obalum S.E. (2013). Maize performance in a sandy loam Ultisol amended with NPK 15-15-15 and poultry manure. In: *The Role of Crop Science in the Agricultural Transformation Agenda* (pp. 135-141), Proc. 1st National Ann. Conf. Crop Sci. Soc. Nigeria (CSSN), 15-19 Sep. 2013, University of Nigeria, Nsukka, Nigeria
- Vasyl C. (2023). Growing sunflowers: High cultivation practices. *EOS Data analysis*. <https://eos.com/blog/how-to-grow-sunflowers/>
- Wahyu H. and Donald S. (2019). Study of NPK fertilizer effect on sunflower growth and yield. *AIP Conf. Proc.* **2120**, 030031. <https://doi.org/10.1063/1.5115635>