



**Bayero Journal of Pure and Applied Sciences, 15(2): 15 - 20**

Received: 11<sup>th</sup> June, 2022

Accepted: 25<sup>th</sup> Sept., 2022

ISSN 2006 – 6996

## **INTEGRATION OF RNAi CONSTRUCT INTO THE GENOME OF COWPEA (*Vigna unguiculata* L.) FOR RESISTANCE TO *Cowpea Aphid Borne Mosaic Virus* (CAMV) AND *Cowpea Severe Mosaic Virus* (CPSMC)**

**Sani<sup>1\*</sup> L. A., Usman<sup>2</sup> I. S., Abdulmalik<sup>2</sup> M. M. and Aragao<sup>3</sup> F. J. L.**

1. Department of Plant Biology, Bayero University Kano, Nigeria

2. Department of Plant Science, Ahmadu Bello University, Zaria, Nigeria

3. EMBRAPA- CENARGEN, Brasilia DF, Brazil

\*Corresponding author: Email: labdul.bot@buk.edu.ng

### **ABSTRACT**

**Improvement of cowpea for resistance to major pests and viral diseases is impeded by the absence of resistance genes in its genome. Genetic engineering is a suitable tool through which genes of interest could be introduced into the genome of cowpea. In order to generate cowpea plant resistant to CAMV and CPSMC, RNA interference construct that target the silencing of CAMV coat protein and CPSMC proteinase cofactor genes were successfully inserted and integrated into cowpea lines using the biolistic gene delivery system. Three transgenic cowpea lines were regenerated from bombardment of 600 embryos with a transformation frequency of 0.5%. Polymerase Chain Reaction (PCR) analysis of all the primary transformed plants (T<sub>0</sub>). T<sub>1</sub> progeny showed the presence of 340bp fragment amplified by primers COMOPTYF389 within the CPSMV sequence and COMOPOTYR729 within the CABMV sequence indicating successful genes insertion and integration of the interfering cassette.**

**Keywords: RNA interference, Cowpea, Virus resistance, Biolistic mediated transformation**

### **INTRODUCTION**

Cowpea (*Vigna unguiculata* L. Walp) is an important source of dietary protein in sub-Saharan Africa for thousands of years. The dry grains contain 23-32% protein and other important food components such as carbohydrate, lipid, fibre, minerals and vitamins (Nielson *et al.*, 1993). About 7.56 million metric tons of cowpea is produced annually over an estimated area of 12.76 million hectares with over 70% of the dry grain production coming from sub-Saharan Africa ([www.iita.org/cms/detail/cowpea\\_project\\_details](http://www.iita.org/cms/detail/cowpea_project_details)). In sub-Saharan Africa, insect pests and viral diseases are critical yield limiting factors which reduce the overall grain yield of cowpea to 0.495t/ha, which is per below the crop's potential estimated at 3.0t/ha (FAO, 2012). Improvement of cowpea for resistance to major pests and viral diseases using conventional breeding is impeded by the absence of resistance genes in the genome of cultivated cowpea varieties. Genetic engineering is a suitable tool through which genes of interest could be introduced into the genome of cowpea. The technology could help breeders overcome major

production challenges in cowpea, especially when applied along with conventional breeding. The *cowpea aphid-borne mosaic virus* (CABMV) belongs to the family *Potyviridae* and genus *potyvirus*. The virus is transmitted by seed and several aphid species which belongs to subfamily *Aphidinae*. CABMV infects cowpea resulting in yield reductions and frequent total crop failure in sub-Saharan countries like Nigeria (Thottappilly and Rossel, 1992). The genome of CABMV is composed of a positive-sense single stranded RNA (ssRNA) molecule with a genome-linked protein (VPg) at the 5' end and a 3' terminal poly(A) tail that contains a single ORF encoding a large precursor polyprotein (Cruz and Aragao, 2014). This polyprotein is processed into P1, helper component (HC), P3, cylindrical inclusion (CI), nuclear inclusion A (NIA), nuclear inclusion B (NIB) and capsid protein (CP), and two small putative proteins known as 6K1 and 6K2 (Chung *et al.*, 2008). The cowpea severe mosaic virus (CPSMV) belongs to the family *Secoviridae* and genus *comovirus*. CPSMV is the most important virus infecting cowpea in America and is among the seven most damaging viruses attacking cowpea in Africa.

It is transmitted either by leaf feeding beetles belonging to the family Chrysomelidae or by seed. The genome of CPSMV consists of two positive-sense single stranded RNA (ssRNA) molecules; RNA1 and 2, which separately encapsulated and expressed by a polyprotein processing strategy (Cruz and Aragao, 2014). RNA1 encodes proteins necessary for replication; the 32K proteinase cofactor, the 58K helicase, the VPg 5'-linked proteins of the genomic RNAs expected set of five mature proteins, 24K proteinase and 87K polymerase, while RNA2 encodes the two capsid proteins and a protein involved in cell-to-cell movement (Chen and Bruening, 1992).


RNA interference (RNAi) is a mechanism that regulates gene expression by activating a sequence-specific RNA degradation process. RNAi has become a valuable biotechnological tool, because double-stranded (ds) RNA generated from the expression of chimerical constructs encoding self-complementary hairpin (hp) RNA can selectively and robustly induce suppression of specific genes of interest (Cruz and Aragao, 2014). This strategy has been used to generate transgenic plants with systemic resistance to viruses confer by sequence specific degradation of the viral RNA (Lopez *et al.*, 2010; Vanderschuren *et al.*, 2012; Aragao, 2014; Cruz and Aragao, 2014; Ammara *et al.*, 2015). This paper reports successful integration of dsRNA

for resistance to CABMV and CPSMV into genome of cowpea.

## MATERIALS AND METHODS

### The Plasmid Vector

A 1063bp interfering cassette containing sequence which corresponds to 502bp fragment from the Coat Protein sequence of *Cowpea Aphid Borne Mosaic Virus* (CABMV) (Gene Bank accession number- KF181277), and 561bp from the Proteinase Cofactor gene (32K protein) of *Cowpea Severe Mosaic Virus* (CPSMV) RNA1 (Gene Bank accession number-M83830), was synthesized at Epoch life Science Inc. USA. Sites for *XbaI*, *SacI*, *ApaI* and *KpnI* were added to allow cloning in sense and antisense orientations in the vector pSIU (Tinoco *et al.*, 2010), generating the vector pSIUCAUPI2. The 1063bp sequence interspersed with intron 3 (370bp) obtained from the malate synthase gene (*ms-i3*) of *Arabidopsis thaliana*. The cassette was removed with enzymes *EcoRI* and *BglII* from the pSIUCAUPI2 vector and cloned into site for *EcoRI* and *BglII* in pBSK vector (Gene Bank accession no.GS54365-1). The pBSK contains the gene *ahas* (which confers tolerance to the herbicide imazapyr) from *Arabidopsis thaliana* and the gene *bla* which confers resistance to ampicillin. The plasmid vector GS54365-1pBSK containing the CABMV/CSMV hairpin sequence was generated and referred to as COMOPOTY.




**Figure 1:** The map of the plasmid vector GS54365-1pBSK-COMOPOTY used to transform the standard cowpea variety, containing the *ahas* gene (*ahas5'*: *ahas* gene promoter; *ahas*: *Arabidopsis thaliana* AHAS coding sequence; *ahas3'*: *ahas* gene terminator), the  $\Delta$ CABMVCSMV cassette (AMV: untranslated leader sequence from *Alfalfa mosaic virus*; intron AtMS: intron 3 from the malate synthase gene from *A. thaliana*). The *bla* gene codes for a  $\beta$ -lactamase that confers resistance to ampicillin.

### Cowpea transformation

The cowpea line (MNC05-828-3-15) was transformed with the plasmid vector GS54365-1pBSK-COMOPOTY according to Cruz and Aragao, (2008). Mature seeds were washed under running tap water and surface sterilized by dipping in 70% ethanol for 1 minute and subsequent shaking in 2% sodium hypochlorite (w/v) for 20 minutes. The seeds were rinsed three times in sterile distilled water then soaked for 16–18 h. Embryonic axes were excised from the sterilized seeds and the apical meristems were exposed by removing both the primary and primordial leaves under microscope. The embryonic axes were then placed with the apical region directed upward in bombardment plates containing basal Murashige and Skoog (MS) medium supplemented with 30g/L sucrose and 8g/L agar. The bombardment was conducted using a high-pressure, helium-driven particle acceleration biolistic machine based on the

procedure described by Aragao *et al.* (2005) and Rech *et al.* (2008). After the bombardment, embryonic axes were immediately transferred to culture bottles containing 80ml of cowpea selection media containing MS medium fortified with 22.2M BAP and 300nM imazapyr. Elongated shoots developed from the bombarded apical meristems after 3 weeks of culture on the selection medium. Regenerated shoots were allowed to reach 4-5 cm in length before they were transferred to hormone free media, without selection pressure for rooting. The transgenic plant lines ( $T_0$  generation) were transferred to plastic cups containing soil:vermiculite (1:1) and covered with transparent polyethylene bags for two weeks for acclimatization. After two weeks of hardening in a small plastic cups, plants were kept for one week before they were transferred to large plastic pots and allowed to grow to maturity.



**Figure 2:** Regeneration of transgenic plant lines and detection of 340bp fragment from COMOPOTY using PCR analyses. (a) Transgenic plant line regenerated on selection media. (b) Acclimatized transgenic seedling (c) Matured and fertile transgenic plant in greenhouse (d) Polymerase chain reaction (PCR) amplification of 340bp fragment (1, 2 and 3), B and C- represent three primary transformed plants, blank (water) and negative control (untransformed cowpea plant).

### Detection of the Transgenes in Primary Transformed ( $T_0$ ) Plants


After one week of hardening, leaf discs were collected from the  $T_0$  and  $T_1$  lines for detection of the interfering cassette. DNA was isolated from leaf discs according to Edwards *et al.*, (1991). PCR was carried out according Bonfim *et al.*, (2007). The primers COMOPTYF389 within the CPSMV sequence and COMOPOTYR729 within the CABMV sequence were used to amplify a 340bp fragment from the interfering cassette- $\Delta$ CABMVCSMV.

### RESULTS

In order to generate transgenic cowpea lines resistant to *Cowpea Aphid Borne Mosaic Virus* (CABMV) and *Cowpea Severe Mosaic Virus* (CPSMV), the plasmid vector GS54365-1pBSK-COMOPOTY was used to deliver the chimeric fragment ( $\Delta$ CSMVCABMV) containing a sequence from the Coat Protein sequence of CABMV and Proteinase Cofactor gene (32K protein) of CPSMV- RNA1 (Figure 1). The results showed that siRNA interfering cassette was successfully delivered into genome of cowpea following bombardment of the apical regions of embryonic axes with the plasmid GS54365-1pBSK-COMOPOTY.

A total of three primary transgenic plant lines (T<sub>0</sub>) harbouring the chimeric fragment containing dsRNA of Coat Protein gene of CABMV and Proteinase Cofactor gene of CPSMV were regenerated from 10 transformation events each containing 60 embryos which corresponds to the transformation frequency of 0.5%. After 3 weeks in culture under selection in the presence of 300 nM imazapyr, shoot buds were observed. Microscopic examination of the apical region showed that numerous shoot buds were induced, but only about 1.0% of shoot buds elongated (Figure 2a). All the primary transformed plants (T<sub>0</sub>) presented normal phenotype (Figure 2c). PCR analysis of genomic DNA isolated from the primary transformed plants (T<sub>0</sub>) showed a successful insertion of the RNAi cassettes ( $\Delta$ CSMVCABMV). This is confirmed by the presence of 340bp fragment amplified by primers COMOPTYF389 within the

CPSMV sequence and COMOPOTYR729 within the CABMV sequence (Figure 2d). The primary transformed plants (T<sub>0</sub>) transferred the transgene to T<sub>1</sub> progeny. When T<sub>1</sub> plants were cultivated in the greenhouse, the plants presented morphologically normal growth and were fertile (Figure 2c). The plants presented normal plant height, site of development of first pod, number of branches, internode length, foliar area, flowers, pods and seeds per pod. The PCR analysis of the T<sub>1</sub> progeny also showed the presence of 340bp fragment amplified by primers COMOPTYF389 within the CPSMV sequence and COMOPOTYR729 within the CABMV sequence indicating successful transfer of the interfering cassette (Figure 3). The results showed that the siRNA cassette inserted in the primary transformed (T<sub>0</sub>) plant has been successfully integrated into the genome of cowpea and transmitted to the T<sub>1</sub> generation.



**Plate VII:** PCR amplification of the 340bp fragment amplifies by the primer (COMOPTYF389 and COMOPOTYR729) from the T<sub>1</sub> progeny. Three T<sub>1</sub> transgenic lines (1, 2 and 3) obtained from the first primary transformed (T<sub>0</sub>) plant, B and C- represent blank and negative control (untransformed cowpea plant).

## **DISCUSSION**

The diseases caused by CABMV and CPSMV constitute important constraints to the production of cowpea especially in America and Africa (Aliyu *et al.*, 2012; Abreu *et al.*, 2012). In this work, a reproducible system was developed in which integration of dsRNA chimeric construct comprising a fragment of the coat protein gene of CABMV and a fragment from the proteinase cofactor gene of CPSMV ( $\Delta$ CSMVCABMV) is routinely achieved. The system is based on the bombardment of pre-exposed apical meristem of cowpea embryonic axes followed by selection of transgenic line using herbicide molecule – Imazapyr. The herbicide molecule belongs to the imidazolinone class (imazapyr), which is capable of systemically translocating and concentrating in the apical meristematic region of the plant (Ivo *et al.*, 2008). Earlier reports (Citadin *et al.*, 2013; Cruz and Aragao, 2014) have shown that this system could be used to introduce genes of agronomic importance into genome of cowpea.

The system target multiple shoots induction in the presence of BAP as means of achieving significant levels of regeneration of transgenic plants. A relatively high transformation frequency (0.5%) was achieved compared to 0.05 - 0.15% reported by Popelk *et al.*, (2006) and Chaudhury *et al.*, (2007) using *Agrobacterium*-mediated transformation system. On the other hand, transformation frequency of 1.41% was reported by Ivo *et al.* (2008), Cruz and Aragao (2014) using a similar approach. In addition to relatively higher transformation frequency, this system is much simpler because it utilizes embryonic axes and is devoid of multiple cultivation steps employed in the *Agrobacterium* mediated transformation of cowpea.

A strategy involving the use of RNAi technology to generate genetically modified plants resistant to RNA viruses have earlier been reported in cowpea (Cruz and Aragao, 2014), sweet potato (Kreuze *et al.*, 2008) and Mexican lime (Lopez *et al.* 2010).

Transgenic plants expressed considerable resistance following mechanical inoculation with the viruses and resistance was found to correspond with the presence of transgene-derived siRNA in the leaves of all transgenic lines. Earlier reports (Cruz and Aragao, 2014) indicated that the presence of the silencing construct in the transgenic cowpea lines induced the silencing of the viral genes by reducing or preventing viral RNA accumulation through sequence specific degradation of the viral RNA

## REFERENCES

- Abreu E. F.M ., Tinoco M. L. P. , Andrade E. C., and Aragão F. J. L., (2012). Diversity among isolates of cowpea severe mosaic virus infecting cowpeas in northeastern Brazil. *Genet. Mol. Res.* 11 (3): 3146-3153. DOI <http://dx.doi.org/10.4238>
- Aliyu, T.H., Balogun, O.S. and Kumar, L., (2012). Survey of the Symptoms and Viruses Associated with Cowpea (*Vigna unguiculata* (L.) in the Agroecological Zones of Kwara State, Nigeria. *Ethiopian Journal of Environmental Studies and Management* Vol.5(4) 613-619.
- Ammara U., Mansoor S., Saeed M., Amin I., Briddon R. B. and Al-Sadi A. M., (2015). RNA interference-based resistance in transgenic tomato plants against *Tomato yellow leaf curl virus-Oman* (TYLCV-OM) and its associated betasatellite. *Virology Journal* 12 (38) 1-12. DOI: [10.1186/s12985-015-0263-y](https://doi.org/10.1186/s12985-015-0263-y)
- Aragao F. J. L., Vianna G. R., Carneiro S. B. R. C. & Rech E. L., (2005) Germ line genetic transformation in cotton (*Gossypium hirsutum* L.) by selection of transgenic meristematic cells with a herbicide molecule. *Plant Science*; 168:1227–1233. DOI:[10.1016/j.plantsci.2004.12.024](https://doi.org/10.1016/j.plantsci.2004.12.024).
- Aragao, F. J. L., (2014). GM plants with RNAi - golden mosaic resistant bean, *BioMed Central Proceedings*; 2014, 8(Suppl. 4):O24.
- Bonfim, K., Faria, J. C., Nogueira, E. O. P. L., Mendes, E. A. & Aragão, F. J. L., (2007): RNAi-Mediated Resistance to Bean golden mosaic virus in Genetically Engineered Common Bean (*Phaseolus vulgaris*). *Molecular Plant-Microbe Interaction*; 20(6):717-726.
- Chaudhury, D., Madanpotra, S., Jaiwal, R., Saini, R., Kumar, A. P. & Jaiwal, P. K., (2007). *Agrobacterium tumefaciens* mediated high frequency genetic transformation of an Indian cowpea (*Vigna unguiculata* L. Walp.) cultivar and transmission of transgenes into progeny. *Plant Science*; 172:692-700. DOI: [10.1016/j.plantsci.2006.11.009](https://doi.org/10.1016/j.plantsci.2006.11.009).
- Chen X, Bruening G, 1992. Cloned DNA copies of cowpea severe mosaic virus genomic RNAs: infectious transcripts and complete nucleotide sequence of RNA 1. *Virology* 191, 607–18. *Plant Pathology* (2014) 63, 831–837
- Chung BY, Miller WA, Atkins JF, Firth AE, 2008. An overlapping essential gene in the Potyviridae. *Proceedings of the National Academy of Sciences, USA* 105, 5897–902.
- Citadin, C. T., Cruz A. R. R. & Aragao F. J. L., (2013). Development of transgenic imazapyr tolerant cowpea (*Vigna unguiculata*). *Plant Cell Report*; 32:537–543, DOI:[10.1007/s00299-013-1385-6](https://doi.org/10.1007/s00299-013-1385-6)
- Cruz, A. R. R. and Aragao, F. J. L., (2014). RNAi-based enhanced resistance to *Cowpea severe mosaic virus* and *Cowpea aphid-borne mosaic virus* in transgenic cowpea. *Plant Pathology* 63, 831–837, DOI: [10.1111/ppa.12178](https://doi.org/10.1111/ppa.12178).
- Edwards, K., Johnstone C. & Thompson C., (1991). A simple and rapid method for the preparation of plant genomic DNA for PCR analysis. *Nucleic Acids Research* 19:1349. doi:[10.1093/nar/19.6.1349](https://doi.org/10.1093/nar/19.6.1349)
- Food and Agriculture Organization, (2012). The second report on the state of the world's plant genetic resources for food and agriculture - <http://www.fao.org>.
- Ivo, N. L., Nascimento, C. P., Vieira, L. S., Campos, F. A. P. & Aragão, F. J. L. (2008). Biolistic mediated genetic transformation of cowpea (*Vigna unguiculata*) and stable Mendelian inheritance of transgenes. *Plant Cell Reports*; 27:1475-83; PMID:[18587583](https://pubmed.ncbi.nlm.nih.gov/18587583/). DOI:[10.1007/s00299-008-0573-2](https://doi.org/10.1007/s00299-008-0573-2).
- Kreuze, J. F., Klein, I. S., Lazaro, M. U., et al., (2008). RNA silencing mediated resistance to crinivirus (Closteroviridae) in cultivated sweet potato (*Ipomoea batatas* L.) and development of sweet potato virus

- disease following co-infection with potyvirus. *Molecular Plant Pathology*; (9):589–98.
- Lopez, C., Cervera, M., Fagoaga, C., *et al.*, (2010). Accumulation of transgene-derived siRNAs is not sufficient for RNAi-mediated protection against Citrus tristeza virus in transgenic Mexican lime. *Molecular Plant Pathology*; 11: 33–41.
- Nielson SS, Brandt WE, Singh BB (1993). Genetic variability for nutritional composition and cooking time of improved cowpea lines. *Crop Sci.* 1993; 33: 469-472
- Popelka, J. C., Gollasch, S., Moore, A., Molvig, L. & Higgins, T. J. V., (2006). Genetic transformation of cowpea (*Vigna unguiculata* L.) and stable transmission of the transgenes to progeny. *Plant Cell Reports*; 25:304-12; PMID:16244884, DOI:10.1007/s00299-005-0053-x.
- Rech, E. L. F., Vianna, G. R. & Aragao F. J. L., (2008). High-efficiency transformation by biolistic of soybean, common bean and cotton transgenic plants. *Nature Protocols*; 3: 410–8.
- Thottappilly, G. and Rossel, H.W., (1992). Virus diseases of cowpea in Tropical Africa. *Tropical Pest Management*; 38(4):337 – 348.
- Tinoco MLP, Dias BBA, Dall’Qastta RC, Pamphile JA, Aragao FJL, 2010. In vivo trans-specific gene silencing in fungal cells by in plant expression of a double-stranded RNA. *BMC Biology* 8, 1–27.
- Vanderschuren, H., Moreno, I., Anjanappa, R. B., *et al.* (2012). Exploiting the combination of natural and genetically engineered resistance to cassava mosaic and cassava brown streak viruses impacting cassava production in Africa. *PLoS One*; 7: e45277.