

## VOLATILE PROFILE OF GREEN COFFEE BEANS FROM *COFFEA ARABICA* L. PLANTS GROWN AT DIFFERENT ALTITUDES IN ETHIOPIA

Girmay Tsegay<sup>1,2</sup>, Mesfin Redi-Abshiro<sup>1</sup>, Bhagwan Singh Chandravanshi<sup>1\*</sup>, Estifanos Ele<sup>1</sup>,  
Ahmed M. Mohammed<sup>1</sup> and Hassen Mamo<sup>3</sup>

<sup>1</sup>Department of Chemistry, College of Natural Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

<sup>2</sup>Agricultural Quality Research Laboratory, Ethiopian Institution of Agricultural Research, P.O. Box 2003, Addis Ababa, Ethiopia

<sup>3</sup>Department of Microbial, Cellular and Molecular Biology, College of Natural Sciences, Addis Ababa University, P.O. Box 1176, Addis Ababa, Ethiopia

(Received March 18, 2019; Revised October 16, 2019; Accepted October 26, 2019)

**ABSTRACT.** This study was aimed to identify volatile compounds of 31 green coffee bean samples and evaluate their correlation with altitude of the coffee plants grown in two zones (Gedeo and Jimma) in Ethiopia. A total of 81 different compounds were detected. The contents of dominant volatile compounds in green coffee beans were in the range: trans-linalooloxide (3.24–19.13%), linalool (1.56–21.76%), 2-methoxy-4-vinylphenol (2.34–15.08%) and cis-linalooloxide (1.03–13.27%). In addition, benzene acetaldehyde (0.45–10.97%), 2-heptanol (0.67–8.49%),  $\alpha$ -terpineol (0.64–6.52%), phenylethyl alcohol (0.44–4.98%) and furfural (0.92–5.3%) were the next dominant compounds. The volatile compounds identified in the green coffee beans are groups of alcohols, aldehydes, ketones, pyrazines, pyridines, and furans. The volatile compounds in green coffee beans showed either weak positive or weak negative correlation with the altitude of coffee plants indicating that variation in altitude of the coffee plants does not significantly influence the volatile compounds of green coffee beans.

**KEY WORDS:** *Coffea arabica* L., Coffee plants, Green coffee beans, Volatile compounds, Effect of altitudes, Heterocyclic compounds

### INTRODUCTION

The name of coffee is derived from the name of the province Keffa (birth place of coffee) where shepherds from Abyssinia/Ethiopia discovered it in the world [1, 2]. Coffee beans are the seeds of a shrub in the genus *Coffea* and belonging to the family Rubiaceae. It has over sixty different species of which only three (*Coffea arabica*, *Coffea robusta (canephora)*, and *Coffea liberica*) have globally commercial values [3]. *Coffea arabica* and *Coffea canephora* are the most important commercial species [4, 5]. Arabica coffee has a higher quality than Robusta coffee according its complex aroma and flavor [6, 7]. Coffee is the most valuable commodities in the world. It is the second largest traded commodity after oil in the world [8] and it is an important non-alcoholic beverage crop [9].

Coffee quality is defined by its sensorial aspects [10]. Quality and content of the chemical composition of coffee beans vary based on the species of coffee, altitude, soil, temperature variations and the place where it grows [7, 8, 11-14]. Recently some studies have been reported on the effect of altitude of the coffee plants on the chemical composition of green coffee beans. Sherge *et al.* [15] have studied the influence of growing altitude, shade and harvest period on quality and biochemical composition of Ethiopian specialty coffee. Hagos *et al.* [3] have studied the correlation between caffeine contents in green coffee beans and altitudes of the coffee plants grown in southwest Ethiopia. Mintesnot and Dechassa [16] have studied the effect of altitude, shade, and processing methods on the quality and biochemical composition (caffeine, trigonelline, and chlorogenic acids) of green coffee beans in Ethiopia. Adem *et al.* [17] have

\*Corresponding author. [bscv2006@yahoo.com](mailto:bscv2006@yahoo.com)

This work is licensed under the Creative Commons Attribution 4.0 International License

reported the effect of altitude on biochemical composition (caffeine, chlorogenic acids and sucrose contents) and quality of green arabica coffee beans. Bodner *et al.* [18] have reported the effect of harvesting altitude on the aroma released by coffee powder. Gebrekidan *et al.* [19] have reported the influence of altitudes of coffee plants on the alkaloids contents of Ethiopian green coffee beans. Tolessa *et al.* [20] have reported the influence of growing altitude, shade and harvest period on quality and biochemical composition of Ethiopian specialty coffee and concluded that coffee from high altitude with open or medium shade and early to middle harvest periods had a superior bean quality and favored the production of beans with lower caffeine. Worku *et al.* [21] have studied the effect of altitude on biochemical composition (the caffeine, chlorogenic acids and sucrose contents) and physical and sensory qualities of green Arabica coffee beans from large plantations in south western Ethiopia.

Coffee taste is different due to the presence of different volatile and nonvolatile chemical constituents. Volatile compounds are fragrant components, generally odorous, which occur in certain indigenous plants or at specified parts of the plants. Their presence in green coffee beans is responsible for the aroma profile [14, 22, 23]. Their profile could be used to confirm the authenticity of *Coffea arabica*. Volatile compounds of coffee beans are diverse in their chemical characteristics. Numerous volatile chemicals have been identified in the green coffee beans [24].

The roasting process mainly induces the majority of volatile chemicals in coffee. Many studies reported the volatile chemical constituents of the brewed coffee. However, there are only a few reports on the identification and quantification of volatile components in the green coffee beans [25]. Toci and Farah [26] identified about 100 volatile compounds in the green coffee beans. Lee and Shibamoto [27] have reported the major constituents of volatile compounds found in green coffee beans. Bertrand *et al.* [28] investigated volatile compounds of green coffee beans. Dong *et al.* [29] identified volatile compounds and bioactive components of seven coffee (*Coffea robusta*) cultivars grown in Hainan Province, China. Toledo *et al.* [30] have comprehensively reviewed the relationship between the different aspects related to coffee quality and their volatile compounds. This study has provided an overview of different aspects related to the quality of coffee beans and their volatile fractions: species/cultivars, geographic origins, bean defects, and types of beverages, processing, roasting, and storage.

The most abundant classes of volatile compounds are alcohols, esters, hydrocarbons, and aldehydes. Heterocyclic compounds are very important components in providing coffee with their characteristic flavors. Some of these compounds have been found useful in differentiating between Robusta and Arabica coffee varieties [31]. Some volatile compounds found in green coffee bean have been indicated as characteristics of bean defects due to certain external causes like over fermentation during processing or insect attacks [32]. However, the vast majority of volatiles compounds are only generated during the roasting of green coffee beans [33].

Characterization of volatile compounds of coffee was used to identify coffee origin and production locations [34]. Risticvic *et al.* [34] have reported profile of volatile flavor chemicals of three roasted Arabica coffee samples obtained from three different countries (Ethiopia, Nicaragua, and Indonesia). The Ethiopian Yirgacheffe coffee produced the highest level of 5-hydroxymethylfurfural (26.8%) in light roasting, furfuryl alcohol (28.2%) in medium roasting and  $\gamma$ -butyrolactone (13.3%) in heavy roasting conditions. In another study, Akiyama *et al.* [35] have reported volatile aroma compounds of Arabica coffee extracts from three production countries (Ethiopia, Tanzania, and Guatemala). Their report showed that the aroma profile of Ethiopian Sidama coffee was discriminately different from Tanzanian and Guatemalan coffees. They further reported that 4-(4-hydroxyphenyl)-2-butanone can characterize the aroma profile of the freshly brewed Ethiopian Sidama coffee. Quantitative analysis also indicated that the lightly roasted Ethiopian coffee contains a higher amount of this compound.

A foregoing review revealed that identification and quantification of volatile compounds in the green coffee beans from different origins including Ethiopia have been reported [15, 16, 18, 20, 23, 24, 26, 28, 29]. However, a comprehensive investigation exclusively devoted to the

characterization of volatile profiles from the various coffee cultivars of Ethiopia is very limited. Therefore, it is necessary to study systematically the volatile compounds of green coffee beans obtained from coffee plants grown at different altitudes in Ethiopia. The objectives of this study were to (i) identify and quantify the volatile compounds of green coffee bean samples from the coffee plants grown at different altitudes in Ethiopia by gas chromatography-mass spectrometry and (ii) evaluate their correlation with altitude of the coffee plants.

## EXPERIMENTAL

### *Chemicals*

Dichloromethane ( $\geq 97\%$ , Sigma-Aldrich, USA) GC/MS grade was used for the extraction of volatile compounds after hydro-distillation of green coffee beans. Anhydrous sodium sulfate (Fluka, Buchs, Switzerland) was used for drying the dichloromethane extract.

### *Sample collection*

Thirty-one green coffee bean samples were collected from Gedeo Zone, Yirgacheffe District (21) and Jimma Zone, Gomma District (10), Ethiopia, during December 2017 to March 2018 from the farmers who produce a special coffee under the supervision of agricultural experts for the export to international markets. The description of green coffee bean samples and sampling sites (sample number, sampling zone, sampling district/woreda and altitude) are given in Table 1. All the green coffee bean samples were obtained from the ripen coffee cherries and processed by sun drying. About 1 kg of green coffee beans was collected from different farmers from each sampling site and stored in paper bags under room temperature conditions. It was ground in an electronic grinder (Moulinex, SEB Group, Selongey, France) for the analysis.

### *Extraction of volatile compounds by hydro-distillation*

The volatile compounds were extracted from green coffee beans by hydro-distillation according to the procedure of Saw *et al.* [24] with some modifications. Each ground sample of green coffee beans (200 g) was transferred into 2 L round bottom flask with 1 L of distilled water. The mixtures were distilled for 6 h at atmospheric pressure. The distillate was transferred to 50 mL of separatory funnel, and 3 mL of dichloromethane was added and shaken properly. The lower organic phase was separated and dried over anhydrous  $\text{Na}_2\text{SO}_4$  to remove residual water content, and the dried extract was transferred into a 2.5 mL glass vial for gas chromatography-mass spectrometry (GC-MS) analysis.

### *Analysis of volatile compounds*

The GC-MS analysis was carried out by using Agilent technologies 7820A GC and 5977E MSD system equipped with auto-sampler. Chromatographic separations were carried out using DB1701 capillary column with 30 m length, 0.25 mm internal diameter and 0.25  $\mu\text{m}$  column phase thickness. Injection mode was split-less, helium was the carrier gas and 1  $\mu\text{L}$  volume of the sample was injected at a constant flow rate of 1 mL/min to the inlet heated to 275  $^{\circ}\text{C}$ . Initial oven temperature was 60  $^{\circ}\text{C}$  with 2 min hold time then heated to 200  $^{\circ}\text{C}$  with ramp of 10  $^{\circ}\text{C}/\text{min}$  and 3  $^{\circ}\text{C}/\text{min}$  to 240  $^{\circ}\text{C}$ . Conditions used for the mass spectrometer were a temperature of 230  $^{\circ}\text{C}$  for the ion source and 150  $^{\circ}\text{C}$  for quadruple, scan range 40-650 m/z, operating in positive electron impact mode with ionization energy of 70 eV. Chromatographic and mass spectral data were processed by using the instrument built in software (MS Mass Hunter; Agilent Technologies, USA). The volatile components were identified from chromatograph library

(NIST-14) and by the MS fragmentation pattern of authentic chemicals. The method validation for the analysis by GC-MS was done by running the blank and standard methyl ester of decanoic acid before the analysis of samples.

Table 1. The description of green coffee bean samples and sampling sites (sample number, sampling zone, sampling district/woreda and altitude).

Sample number	Sampling zone	District/woreda	Altitude (m)
J32	Jimma	Gomma	1515
J31	Jimma	Gomma	1579
J24	Jimma	Gomma	1627
J30	Jimma	Gomma	1727
J25	Jimma	Gomma	1737
J29	Jimma	Gomma	1782
G12	Gedeo	Yirgacheffe	1840
G11	Gedeo	Yirgacheffe	1855
G13	Gedeo	Yirgacheffe	1860
G10	Gedeo	Yirgacheffe	1878
G03	Gedeo	Yirgacheffe	1891
G04	Gedeo	Yirgacheffe	1904
G09	Gedeo	Yirgacheffe	1916
G22	Gedeo	Yirgacheffe	1918
G02	Gedeo	Yirgacheffe	1922
G21	Gedeo	Yirgacheffe	1945
G01	Gedeo	Yirgacheffe	1971
J33	Jimma	Gomma	1972
J28	Jimma	Gomma	2010
G16	Gedeo	Yirgacheffe	2037
G20	Gedeo	Yirgacheffe	2044
G19	Gedeo	Yirgacheffe	2046
G15	Gedeo	Yirgacheffe	2050
G17	Gedeo	Yirgacheffe	2050
G18	Gedeo	Yirgacheffe	2063
G14	Gedeo	Yirgacheffe	2068
J26	Jimma	Gomma	2137
G06	Gedeo	Yirgacheffe	2139
J27	Jimma	Gomma	2163
G05	Gedeo	Yirgacheffe	2200
G07	Gedeo	Yirgacheffe	2220

## RESULTS AND DISCUSSION

### *Identification and quantification of volatile compounds of green coffee beans*

Volatile components isolated from green coffee beans (*Coffea arabica* L.) were identified using the comparison of the corresponding mass spectra in GC-MS with data of NIST-14 mass spectral data library found in GC-MS database. Their quantification was carried out according to their peak area of the chromatogram.

Figure 1 shows chromatogram peak profile of volatile compounds presents in green coffee beans detected by GC-MS. Most of the compounds are found in the range of 4.2 min up to 28.5 min in all the samples, but the compounds detected after 25.5 min were not identified. The chromatogram shows that the most intense compounds are trans-linalooloxide at a retention time

of about 8.2 min, cis-linalooloxide (8.56 min), linalool (8.75 min), 2-methoxy-4-vinylphenol (12.77 min).




Figure 1. The chromatogram of volatile compounds of green coffee bean (sample number G04) from Gedeo zone.

In this study, 81 volatile compounds with 18 chemical classes were detected as shown in Figure 1. The peak number, name of volatile compounds, their chemical class, retention time (RT) and range of percentage peak area of the identified compounds in the 31 green coffee bean samples by GC-MS are given in Table 2. Some compounds in green coffee beans reported in the literature by Saw *et al.* [24] and Bertrand *et al.* [28] (such as  $\alpha$ -terpineol, trans-linalool oxide, cis-linalool oxide, 2-heptanol, furfural, linalool, benzyl alcohol, indole, methyl stearate) were also identified in the green coffee bean samples of the present study. The components detected in roasted coffee beans like furans, pyrazines, and pyridines were also identified in the present study. Among these, the dominant compounds (mean/range) are: trans-linalool oxide (12.2%, 3.24–19.13%), linalool (9.28%, 1.56–21.76%), 2-methoxy-4-vinylphenol (8.47%, 2.34–15.08%), cis-linalooloxide (7.31%, 1.03–13.27%), with the retention time of 8.20, 9.28, 12.8 and 8.47 min, respectively. Benzene acetaldehyde (5.18%, 0.45–10.97%), 2-heptanol (4.86%, 0.67–8.49%),  $\alpha$ -terpineol (4.85%, 0.64–6.52%), phenylethyl alcohol (4.07%, 0.44–4.98%) and furfural (3.15%, 0.92–5.3%) are the next dominant compounds, respectively.

Table 2. The peak number, name of volatile compounds, their chemical class, retention time (RT) and range of percentage peak area of the identified compounds in the 31 green coffee bean samples by GC-MS.

Peak No.	Compound name	Class	RT	Range of peak area (%)
01	3-Methyl-2-butenal	Aldehyde	4.28	0.13–4.35
02	Ethyl-3-methyl, butanoate	Ester	4.80	0.63–3.23
03	2,6-Lutidine	Pyridine	5.01	0.16–0.94
04	3,5-Dimethylpyrazole	Pyrazole	5.28	0.53–3.69
05	Pyridine, 2,3-dimethyl-	Pyridine	5.30	0.61–1.37
06	Furfural	Furan	5.32	0.92–5.3
07	3-Furfural aldehyde	Furan	5.36	0.62–4.5
08	1H-Imidazole,1,4-dimethyl-	Imidazole	5.38	0.35–1.30
09	2-Heptanol	Alcohol	5.63	0.67–8.49

10	1,4-Hexadiene, 4-methyl	Hydrocarbon	6.36	0.13–2.5
11	Eucalyptol/1,3,3-trimethyl-2-oxabicyclo[2.2.2]octane	Ether	6.94	0.34–5.27
12	Benzaldehyde	Aldehyde	7.08	0.65–3.95
13	1-Butanamine, 4[2,4-bis(1,1-dimethylpropyl)phenoxy]-N-propylidene	Amine	7.32	0.36–2.48
14	N-Ethylidene t-butylamine	Amine	7.36	0.34–1.85
15	1-Pyrolidineethaneamine/1-(2-aminoethyl)pyrrolidine	Pyrolidine	7.41	0.30–3.21
16	Cyclohexanol, 2,2-dimethyl-	Alcohol	7.72	0.25–1.77
17	3-Ethyl-4-methylpentan-1-ol	Alcohol	7.79	0.22–1.9
18	2-Ethyl piperidine	Piperidine	7.81	0.27–1.98
19	1-Pyrolid-2-one, N-carbamoyl-	Pyrolidine	7.84	0.21–1.90
20	1,6-Dioxaspiro[4.4]nonane, 2-ethyl	Hydrocarbon	7.96	0.20–1.64
21	Pyrazole, 3,5-dimethyl-1-allyl-	Pyrazole	8.12	0.16–5.01
22	1H-1,2,3-Triazole-5-carboxamide, 4-amino-	Triazole	8.01	0.20–4.22
23	Trans-linalooloxide	Furan	8.20	3.24–19.13
24	Cis-linalooloxide	Furan	8.49	1.03–13.27
25	Benzeneacetaldehyde	Aldehyde	8.60	0.45–10.97
26	Linalool	Alcohol	8.75	1.56–21.76
27	Benzyl alcohol	Alcohol	8.94	0.38–2.47
28	1,3-Dioxolane-2-propanal, 2-methyl	Aldehyde	9.16	0.17–0.67
29	2-Cyclohexene-1-ol, 1-methyl-4-[1-methylethenyl]-, trans-	Alcohol	9.57	0.23–1.39
30	Pyrazine, 3,5-diethyl-2-methyl-	Pyrazine	9.23	0.11–1.56
31	6-Pethyl-3,5-heptadiene-2-one	Ketone	9.44	0.53–1.25
32	1,3,8- <i>p</i> -Menthatriene	Hydrocarbon	9.55	0.12–6.24
33	Pyrazine, 2-methoxy-3-[2-methylpropyl]-	Pyrazine	9.40	0.09–1.68
34	Benzen, 1,2-dimethoxy	Ether	9.86	0.32–4.73
35	Phenylethyl alcohol	Alcohol	10.0	0.44–4.98
36	2-Butanone, 4-[5-methyl-furanyl]	Furan	10.0	0.08–4.60
37	$\alpha$ -Terpineol	Alcohol	10.3	0.64–6.52
38	Methyl salicylate	Ester	10.4	0.81–4.98
39	S-Benzoyl(thiohydroxylamine)	Amine	10.5	0.08–10.51
40	Benzene, 1-methyl-4-[1-methylethylene]-/ <i>p</i> -Isopropenyl toluene	Hydrocarbon	10.7	0.07–0.53
41	Cyclopentane, 1,3-bis[methylene]	Hydrocarbon	10.7	0.1–0.90
42	Hexane, 1-chloro-5-methyl	Hydrocarbon	10.8	0.07–1.30
43	3-Carene/3,7,7-Trimethylbicyclo[4.1.0]hept-3-ene	Hydrocarbon	10.89	0.35–1.88
44	1H-Imidazole, 2-ethyl-4-methyl-	Imidazole	10.93	0.06–4.17
45	Phenylephrine	Alcohol	10.93	0.58–0.86
46	3,4,5-Trimethylpyrazole	Pyrazole	10.94	0.18–0.97
47	(3E,5E)-2,6-Dimethylocta-3,5,7-trien-2-ol	Alcohol	11.13	0.17–1.32
48	Benzene acetic acid, ethyl ester	Ester	11.23	0.22–2.68
49	2,6-Octadien-1-ol, 3,7-dimethyl-,(z)/trans-farnesol	Alcohol	11.28	1.22–3.08
50	Benzoic acid, 2-hydroxy-, ethyl ester	Ester	11.48	0.15–2.76
51	Citral	Aldehyde	11.75	0.22–1.23
52	Benzeneacetaldehyde, .alpha.-ethylidene	Aldehyde	12.19	0.08–0.84
53	2,4-Decadienal, (E,E)-	Aldehyde	12.41	0.19–0.54
54	2-Methoxy-4-vinylphenol	Alcohol	12.77	2.34–15.08
55	3-Methoxyacetophenone	Ketone	12.00	0.16–7.91

56	Phenol, <i>p</i> -tert-butyl-	Alcohol	12.88	0.46–1.12
57	2-Butene-1-one, 1-[2,6,6-trimethyl-1,3-cyclohexadien-1-yl]-, (E)-	Ketone	13.03	0.16–1.53
58	Indole	Indole	13.52	0.45–0.94
59	3-Amino-1,2,4-triazole-5-carboxylic acid	Triazole	13.85	0.20–0.50
60	2[3H]-Furanone, dihydro-5-pentyl-	Furan	14.04	0.12–0.50
61	Acetamide, N-phenyl-	Amide	15.47	0.15–0.25
62	$\alpha$ -Isomethylionone	Kitone	14.21	0.21–0.7
63	Lilial	Aldehyde	14.96	0.09–1.88
64	2,4-Di-tert-butylphenol	Alcohol	15.09	0.16–0.45
65	Octadecane	Hydrocarbon	16.36	0.08–0.79
66	Pentadecanal	Aldehyde	16.57	0.15–4.09
67	Hexadecanal	Aldehyde	16.57	0.15–3.01
68	Octanal, 2-{phenylmethylene}-	Aldehyde	17.65	0.84–1.74
69	2-Pentadecanone, 6,10,14-trimethyl	Kitone	18.22	0.36–1.16
70	Methylpent-4-enylamine	Amine	18.24	0.24–0.91
71	2,7-Diphenylindole	Indole	18.69	0.14–1.52
72	2-Butene-1,4-diamine, N,N'-dimethyl-	Amine	18.93	1.21–2.69
73	Hexadecanoic acid, methyl ester	Ester	19.93	1.04–12.71
74	Caffeine	Amide	21.36	0.50–1.60
75	7-Octadecenoic acid, methyl ester	Ester	21.98	0.13–3.5
76	9,12-Octadecadienoic acid, methyl ester	Ester	22.08	0.26–7.73
77	Methyl stearate	Ester	22.24	0.24–6.46
78	<i>p</i> -Hydroxynorephedrine	Ephedrine	23.15	0.18–0.72
79	Linoleic acid, methyl ester	Ester	23.25	0.20–3.51
80	<i>o</i> -Veratramide	Amide	23.26	0.26–3.6
81	Stigmasterol	Alcohol	25.99	0.58–2.73

According to Lee and Shibamoto [27] and Bertrand *et al.* [28] about 74 compounds were identified in green coffee beans from different countries. Among the major components of volatile compounds found in green coffee beans were 3-methyl butanoic acid, phenyl ethyl alcohol, hexanol, 4-hydroxy-3-methylacetophenone, 3-methyl butanol, 2-methoxy-3-(2-methylpropyl)-pyrazine. Brevard and Yeretizian [36] identified about 219 volatile compounds and approximately 100 different volatile compounds were identified in green coffee beans [26]. Poyraz *et al.* [37] have reported about 40 compounds in green coffee beans. In their report, the components of the green coffee beans were identified as isoamyl alcohol (10.4%), hexanal (10.4%) and hexacosane (8.2%) while furfuryl alcohol (13.6%), furfurylacetate (10.7%) and 5-methyl furfural (9.27%) were identified as the main components of the roasted coffee obtained from a local market in Turkey in 2014. While the major compounds identified in present study are: trans-linalool oxide (12.2%), linalool (9.28%), 2-methoxy-4-vinylphenol (8.47%), cis-linalooloxide (7.31%), benzene acetaldehyde (5.18%), 2-heptanol (4.86%),  $\alpha$ -terpineol (4.85%), phenylethyl alcohol (4.07%) and furfural (3.15%) which are different from the compounds reported by Poyraz *et al.* [37].

The maturation stage of the coffee beans is an important factor for the volatile composition of green coffee beans [37]. Volatile compounds identified in the extracted green coffee beans of the present study are a group of alcohols, aldehydes, ketones, esters, hydrocarbons, pyrroles, furans, and other compounds as listed in Table 3, which are accountable to flavor profiles. In this study, the first dominant class of compounds in the green coffee beans is an alcohol with the average value of 31.1%. This is comparable with the results reported by Brevard and Yeretizian [36]. Green coffee beans are also very rich in aldehydes and esters. Esters are important compounds in sensory point of view. The highest average value of chemical classes of volatile compounds present in 31 green coffee bean samples are shown in Figure 2. Alcohols, furans,

and aldehydes are the most abundant classes of compounds in the Ethiopian green coffee beans of the present study. In another study, the major volatile compounds in green coffee beans were aldehydes (benzaldehyde) and alkanes [37]. Whereas the major volatile compounds in roasted beans determined were a group of heterocyclic compounds as furans, pyrazines, and pyridines [38].




Figure 2. Average percentage peak area for chemical classes of volatile compounds that constitute in 31 green coffee bean samples.

Heterocyclic compounds are responsible for the characteristic flavor of coffee as flavoring agents and these compounds are also used as stimulants and medication. Their existence in green coffee beans is somewhat different from roasted coffee beans. Over 350 heterocyclic compounds, including pyrroles, furans, pyrazines, thiazoles, oxazoles, thiophenes and imidazoles, have been identified in the roasted coffee beans. Lee and Shibamoto [27] have reported that heterocyclic compounds were not found in green coffee beans, except for 2-methoxy-3-(2-methylpropyl)-pyrazine. In the present study, furans, pyrazines, imidazole, indole, piperidine, pyrazole, pyridine, triazole and pyrrolidine, even 2-methoxy-3-(2-methylpropyl)-pyrazine were identified in the extract of green coffee beans. Their presence (or absence) in the individual green coffee bean samples are summarized in Table 3. These heterocyclic compounds are among the important contributors to the coffee flavor [39] and they are used as flavoring agents like 2-methoxy-4-vinylphenol. Many compounds responsible to spicy, earthy (pyrazine, 2-methoxy-3-[2-methylpropyl]) and buttery flavor were also identified in the present study. Nitrogen-containing heterocyclic compounds are found more in green coffee beans collected from Jimma than from Gedeo. Therefore, it can be concluded that coffee plants grown in Jimma could be distinguished from coffee plants grown in Gedeo samples according to nitrogen heterocyclic compounds (Table 3).

Pyridine (in samples J25, J26, J29, J30), triazole (in samples J25, J26, J27, J29, J30, J31, J33) and piperidine (in samples J26, J27, J29, J30, J31, J33) are only found in green coffee bean samples from Jimma. In addition, pyrazine is only found in four samples from Gedeo (G10, G12, G20, G22) but in many sample from Jimma and ephedrine is only found in sample J24 and



sample J33 (Jimma) (Table 3). Ephedrine is a medication, stimulant and it is used to prevent low blood pressure during spinal anesthesia. The volatile compounds of green coffee beans have an antioxidant activity [27]. The antioxidant activities of volatile compounds in green coffee beans are due to the presence of several components. Compounds responsible for antioxidants activities such as benzyl alcohol, methyl salicylate, and 4-hydroxy-3-methylacetophenone are identified in the research of Lee and Shibamoto [27] and these compounds are also identified in the present study.

Table 3. Total peak area (percentage) of each chemical class in individual green coffee bean samples.

Sample number	Altitude (m)	Class of compounds*																	
		Alc	Ald	Am	Ami	Eph	Est	Eth	Fur	Hyd	Ime	Ind	Ket	Pip	Pyr	Prz	Pyri	Pyro	Tri
J32	1515	31.4	10.4	1.80	0.94	-	23.9	0.81	13.8	2.72	2.88	-	0.95	-	-	-	-	-	-
J31	1579	42.9	18.0	0.26	11.03	0.72	8.17	2.66	21.7	1.12	0.86	1.52	2.22	1.56	0.79	0.50	-	1.90	0.98
J24	1627	7.50	3.92	-	-	0.25	37.4	3.71	5.40	0.38	4.17	0.14	0.63	-	-	-	-	-	-
J30	1727	40.0	15.0	0.26	3.33	0.72	7.19	7.21	10.2	3.87	1.60	0.75	3.36	1.56	0.79	4.46	0.94	4.38	0.98
J25	1737	29.2	18.4	-	-	0.25	6.25	1.58	14.4	5.91	4.17	0.50	1.71	-	-	0.68	0.73	-	0.96
J29	1782	52.5	11.5	0.51	4.13	0.72	6.87	3.77	8.94	6.14	2.42	0.72	3.45	1.56	0.79	4.42	0.64	1.90	1.31
G12	1840	26.0	6.64	1.77	2.18	-	11.0	1.39	14.3	2.24	-	-	-	1.89	-	-	-	-	-
G11	1855	19.5	6.37	-	1.60	-	5.52	1.22	12.9	5.20	1.84	-	0.59	-	-	-	-	-	-
G13	1860	29.1	11.9	-	2.06	-	3.24	1.28	16.8	2.70	2.43	-	1.05	-	-	-	-	-	-
G10	1878	22.2	7.17	-	1.04	-	10.3	1.35	10.3	2.44	0.61	-	2.40	-	0.99	-	-	-	-
G03	1891	35.5	12.9	-	0.91	-	1.26	2.54	32.1	3.48	-	1.98	1.12	-	-	-	-	-	-
G04	1904	40.9	12.9	-	2.05	-	2.24	2.49	31.6	3.80	-	1.08	1.51	-	-	-	-	3.67	-
G09	1916	40.2	13.8	-	3.83	-	15.2	2.75	21.0	4.14	-	2.58	2.21	-	-	2.34	-	3.67	-
G22	1918	26.1	6.13	-	2.22	-	11.2	1.35	19.0	5.09	-	-	1.44	-	1.08	-	-	-	-
G02	1922	42.3	10.1	1.33	2.74	-	5.07	2.06	26.8	5.71	3.72	-	-	-	-	-	-	-	-
G21	1945	10.7	2.51	0.30	0.88	-	17.7	0.60	8.58	2.53	0.64	-	2.54	-	-	-	-	-	-
G01	1971	24.3	5.00	-	-	-	9.50	1.34	14.6	1.72	-	-	1.67	-	-	-	-	-	-
J33	1972	29.8	12.8	0.26	1.27	0.72	27.2	3.76	9.08	1.68	0.86	0.83	2.57	1.56	0.79	0.50	-	1.90	0.98
J28	2010	15.9	7.94	-	-	-	20.5	3.53	-	-	-	-	0.84	-	-	2.22	-	-	-
G16	2037	36.8	12.1	-	3.90	-	8.54	2.53	26.7	3.35	-	0.80	2.50	-	-	1.31	-	0.65	-
G20	2044	26.5	8.92	-	1.23	-	7.47	1.77	16.5	4.37	-	-	0.52	-	1.49	-	-	-	-
G19	2046	38.1	15.3	-	4.30	-	9.54	2.53	34.7	3.62	-	0.93	1.73	-	-	3.07	-	3.21	-
G15	2050	27.9	15.4	-	4.00	-	12.8	2.33	24.3	2.90	-	0.95	3.13	-	-	3.01	-	0.30	-
G17	2050	20.2	7.88	-	-	-	1.37	1.27	10.8	0.92	-	-	2.52	-	-	-	-	-	-
G18	2063	36.8	1.75	-	2.42	-	24.8	2.27	23.6	2.27	0.93	-	1.39	-	-	-	-	-	-
G14	2068	15.1	7.66	6.75	1.50	-	6.51	0.85	6.88	4.50	1.75	-	0.63	-	-	-	-	-	-
J26	2137	32.9	8.99	0.51	3.10	0.72	10.5	3.53	10.3	2.70	2.68	0.35	1.20	1.56	0.79	3.92	0.58	1.90	1.18
G06	2139	27.1	4.38	0.74	1.88	-	8.06	1.18	14.7	3.70	1.45	-	-	-	-	-	-	-	-
J27	2163	51.8	9.18	0.51	4.13	0.72	8.77	0.00	8.98	6.14	1.54	0.72	3.45	1.56	0.79	1.32	-	1.90	1.31
G05	2200	45.4	13.5	-	3.51	-	7.77	1.96	26.9	3.94	-	1.04	2.32	-	-	3.48	-	3.67	-
G07	2220	40.8	10.0	-	2.31	-	9.47	3.20	28.4	3.30	-	-	-	-	-	-	-	-	-
Mean		31.1	9.95	0.52	2.34	0.18	11.1	2.2	16.9	3.31	1.11	0.53	1.67	0.32	0.33	1.08	0.10	0.97	0.26

\*Note: Alc (alcohol), Ald (aldehyde), Am (amide), Ami (amine), Eph (Ephedrine), Est (ester), Eth (ether), Fur (furan), Hyd (hydrocarbon), Ime (imidazole), Ind (indole), Ket (ketone), Pip (piperidine), Pyr (pyrazine), Prz (pyrazole), Pyri (pyridine), Pyro (pyrrolidine), Tri (triazole).

#### *Correlations between the chemical classes of volatile compounds in green coffee beans with an altitude of the Ethiopian coffee plants*

The Pearson correlation shows weak, moderate or strong relationship among variables in the positive or negative direction [40]. The volatile compounds were grouped according to their chemical classes and correlations were evaluated with an altitude of Ethiopian coffee plants. The correlation coefficients of main chemical classes of alcohol, furan, and hydrocarbon with

altitude have a weak significant positive correlation ( $p = 0.01$ ) which indicates that the percentage of these compounds increases with increasing altitude. While aldehyde, amine, ester, and ether have a weak negative significant correlation ( $p = 0.01$ ) which indicates that the percentage of these compounds decreases with increasing altitude as shown in Figure 3. The correlation coefficients of all chemical classes of volatile compounds in the green coffee beans with respect to the altitudes of coffee plants are listed in Table 4. Since the two zones (Jimma and Gedeo) have variable altitudes it is not possible to differentiate the compositions of the eight classes of compounds used to evaluate the correlation coefficients.

Table 4. Correlation coefficients between the main chemical classes of volatile compounds and altitude of Ethiopian coffee plants.

Compound	Alcohol	Aldehyde	Amine	Ester	Ether	Furan	Hydrocarbon	Ketone
Correlation coefficient (R)	0.111	-0.215	-0.095	0.237	-0.181	0.195	0.124	0.068




Figure 3. Correlation between the main eight chemical classes of volatile compounds in the green coffee beans and the altitude of coffee plants grown in Ethiopia.

There is lack of relevant literature on the correlation of volatile compounds in the green coffee beans with the altitude of the coffee plants. No specific study on the correlation of altitudes and volatile compounds has been reported in the literature. To the best of our knowledge, this is the first investigation on the correlation of volatile compounds in green coffee beans and the altitudes of the coffee plants.

## CONCLUSION

A total of 81 different volatile compounds with a group of 18 chemical classes were identified in Ethiopian green coffee bean samples. The major volatile components identified in green coffee

beans are trans-linalool oxide, linalool, 2-methoxy-4-vinylphenol, cis-linaloloxide, benzene acetaldehyde, 2-heptanol, and  $\alpha$ -terpineol. About 80% of the total volatile compounds are a class of alcohols, furans, aldehydes and esters and very few heterocyclic compounds are identified in the green coffee beans. Nitrogen-containing heterocyclic compounds are found more in green coffee beans collected from Jimma than from other sites. Therefore, it can be concluded that coffee plants grown in Jimma could be distinguished from coffee plants grown in Gedeo samples according to nitrogen heterocyclic compounds. This could be due to genetics, soil composition, climate, and rainfall differences. Altitudes have a weak negative effect on the esters content of the Ethiopian green coffee beans. The quality of coffee depends on many parameters including chemical composition, soil characteristics, altitudes and climatic conditions. Therefore, it is not possible to determine the quality of coffee from the composition of volatile compounds in the green coffee beans only.

#### ACKNOWLEDGMENTS

The authors express their gratitude to the Department of Chemistry for providing laboratory facilities. Girmay Tsegay is thankful to Ethiopian Institution of Agricultural Research, Addis Ababa, Ethiopia, for sponsoring his study.

#### REFERENCES

1. Amamo, A.A. Coffee production and marketing in Ethiopia. *Eur. J. Bus. Manage.* **2014**, *6*, 109–122.
2. Weldegebreal, B.; Redi-Abshiro, M.; Chandravanshi, B.S. Development of new analytical methods for the determination of caffeine content in aqueous solution of green coffee beans. *Chem. Cent. J.* **2017**, *11*, Article 126, DOI: 10.1186/s13065-017-0356-3.
3. Hagos, M.; Redi-Abshiro, M.; Chandravanshi, B.S.; Ele, E.; Mohammed, A.M.; Mamo, H. Correlation between caffeine contents in green coffee beans and altitudes of the coffee plants grown in southwest Ethiopia using ultraviolet-visible spectrophotometry. *Bull. Chem. Soc. Ethiop.* **2018**, *32*, 13–25.
4. Yisak, H.; Redi-Abshiro, M.; Chandravanshi, B.S. New fluorescence spectroscopic method for the determination of alkaloids in green coffee beans. *Chem. Cent. J.* **2018**, *12*, Article 59, DOI: 10.1186/s13065-018-0431-4.
5. Yisak, H.; Redi-Abshiro, M.; Chandravanshi, B.S. Selective determination of caffeine and trigonilline in aqueous extract of green coffee beans by FT-MIR-ATR spectroscopy. *Vib. Spectrosc.* **2018**, *97*, 33–38.
6. Mehari, B.; Redi-Abshiro, M.; Chandravanshi, B.S.; Combrinck, S.; Atlabachew, M.; McCrindle, R. Profiling of phenolic compounds using UPLC–MS for determining the geographical origin of green coffee beans from Ethiopia. *J. Food Compos. Anal.* **2016**, *45*, 16–25.
7. Mehari, B.; Redi-Abshiro, M.; Chandravanshi, B.S.; Atlabachew, M.; Combrinck, S.; McCrindle, R. Simultaneous determination of alkaloids in green coffee beans from Ethiopia: Chemometric evaluation of geographical origin. *Food Anal. Methods* **2016**, *9*, 1627–1637.
8. Mussatto, S.I.; Machado, E.M.S.; Martins, S.; Teixeira, J.A. Production, composition, and application of coffee and its industrial residues. *Food Bioprocess. Technol.* **2011**, *4*, 661–672.
9. Kathurima, C.; Gichimu, B.M.; Kenji, G.M.; Muhoho, S.M.; Boulanger, R. Evaluation of beverage quality and green bean physical characteristics of selected Arabica coffee genotypes in Kenya. *Afr. J. Food Sci.* **2000**, *3*, 365–371.
10. Fassio, L.O.; Malta, M.R.; Carvalho, G.R.; Liska, G.R.; Lima, P.M.; Nadaleti, D.H.S.; Pimenta, C.J. Fatty acids profile of *Coffea arabica* L. resistant to leaf rust grown in two

- environments of Minas Gerais, Brazil. *J. Agric. Sci.* **2017**, *9*, 88–98.
11. Bertrand, B.; Vaast, P.; Alpizar, E.; Etienne, H.; Davrieux, F.; Charmetant, P. Comparison of bean biochemical composition and beverage quality of Arabica hybrids involving Sudanese-Ethiopian origins with traditional varieties at various elevations in Central America. *Tree Physiol.* **2006**, *26*, 1239–1248.
  12. Villarreal, D.; Laffargue, A.; Posada, H.; Bertrand, B.; Lashermes, P.; Dussert, S. Genotypic and environmental effects on coffee (*Coffea arabica* L.) bean fatty acid profile: Impact on variety and origin chemometric determination. *J. Agric. Food Chem.* **2009**, *57*, 11321–11327.
  13. Mehari, B.; Redi-Abshiro, M.; Chandravanshi, B.S.; Combrinck, S.; McCrindle, R. Characterization of the cultivation region of Ethiopian coffee by elemental analysis. *Anal. Lett.* **2016**, *49*, 2474–2489.
  14. Mehari, B.; Redi-Abshiro, M.; Chandravanshi, B.S.; Combrinck, S.; McCrindle, R.; Atlabachew, M. GC-MS profiling of fatty acids in green coffee (*Coffea arabica* L.) beans and chemometric modelling for tracing geographical origins from Ethiopia. *J. Sci. Food Agric.* **2019**, *99*, 3811–3823.
  15. Sherge, K.T.; D'heer, J.; Duchateau, L.; Boeckx, P. Influence of growing altitude, shade and harvest period on quality and biochemical composition of Ethiopian specialty coffee. *J. Sci. Food Agric.* **2017**, *97*, 2849–2857.
  16. Mintesnot, A.; Dechassa, N. Effect of altitude, shade, and processing methods on the quality and biochemical composition of green coffee beans in Ethiopia. *East African J. Sci.* **2018**, *12*, 87–100.
  17. Adem, M.W.; Meulenaer, B.D.; Duchateau, L.; Boeckx, P. Effect of altitude on biochemical composition and quality of green Arabica coffee beans can be affected by shade and postharvest processing method. *Food Res. Int.* **2018**, *105*, 278–285.
  18. Bodner, M.; Morozova, K.; Kruathongsri, P.; Thakeow, P.; Scampicchio, M. Effect of harvesting altitude, fermentation time and roasting degree on the aroma released by coffee powder monitored by proton transfer reaction mass spectrometry. *Eur. Food Res. Technol.* **2019**, *245*, 1499–1506.
  19. Gebrekidan, M.; Redi-Abshiro, M.; Chandravanshi, B.S.; Ele, E.; Mohammed, A.M.; Mamo, H. Influence of altitudes of coffee plants on the alkaloids contents of green coffee beans. *Chem. Int.* **2019**, *5*, 247–257.
  20. Tolessa, K.; D'heer, J.; Duchateau, L.; Pascal Boeckx, P. Influence of growing altitude, shade and harvest period on quality and biochemical composition of Ethiopian specialty coffee. *J. Sci. Food Agric.* **2017**, *97*, 2849–2857.
  21. Worku, M.; Meulenaer, B.D.; Duchateau, L.; Boeckx, P. Effect of altitude on biochemical composition and quality of green Arabica coffee beans can be affected by shade and postharvest processing method. *Food Res. Int.* **2018**, *105*, 278–285.
  22. Czerny, M.; Mayer, F.; Grosch, W. Sensory study on the character impact odorants of roasted Arabica coffee. *J. Agric. Food Chem.* **1999**, *47*, 695–699.
  23. Knysak, D. Volatile compounds profiles in unroasted *Coffea arabica* and *Coffea canephora* beans from different countries. *Food Sci. Technol.* **2017**, *37*, 444–448.
  24. Saw, A.K.-C.; Yam, W.-S.; Wong, K.-C.; Lai, C.-S. A comparative study of the volatile constituents of Southeast Asian *Coffea arabica*, *Coffea liberica* and *Coffea robusta* green beans and their antioxidant activities. *J. Essential Oil Bearing Plants* **2015**, *18*, 64–73.
  25. Czerny, M.; Grosch, W. Potent odorants of raw arabica coffee: Their changes during roasting. *J. Agric. Food Chem.* **2000**, *48*, 868–872.
  26. Toci, A.T.; Farah, A. Volatile fingerprint of Brazilian defective coffee seeds: Corroboration of potential marker compounds and identification of new low quality indicators. *Food Chem.* **2014**, *153*, 298–314.
  27. Lee, K.; Shibamoto, T. Analysis of volatile components isolated from Hawaiian green

- coffee beans (*Coffea arabica* L.). *Flavor Fragrance J.* **2002**, 17, 349–355.
28. Bertrand, B.; Boulanger, R.; Dussert, S.; Ribeyre, F.; Berthiot, L.; Descroix, F.; Joët, T. Climatic factors directly impact the volatile organic compound fingerprint in green Arabica coffee bean as well as coffee beverage quality. *Food Chem.* **2012**, 135, 2575–2583.
  29. Dong, W.; Tan, L.; Zhao, J.; Hu, R.; Lu, M. Characterization of fatty acid, amino acid and volatile compound compositions and bioactive components of seven coffee (*Coffea robusta*) cultivars grown in Hainan Province, China. *Molecules* **2015**, 20, 16687–16708.
  30. Toledo, P.R.A.B.; Pezza, L.; Pezza, H.R.; Toci, A.T. Relationship between the different aspects related to coffee quality and their volatile compounds. *Compr. Rev. Food Sci. Food Saf.* **2016**, 15, 705–719.
  31. Franca, A.S.; Oliveira, L.S.; Oliveira, R.C.S.; Agresti, P.C.M.; Augusti, R. A preliminary evaluation of the effect of processing temperature on coffee roasting degree assessment. *J. Food Eng.* **2009**, 92, 345–352.
  32. Toci, A.T.; Farah, A. Volatile compounds as potential defective coffee beans' markers. *Food Chem.* **2008**, 108, 1133–1141.
  33. Yeretizian, C.; Jordan, A.; Badoud, R.; Lindinger, W. From the green bean to the cup of coffee: Investigating coffee roasting by on-line monitoring of volatiles. *Eur. Food Res. Technol.* **2002**, 214, 92–104.
  34. Risticvic, S.; Carasek, E.; Pawliszyn, J. Headspace solid-phase microextraction gas chromatographic time-of flight mass spectrometric methodology for geographical origin verification of coffee. *Anal. Chim. Acta* **2008**, 7, 72–84.
  35. Akiyama, M.; Murakami, K.; Hirano, Y.; Ikeda, M.; Iwatsuki, K.; Wada, A.; Iwabuchi, H. Characterization of headspace aroma compounds of freshly brewed arabica coffees and studies on a characteristic aroma compound of Ethiopian coffee. *J. Food Sci.* **2008**, 73, 335–346.
  36. Brevard, H.; Yeretizian, C. Characterisation of the aroma of green Mexican coffee and identification of mouldy/earthy defect. *Eur. Food Res. Technol.* **2001**, 212, 648–657.
  37. Poyraz, İ.E.; Öztürk, N.; Kiyani, H.T.; Demirci, B. Volatile compounds of *Coffea arabica* L. green and roasted beans. *Anadolu University J. Sci. Technol. C - Life Sci. Biotechnol.* **2016**, 5, 31–35.
  38. Somporn, C.; Kamtuo, A.; Theerakulpisut, P.; Siriamornpun, S. Effects of roasting degree on radical scavenging activity, phenolics and volatile compounds of Arabica coffee beans (*Coffea arabica* L. cv. Catimor). *Int. J. Food Sci. Technol.* **2011**, 46, 2287–2296.
  39. Petisca, C.; Pérez-Palacios, T.; Farah, A.; Pinho, O.; Ferreira, I.M.P.L.V.O. Furans and other volatile compounds in ground roasted and espresso coffee using headspace solid-phase microextraction: Effect of roasting speed. *Food Bioprod. Process.* **2013**, 91, 233–241.
  40. Miller, J.N.; Miller, J.C. *Statistics and Chemometrics for Analytical Chemistry*, 5th ed., Pearson Practice Hall: England; **2005**; pp 55–72.