



## **Oxytetracycline in the possible management of African trypanosomosis**

**Justine T. EKANEM\* and Iyabo S. ADENIRAN**

*Department of Biochemistry, University of Ilorin, Ilorin, Nigeria*

Received 20 October 2003

MS/No BKM/2003/055, © 2003 Nigerian Society for Experimental Biology. All rights reserved.

---

### **Abstract**

We tested the antimicrobial drug Oxytetracycline-HCl against the infections of the protozoan parasite *Trypanosoma brucei* which causes the disease African Trypanosomosis. Oxytetracycline-HCl was used to treat rats infected with the disease at early and late stages of infections. Prophylactic administration was also carried out. The results showed that prophylactic treatment only did not inhibit the proliferation of the parasite but with continued administration kept parasitaemia low. Both prophylactic and early stage treatment of the infection extended the lifespan of the animals. Oxytetracycline-HCl was ineffective for the treatment of the late stage of infection. Oxytetracycline with its iron chelating property could be a useful inhibitor of ribonucleotide reductase, a cell cycle regulated enzyme that in turn regulates cell proliferation. Our experiment suggests that oxytetracycline-HCl could be a good management therapy for the African trypanosomosis if administered at the early stage of infection.

**Key words:** oxytetracycline, sleeping sickness, management

---

\*E-mail: jtekanem@scientist.com Tel: 08033880310

## INTRODUCTION

African trypanosomiasis, the human sleeping sickness is a great threat to more than 60 million people in 36 countries of sub-Saharan Africa (WHO, 1998). The disease is caused by *Trypanosoma brucei*, a protozoan parasite transmitted from host to host by the blood sucking tsetse fly (Seebeck *et al.*, 1999). Trypanosomes are a group of unicellular haemoflagellates of the genus *Trypanosoma* (WHO, 1986). The parasite lives and multiplies in the host where it is able to establish its population through binary cell division (Vickerman, 1985; Ekanem, 2001). During the course of the infection there is a constant rising and falling parasitaemia resulting from the generation of sub-populations with antigenically different variant-specific glycoproteins (VSGs) at the cell surface (Rudenko *et al.*, 1990). This property helps trypanosomes to evade the immune system, shield other proteins e.g. permeases on the surface from immune attack and inhibit phagocytosis (Rudenko *et al.*, 1990). Due to this constant change in variable surface glycoproteins, termed antigenic variation, it has become quite difficult to develop vaccines against the disease trypanosomiasis (Gray and Luckins, 1978; WHO, 1998). Most of the very limited drugs currently in use to combat it are either too expensive or does not satisfy even the minimal standards set today for drug safety and efficiency (Seebeck *et al.*, 1999).

Iron is essential to replication and growth (Romeo *et al.*, 2001) thus all organisms have an ambiguous relationship with it (Hill, 1983; Halliwell and Gutteridge, 1990), a relationship pathogens and their hosts also share (Cabantchik *et al.*, 1999). This dependence of pathogens on their hosts' supply of iron makes them susceptible to the potential bacteriostatic effects of host iron limitation or chelation (Romeo *et al.*, 2001). Much evidence abound to indicate that manipulation of the host iron status does lead to modulation of the proliferation and virulence of bacteria and protozoa (Richardson and Ponka, 1997; Mabeza *et al.*, 1999). The essentiality of iron cannot be overlooked due to its role as prosthetic group in a variety of enzymes, electron transport proteins required for

energy metabolism, for the biosynthesis of deoxyribonucleotides required for DNA synthesis and repair (Aisen *et al.*, 1999), cell viability and proliferation (Richardson and Ponka, 1997). It is important in the G1 phase of the cell cycle and having its role in DNA synthesis linked to the activity of ribonucleotide reductase (RNR), an iron-requiring enzyme responsible for the synthesis of deoxyribonucleotides (Thelander and Reichard, 1979; Nuffer and Cheng 1987; Reichard, 1993; Jordan and Reichard, 1998).

Ribonucleotide reductase plays a central role in DNA replication, cell division and proliferation by providing the deoxyribonucleotide (dNTP) precursors required (Nocentini, 1996; Ingram and Kinnaird, 1999). It is also the only enzyme along the cell division cycle that requires iron not as part of a prosthetic group but as part of the free iron pool (Thelander *et al.*, 1983; Cooper *et al.*, 1996). The role of iron in RNR activity has been elucidated through various studies by iron chelators to alter cellular iron status (Hoyes *et al.*, 1992, Nyholm *et al.*, 1993; Cooper *et al.*, 1996) thus confirming the link between RNR activity and iron transport into proliferating cells (Chitambar *et al.*, 2000). The essentiality of iron in RNR has been proved by studies which showed that when iron delivery is limited, the tyrosyl radical signal diminishes and the R2 subunit exists as a functionally inactive apo-R2 protein (Nyholm *et al.*, 1993).

The structure of Ribonucleotide reductase has been described and classed into three different classes I (a and b), II and III (Chabes *et al.*, 2000). Ribonucleotide reductase has been reported present in *T. brucei* (Hofer *et al.*, 1997; 1998). Trypanosome RNR has binding site for iron (Hofer *et al.*, 1997, Dormeyer *et al.*, 1997). Unlike mammalian cells, bloodstream forms of *T. brucei* have a low iron content (Schell *et al.*, 1991) which is attributed to its lack of cytochromes and like all living organisms trypanosomes require iron for growth (Schell *et al.*, 1991; Steverding, 1998). The delivery of iron into bloodstream forms of trypanosome is mediated by host transferrin that is taken up via a unique receptor (Fast *et al.*, 1999). The growth of trypanosomes has been successfully inhibited

by iron chelation (Breidbach *et al.*, 2002). The enzyme has also been reported as a possible target for drug discovery against sleeping sickness (Ekanem, 2001).

Oxytetracycline belongs to a class of antibiotics of naturally occurring tetracyclines and has found much extensive use in combating both plant (McManus *et al.*, 2002) and animal infections (Lents *et al.*, 2002). The tetracyclines, oxytetracycline inclusive have been found to have strong iron chelating properties (Grenier *et al.*, 2000). Oxytetracycline-HCl could be a possible drug against sleeping sickness more so when a related drug tetracycline-HCl has been reported as a possible cheap drug against the disease, African trypanosomosis (Ekanem *et al.*, 2002; Johnson and Ekanem, 2003). In this report we have investigated the effect of oxytetracycline-HCl on *T. brucei* infected rats at early and late stages of infection. We have also assessed the effect of prophylactic treatment of the drug against the disease.

## MATERIALS AND METHODS

### *Parasite*

Federe strain of African trypanosomes, *Trypanosoma brucei* was obtained from the Veterinary and Livestock Studies Department, Nigerian Institute for Trypanosoma Research (VLS – NITR), Vom near Jos in Plateau State of Nigeria. The parasite was injected into the rats intraperitoneally and maintained through repeated passages into other rats.

### *Experimental animals*

Albino rats (*Rattus norvegicus* species) of an average weight of 230g were used for the experiment. They were obtained from the Animal house of the Department of Biochemistry, University of Ilorin, Ilorin. The animals were kept under standard laboratory conditions and they were maintained on growers mash obtained from Bendel Feed and Flour Mills Ltd., Km 100 Benin-Auchi Road, Ewu, Edo State, Nigeria.

### *Reagents*

Research grade reagents were used for the experiments. NaCl was a product of BDH Chemical Limited, Poole, England. Injectable form of Oxytetracycline-HCl 50mg/ml was a product of Hamexmedica Limited, Teriot Street, London, U.K.

### *Administration of oxytetracycline-HCl*

Oxytetracycline-HCl was injected intraperitoneally into the animals twice daily at various concentrations taking the weights of the rats into consideration.

### *Inoculation of parasites*

The tail of an infected rat was cleaned with a damp cloth and the tip of the tail was cut with a clean pair of scissors. Then the blood was extruded into 0.5ml normal saline solution. The extruded blood and saline solution was swirled to mix and then drawn into a 1ml syringe. A drop of the solution was placed on a microscope slide and observed under the light microscope to ascertain the presence of the parasite in the solution. Inoculation into the peritoneal cavity of an uninfected rat was carried out when parasite suspension contained 3 or 4 trypanosomes per view at x100 magnification.

### *Parasitaemia (parasite count)*

Parasitaemia was determined by counting the number of trypanosomes per view under light microscope at x100 magnification from a thin blood smear obtained from the tip of the tail of an infected rat. The parasitaemia was recorded at 12 hr interval until the rats died.

### *Prophylactic treatment*

Three groups of three rats were treated with the drug for 72 hours at 12 hr interval before trypanosome infection and as the infection progresses until the death of the animals. The dosages used were 15mg/kg, 30mg/kg and 40mg/kg body weight of the rats.

### *Early stage treatment*

Five groups of three rats were treated with 10mg/kg, 15mg/kg, 20mg/kg, 30mg/kg and 40mg/kg body weights of the rats from the time the parasite was first sighted in the blood until the death of the rats.

### *Late stage treatment*

Three groups of three rats were used and treated with 10mg/kg, 20mg/kg and 40mg/kg body weights of the rats 48 hours after the parasite was first sighted in the blood until the animals died.

### *Control*

For each experiment control rats (n=3) were injected with the parasite and allowed to develop without treatment. The parasitaemia was recorded until the death of the animals. Another group of three rats was also treated with oxytetracycline-HCl at 40mg/kg body weight of animals for 14 days without infection.

## **RESULT**

### *Prophylactic treatment*

Fig. 1 shows the result obtained when rats were treated with oxytetracycline-HCl 72 hours before infection. At 15mg/kg body weight, the life span of the animals were increased beyond day 5 at which the control animals died to day 10 and the animals lived longer than control by 5 days. The parasitaemia increased steady from the day of infection and peaked the morning of day 7 after which it reduced to 3 parasites per view on day 8 before the death of the animals on day 10 with increased parasitaemia as indicated by the count for day 9.

At double the dosage i.e. 30mg/kg, the parasitaemia increased steadily from the day of infection and reached its peak by the morning of day 6 then it dropped until one parasite per view was observed on day 10. The rats died on the morning of day 13 compared with the control animals, their life span were extended by 8 days.

With a dosage of 40mg/kg, the parasitaemia gave steady increase and peaked on day 5. The animals died on day 11 having lived 6 days longer than the control animals. The parasitaemia was relatively low throughout the duration of infection when compared with the control.

### *Early stage*

The results obtained when the infected rats were treated with oxytetracycline-HCl immediately parasite was sighted are shown in Fig. 2. At 10mg/kg, the parasitaemia increased steadily and the animals died on day 4. At 15mg/kg there was a steady increase in parasitaemia till it peaked on day 5 when the rats eventually died. Treatment at 20mg/kg showed steady increase in parasitaemia until day 4 which was followed by reduction till day 7 when the parasitaemia increased and it reached its peak before the death of the rats on day 8. The lifespan of the animals were thus extended by 3 days compared with the control animals.

At 30mg/kg, the lifespan of the rats were increased by 4 days. The parasitaemia increased steadily and reached its peak on day 9 after which death occurred. At 40mg/kg dosage, the parasitaemia increased steadily and reached a peak on day 11. The animals died on day 12 having had their lifespan extended by 7 days compared with the control.

### *Late stage*

Fig. 3 shows the result of treatment with oxytetracycline-HCl 48 hours after parasite was first sighted in the blood. At 10mg/kg, the parasitaemia increased steadily and peaked on day 3. The rats died on day 5. At 20mg/kg dosage, the parasitaemia increased steadily and reached its peak by day 3 when death occurred. At double the dosage i.e. 40mg/kg dosage, the parasitaemia increased steadily till it peaked on day 5 when the animals died. There was no extension in the lifespan of all the animals used in the late stage treatment compared with the control.




Fig.1: Prophylactic treatment with Oxytetracycline HCl. Parasitaemia of *T. brucei* infected rats treated twice daily with Oxytetracycline-HCl 72 hours before infection. The parasitaemia (number of cells per view at x100) was monitored until the rats died. Treatments of rats were with 15 mg/kg, 30 mg/kg and 40 mg/kg body weights of animal dosage. Control rats were infected but not treated. Each point is an average count from three rats.




Fig. 2: Early stage treatment with Oxytetracycline-HCl. Parasitaemia of *T. brucei* infected rats treated twice daily with Oxytetracycline-HCl as the infection progressed until death. Treatment of rats started on the day parasite was first sighted in the blood with 10 mg/kg, 15 mg/kg, 20 mg/kg, 30 mg/kg and 40 mg/kg body weight of rats. Control rats were infected but not treated. The parasite count was made at 12-hour interval. Each point is an average count of trypanosomes per view at x100 from three rats.




Fig. 3: Late stage treatment with Oxytetracycline HCl. Parasitaemia of *T. brucei* infected rats treated with Oxytetracycline HCl twice daily as the infection progressed until death. Treatment of rats started 48 hours after parasite was first sighted in the blood with 10 mg/kg, 20 mg/kg and 40 mg/kg body weight of animal dosage. Control rats were infected but not treated. Each point is an average count (number of trypanosomes per view at  $\times 100$ ) from three rats.

#### Oxytetracycline-HCl treatment of uninfected rats

Uninfected rats that were treated with oxytetracycline-HCl for 14 days showed no abnormality and no considerable loss in weight at the end of the experiment. None of the experimental rats survived beyond 14 days.

#### DISCUSSION

When trypanosome enters into a system, the parasite proliferates to establish its population in the infected host through binary cell division (Vickerman, 1985; Ekanem, 2001). They release their toxins into the mammalian system (Ekanem *et al.*, 1996a,b) and eventually kill their host. Iron plays a pivotal role in the synthesis of DNA in the trypanosomal system through the enzyme ribonucleotide reductase (Dormeyer *et al.*, 1997; Hofer *et al.*, 1997). The deprivation of a biological system of iron brings about the inactivation of the ribonucleotide reductase enzyme (Cooper *et al.*, 1996; Chabes *et al.*, 2000). It has also been established

previously that the deprivation of iron inhibits the growth of cells, even those of trypanosomes (Fast *et al.*, 1999). Iron chelation has thus been adopted as an attractive and possible strategy for new antiparasite drugs and treatment of sleeping sickness (Ekanem, 2001; Breidbach *et al.*, 2002). Grenier *et al.* (2000) reported iron chelating properties of oxytetracycline.

Our study revealed that early stage treatment and prophylactics at high dosage of 30mg/kg and 40mg/kg extended the life span of the animals. In the case of the prophylactic treatment very low parasitaemia with almost total clearance of parasite in the system was observed. This is probably as a result of the drug chelating iron present within the system since it had been introduced into the host system long before parasite invasion of the system unlike during the early stage treatment whereby the drug was not administered until the parasite was sighted present in the blood. Also, results obtained showed that at high dosage of 30mg/kg and 40mg/kg body weight the parasitaemia was less

than those recorded with the lower dosage of 10mg/kg, 15mg/kg and 20 mg/kg although the administration of 15 mg/kg dosage for prophylactic treatment brought about death a day before the animals treated with 30 mg/kg body weight dosage died (Figs. 1 & 2). This suggests that high dosage of oxytetracycline is considerably effective for removal of the parasite from the system.

Oxytetracycline seems to have been recognized as a potential trypanocide or combined therapy trypanocide judging by the interest it aroused in some non-professionals who exploited it to combat the disease trypanosomosis. Some Maasai pastoralists in Kenya administered oxytetracycline with other trypanocides as soon as the disease was recognized (Rodericks *et al.*, 2000) but not as prophylactics. Our study established that the use of oxytetracycline-HCl for prophylactics kept parasitaemia low and considerably extended the lifespan of infected

animals. Severity of the disease correlates with the level of parasitaemia (Murray *et al.*, 1988).

The late stage of trypanosomosis is usually characterized by high parasitaemia, reduced packed cell volume, and pronounced haemolytic anemia (Murray *et al.*, 1988) due to the lysing of the erythrocyte of the animals. The late stage treatment by oxytetracycline-HCl was not effective since death resulted at about the same time as the control (Fig. 3). Obviously, the toxic substances released by the parasites into the blood stream (Ekanem, 1989; Ekanem *et al.*, 1986a,b) completely overwhelms the hosts' system ultimately resulting in the death of the animals as the disease takes its toll on the animals.

We suggest that oxytetracycline-HCl could be a useful drug in the management of African sleeping sickness since it extends the lifespan of infected rats when treated at early stage.

## REFERENCES

**Aisen, P., Wessling- Resnick, M and Leibold, E. A. (1999)** Iron Metabolism. *Curr. Op. Chem. Biol.* **3**:200-206.

**Breidbach, T., Krauth-Siegel, R. L and Steverding, D. (2002)** Ribonucleotide reductase is regulated via the R2 subunit during the life cycle of *Trypanosoma brucei*. *FEBS. Lett.* **473**:212-216.

**Cabantchik, Z. I., Moody- Haupt, S. and Gordeuk, V. R. (1999)** Iron chelators as anti-infectives: Malaria as paradigm. *FEMS Immun. Med. Microbiol.* **26**:289-298.

**Chabes, A., Domkin, V., Larsson, G., Liu, A., Graslund, A., Wijmenga, S. and Thelander, L. (2000)** Yeast ribonucleotide reductase has heterodimeric iron-radical-containing subunit. *Proc. Natl. Acad. Sci. USA* **97**:2447-2479.

**Chitamber, C.R., Wereley, J.P., Heiman, T., Antholine, W.E and O'Brien, W.J. (2000)**

Cellular adaptation of down-regulated iron transport into lymphoid leukaemia cells : effects on the expression of the gene for ribonucleotide reductase. *Biochem. J.* **345**:681-685.

**Cooper, C. E., Lynagh, G. R., Hoyes, K. P., Hider, R. C., Cammack, R. and Porter, J. B. (1996)** The relationship of intracellular iron chelation to the inhibition and regeneration of human ribonucleotide reductase. *J. Biol. Chem.* **271**: 20291-20299.

**Dormeyer, M, Schoneck, R., Dittmar, G.A.G., Krauth-Siegel, R. L., (1997).** Cloning, sequencing and expression of ribonucleotide reductase R2 from *Trypanosoma brucei*. *FEBS Lett.* **415**: 449-453.

**Ekanem, J. T. (1989)** Extracellular fractions derived from *Trypanosoma brucei* activate erythrocyte Ca<sup>+2</sup>ATPase. *Med. Sci. Res.* **17**:739-740.

- Ekanem, J. T. (2001).** Ribonucleotide reductase as target for drug discovery against African sleeping sickness. *NISEB Journal* 1:287-292.
- Ekanem, J.T., Akanji, M.A and Odutuga, A.A (1996a)** Elevated erythrocyte calcium and Ca<sup>2+</sup>-ATPase activity in *Trypanosoma brucei*-infected rats. *Biomed. Lett.* 54:7-11
- Ekanem, J. T., Odutuga, A. and Akanji, M. A. A. (1996b)** Extracellular proteins of *Trypanosoma brucei* origin lyse erythrocytes of rat *in vitro*. *Biokemistri* 6:21-29.
- Ekanem, J. T., Johnson T. O. and Obaleye, J. A. (2002)** Tetracycline: possible cheap drug in the management of African sleeping sickness. *NISEB Journal* 2:83-87.
- Fast, B., Kremp, K., Boshart, M. and Steverding, D. (1999)** Iron-dependent regulation of transferrin receptor expression in *Trypanosoma brucei*. *Biochem. J.* 342:691-696.
- Gray, A. R. and Luckins, A. G. (1978)** Antigenic variations in Salivarian trypanosomes. In: *Biology of the kinetoplastida*. Lumsden W.H.R. & Evans D. N. (Eds). New York. Acad. Press. 1:493.
- Grenier, D., Huot, M. and Mayrand, D. (2000)** Iron chelating activity of Tetracyclines and its impact on the susceptibility of *Actinobacillus antinomycetemcomitans* to these Antibiotics. *Antimicrob. Agents Chemother.* 106:763-766.
- Halliwell, B. and Gutteridge, J.M.C. (1990)** Role of free radicals and catalytic metal ions in human diseases: An overview. *Methods Enzymol.* 86:1-85.
- Hill, H. A. O. (1983)** Biological Aspects of metals and metal-related diseases (Sarkar B.ed.) p 15-21. Raven Press Ltd., New York.
- Hofer, A., Ekanem, J. T. and Thelander, L. (1998)** The allosteric regulation of *Trypanosoma brucei* Ribonucleotide reductase studied *in vitro* and *in vivo*. *J. Biol. Chem.* 273:34098-34104.
- Hofer, A., Schmidt, P. P., Graslund, A. and Thelander, L (1997)** Cloning and characterization of the R1 and R2 subunits of ribonucleotide reductase from *Trypanosoma brucei*. *Proc. Natl. Acad. Sci. USA.* 94:6959-6964.
- Hoyes, K. P., Hider, R. C. and Porter, J. B. (1992)** Cell cycle synchronization and growth inhibition by 3-hydroxypridin-4-one iron chelators in leukaemia cell lines. *Cancer Res.* 52:4591-4599.
- Ingram, G. M. and Kinnaird, J. H. (1999)** Ribonucleotide reductase. A new target for antiparasite therapies. *Parasitol. Today* 15:338-342.
- Johnson T. O. and Ekanem J. T. (2003)** Effect of tetracycline on late-stage African trypanosomiasis in rats. *Biokemistri* 14:51-56
- Jordan, A. and Reichard, P. (1998)** Ribonucleotide reductase. *Ann. Rev. Biochem.* 67:71-98.
- Lents, C. A., Wettermann, P. R., Paape, M. J., Vizcarra, J. A., Looper, M. L., Buchanna, D.S. and Lusby, K. S. (2002)** Efficacy of intramuscular treatment of beef cows with Oxytetracycline to reduce mastitis and to increase calf growth. *J. Anim. Sci.* 80:1404-1412.
- Mabeza, G. F., Loyevsky, M., Gordeuk, V. R. and Weiss, G. (1999)** Iron chelation therapy for Malaria: a review. *Pharmacol. Ther.* 81:53-75.
- McManus, P. S., Stockwell, V. O., Sundin, G. W. and Jones, A. L. (2002)** Antibiotics use in Plant Agriculture. *Annu. Rev. Phytopathol.* 40:443-465.
- Murray, M., Morrison, W. Y. and White Kiv, D. V. (1988)** Host susceptibility to African trypanosomiasis; Trypanotolerance. *Adv. Parasitol.* 21:52-57.
- Nocentini, G. (1996)** Ribonucleotide reductase inhibitors – new strategies for cancer chemotherapeutics. *Crit. Rev. Oncol. Hematol.* 22:89-126.



**Nuffer, L. M. and Cheng, Y. C. (1987)** Inhibitors of ribonucleotide diphosphate reductase activity (Cory, J. G. and Cory, A. H. eds.). pp 37-54.

**Nyholm, S., Mann, G. J. Johansson, A. G., Bergerson, R. J., Graslund, A. and Thelander, L. (1993)** Role of ribonucleotide reductase in inhibition of mammalian cell growth by potent iron chelators. *J. Biol. Chem.* **268**:26200-26205.

**Reichard, P. (1993)** From RNA to DNA, why so many ribonucleotide reductases. *Science* **260**:1773-1777.

**Richardson, D. R. and Ponka, P. (1997)** The molecular mechanisms of the metabolism and transport of iron in normal and neoplastic cells. *Biochim. Biophys. Acta.* **1331**:1-40.

**Rodericks, S., Stevenson P., Mwendia, C. and Okech, G. (2000)** The use of trypanocides and antibiotics by Maasai pastoralists. *Trop. Anim. Hlth. Prod.* **32**:361-374.

**Romeo, A. M., Christen, L., Niles, E. G. and Kosman, D. J., (2001)** Intracellular chelation of iron by Bipyridyl inhibits DNA virus replication (Ribonucleotide reductase maturation as a probe of intracellular iron pools). *J. Biol. Chem.* **276**:24301-24308.

**Rudenko, G., Cross, M. and Borst, P. (1998)** Changing the end antigenic variation. Orchestrated at the telomeres of African trypanosomes. *Trends Microbiol.* **6**:113-116.

**Schell, D., Borowy, N. K. and Overath, P. (1991)** Transferrin is a growth factor for the bloodstream form of *Trypanosoma brucei*. *Parasitol. Res.* **77**: 558-560.

**Seebeck, T., Naula, C., Shalaby, T. and Gong, K. (1999)** Human sleeping sickness – current approaches to chemotherapy. *Nova.Acta. Leopoldina. NF78, NT* **307**:227-241.

**Steverding, D. (1998)** Bloodstream forms of *Trypanosoma brucei* require only small amounts of iron for growth. *Parasitol. Res.* **84**:59-62.

**Thelander, L., Graslund, A. and Thelander, M. (1983)** Continual presence of oxygen and iron required for mammalian ribonucleotide reduction: Possible regulation mechanisms. *Biochem. Biophys. Res. Comm.* **110**:859-865.

**Thelander, L., and Reichard, P. (1979)** Ribonucleotide reductase. *Ann. Rev. Biochem.* **48**:133-158.

**Vickerman, K. (1985)** Developmental cycles and biology of pathogenic trypanosomes. *Brit. Med. Bull.* **41**:105-144.

**W.H.O. (1986)** Epidemiology and Control of African Trypanosomosis. Report of a WHO Expert Committee Technical Report Series World Health Organisation, Geneva.

**W.H.O. (1998)** Control and Surveillance of African Trypanosomosis. Report of WHO expert committee. *WHO Tech. Rep. Ser.* 881.I-VI, 1-114.