

Bacterial Cell Envelope Lysis and Hemotoxicity of Peptides Previously isolated From African Catfish, *Clarias gariepinus*

Hedmon Okella^{*a}, Clement Olusoji Ajayi^a, Hilda Ikiriza^a, Andrew Glory Mtewa^e, Bruhan Kaggwa^{a,c}, Jacqueline Aber^{a,d}, Charles Drago Kato^b, Patrick Ogwang Engeu^a

^aPharm-Biotechnology and Traditional Medicine Centre, Mbarara University of Science and Technology, Mbarara, Uganda, ^bCollege of Veterinary Medicine, Animal Resources and Biosecurity, Makerere University, Kampala, Uganda, ^cDepartment of Pharmacy, College of Health Sciences, Makerere University, Kampala, Uganda, ^dDepartment of Pharmacy, Faculty of Medicine, Gulu University, Gulu, Uganda, ^eChemistry Section, Malawi Institute of Technology, Malawi University of Science and Technology, Limbe, Malawi

Correspondence to Hedmon Okella (hokella@std.must.ac.ug)

ABSTRACT

Background: The skin mucus layer of fish is endowed with biologics including, Antimicrobial peptides (AMPs) that offer a first line of defence against pathogens. Such peptides can either inhibit bacterial growth or completely kill the bacteria and hence are regarded as a viable alternative to traditional antibiotics, in addressing the ever-increasing incidences of antimicrobial resistance. However, one of the major hurdles to AMPs use is their poor haemolytic profile. As a result, a thorough evaluation of prospective AMPs' bacterial cell membrane disruption and hemolytic potentials in the early phases of drug discovery is critical. The current study presented cell membrane destruction as well as hemo-compatibility of antimicrobial peptides previously isolated from skin mucus of African catfish, *Clarias gariepinus*.

Methods: A previously isolated antimicrobial peptide in the skin mucus of African catfish, *C. gariepinus* were profiled using 15% Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE). The electrical conductivity and alkaline phosphatase assays were utilised to measure bacterial cell envelope lysis activity as a classical mode of action of the antimicrobial peptides. Afterwards, fresh Rabbit blood cells were then utilised for in vitro hemolytic assay.

Results: The peptides were found to be about 5 kDa molecular weight with, ability to damage the bacterial cell envelope causing significant leakage in periplasmic alkaline phosphatase enzyme and cytoplasmic electrolytes. Even at the highest peptide extract concentration of 100 µg/mL, no significant hemolysis was observed on the fresh rabbit blood cells [3.63%; P>.05], signifying their safety on normal mammalian cells.

Conclusion: The findings of this study pointed out that antimicrobial peptides in skin mucus of *C. gariepinus* are potentially safe source of antimicrobial drug leads; however, further studies are still required to search for possibly maximum dose that is safe to host cells but still effective against infecting bacteria.

INTRODUCTION

Pathogenic bacteria, fungi, viruses and protozoa flourish in the same aquatic ecosystem in which fish live.¹ Such habitats predispose fish to higher infection risks compared to terrestrial vertebrates,^{2,3} thereby calling for a more effective natural defence mechanism. Given their underdeveloped adaptive immunity, fish mostly utilise a more complex innate defence mechanism comprising three major components: physical, phagocytic cells and chemical mediators.^{4,5} Over decades, it has been demonstrated that the fish mucus layer on the skin, gills, nose and gut remains the principal first-line physical defence against infections.^{6,7} The mucus layer comprises a cocktail of biologics including peptides, acute phase proteins, glycoproteins, enzymes, immunoglobulin, lectins and hemolysin that are essential in contributing potential leads in the field of drug discovery.^{8,9} However, the aquatic exploration has largely been for food sourcing with far less attention to fish-derived drug leads. This

has left little known about the toxicological profiles and bactericidal efficacy of fish-derived antimicrobial peptides. Yet, exploring such may be of relevance to new alternative antimicrobial drug candidates in the era of new antimicrobial drug search.¹⁰

To this effect, the fish skin mucus has increasingly enthralled the search for new potential bactericidal drug candidates. Natively available fish biologics are gradually gaining pursuits as potential therapeutic candidates, due to their safety, low cost of production and rapid mode action¹¹.

Accordingly, several studies have investigated the antimicrobial activity of different fish species, notable examples include antimicrobial activity of skin mucus extracts of *Hypophthalmichthys nobilis*,^{12,13} *Clarias batrachus*,¹⁴⁻¹⁶ *Heteropneustes fossilis* and *Clarias batrachus*,¹⁷ *Channa striatus*,¹⁸ *Catla catla*,¹⁹ *Rutilus frisii*,²⁰ *Periophthalmodon schlosseri*²¹ and *Anabas testudineus*²² among others.

However, studies on the antimicrobial potentials of *Clarias gariepinus* inhabiting any of the three major lakes of Uganda have not been reported. Besides, the bacterial cell lytic activity just like hemo-compatibility of such peptide extracted from the skin mucus of *C. gariepinus* in the African region remains unknown. The current work builds on a previous study where antimicrobial peptides from the skin mucus of the African catfish, *C. gariepinus* sourced from Lake Albert, Uganda were isolated.²³ In fact, the previous study was only limited to Sephadex G-25 peptide isolation. Therefore, in the quest for potential drug leads in the era of antimicrobial resistance, the current study presents the first report on bacterial cell envelope disruption and hemolytic profiles of such peptides.

MATERIALS AND METHODS

Preparation of antimicrobial peptides

Two lyophilized archived antimicrobial peptide fractions (Peak I and Peak II) previously isolated from 24 live mature *C. gariepinus* (Figure 1) in the family *Clariidae* (mean weight 300.50 ± 5.98 g, mean length 30.60 ± 2.11 cm) using Sephadex G-25 gel filtration chromatography,²³ were utilised in the present study. Guided by our previous study, only fractions (Peak I) with demonstrable antimicrobial activity *E. coli* [(MIC: 0.31 ± 0.16 µg/ml) and *S. aureus* (MIC: 1.99 ± 0.13 µg/ml)]²³ and clear band upon resolving on a 15 (%) Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) was considered. In addition, only the peptides extracted through Solid Phase Extraction (SPE) technique prior to Sephadex G25 bio-guided fractionation were employed in this study. This was due to the fact that cartridge's solid-phase hydrophobic matrix optimally captures the hydrophobic peptides that are later recovered through organic solvent systems.²⁴

Peptide Profiling

To establish the peptide profiles, Sodium Dodecyl Sulfate-Polyacrylamide Gel Electrophoresis (SDS-PAGE) was run as described by Laemmli with minor modifications.²⁵ A 15% separating gel and 4% stacking gel was used. Here, 0.01 g of lyophilized Peak I fractions and 0.01 g of lyophilized C₁₈ SPE elute were dissolved in 50 µl deionized water, respectively. Thereafter, 40 µl of samples and 40 µl of sample loading buffer [12% SDS (w/v), 6% mercaptoethanol (v/v), 30% glycerol (w/v), 0.05% Coomassie blue G-250, 150 mM Tris/HCl, pH 7.0] were mixed, boiled for 5 minutes at 100 °C utilising heating block (Biobase, Shangdon, China). Then, 10 µl of the samples were then loaded into each well of the gel. Later, 8 µl of pre-stained SDS-PAGE standard markers (Thermo Fisher Scientific, Waltham, USA) were included to estimate the molecular mass of the proteins.

Electrophoresis was run at a constant voltage of 150 V for 55 minutes, until the dye front migrated to 2 cm from the bottom of the gel (Bio-Rad, Hercules, USA). Later, the gel was directly stained in 100 ml staining solution (1 g Coomassie R250, 30 ml, methanol, 65 ml deionized water, 50 ml acetic acid) for 4 hours.²⁶ Peptides were then directly visualized in destaining solution I (10 ml methanol, 10 ml Glacial acetic acid, and 80 ml deionized water) at 55 rpm rotation at room temperature; 25 °C (Edmund Buhler GmbH, Bodelshausen, Germany), until fairly clear bands were observed.

Alkaline Phosphatase Activity

Cell wall destruction was determined by measuring Alkaline Phosphatase Activity (ALP). An alkaline phosphatase assay kit (QuantiChrom BioAssay Systems, Hayward, USA) with *p*-nitrophenylphosphate (*p*NPP) as a substrate was used to measure the ALP activity of the cell lysate.²⁷ Briefly, 12 hour cultured *E. coli* were dissolved in fresh sterile medium to 106 CFU/ml. About 5 mg/ml of laboratory prepared antimicrobial peptide in 0.01 M Phosphate Buffer Solution (PBS) was added to the medium and cultured at 37°C (Heraeus, Hanau, Germany) for 30 minutes. The reaction mixture (1 ml) was then collected every hour, centrifuged at 5000 x g (Eppendorf, Dubai, UAE) for the next 20 hours. Thereafter, 150 µl of disodium *p*-nitrophenyl phosphate substrate buffer was added in the 50 µl supernatant and the mixture incubated in a 40 °C water bath (Grants, Cambridge-shire, UK) for 4 minutes. Absorbance was then measured at 405 nm using Microtiter plate reader (Biochrom, Cambridge, England) while 0.01 M PBS was used as a negative control. The experiment was carried out in triplicate before the Mean and Standard Error of Mean (SEM) were then calculated.

Electrical Conductivity Detection

Electrical conductivity of cytoplasmic fluid was determined as previously described by Lee et al.²⁸ Here, a 12 hour cultured *E. coli* was dissolved in fresh sterile medium to 106 CFU/ml. Thereafter, 0.05 mg/ml of antimicrobial peptides in 0.01 M PBS was added and cultured at 37°C (Heraeus, Hanau, Germany) for 20 minutes. The culture mixture (1 ml) was then collected every hour, centrifuged at 5000 x g (Eppendorf, Dubai, UAE) for the next 20 hours. The electrical conductivity was detected with Seven Go Duo SG-23 digital conductivity meter (Mettler-Toledo, Columbus, USA). The experiment was carried out in triplicate before the Mean and Standard Error of Mean (SEM) was then calculated, taking 0.01 M PBS as a negative control.

FIGURE 1. African Catfish



Clarias gariepinus (Burchell, 1822). Common name, African Catfish; Lango, Rec Lango/Twang; Luganda, Semutundu; Alur, Nyaii; Kiswahili, Kambale. A benthopelagic freshwater scales-less fish, with four pairs of slender, whisker-like sensory organs (barbel) near the mouth. An original photo by Hedmon Okella

Hemolytic Activity Testing

Hemolytic activity was assayed with a modified Rabbit blood cells method described by Lin and others.²⁹ Briefly,

fresh Rabbit blood cells were obtained by centrifuging whole blood from live rabbit in EDTA-coated Vacutainer (Becton & Dickinson, New Jersey, USA) in a refrigerated microfuge (Eppendorf, Dubai, UAE) at 211 x g for 5 min at room temperature (25 °C). Blood cells were washed three times with PBS, and then diluted with PBS to a blood cell concentration of approximately 10% (v/v).

A portion of the Rabbit blood cell suspension (500 µl) was transferred to six micro-centrifuge tubes (CellTreat, Massachusetts, USA), and mixed with 500 µL of antimicrobial peptide extract solution in 0.01 M PBS at the desired concentrations (1, 20, 40, 60, 80 and 100 µg/ml). The tubes were then incubated at 37°C (Heraeus, Hanau, Germany) to induce hemolysis. After 30 minutes of incubation, non-hemolysed blood cells were separated by centrifugation at 211 x g for 5 minutes at room temperature. Aliquots (100 µl) of the supernatant were transferred to a 96-well plate (Corning, New York, USA), and hemoglobin release was monitored by measuring the absorbance of the supernatant at 540 nm using microtiter plate reader (Biochrom, Cambridge, England). A blood cell solution treated with 1% Triton X-100 (to induce 100% lysis) was employed as a positive control, and an untreated blood cell suspension in 0.01 M PBS alone was used as negative control. Each assay was performed in triplicate for three independent experiments, and data were expressed as the mean and SEM. The percentage of hemolysis was calculated using the following formula³⁰

$$\text{Hemolysis (\%)} = ((A_e - A_n) / (A_p - A_n)) \times 100$$

Where A_e is Absorbance of the extracts, A_p is the Absorbance of the positive control and A_n is the absorbance of the Negative control.

Statistical Analysis

Tableau Software v2019.4 (Tableau, Seattle, U.S.A) was used to present data. All experiments were carried out in triplicate and expressed as Mean and Standard Error of Mean (SEM) using Prism 5.0 Statistical software (GraphPad, San Diego, U.S.A) and SPSS v16.0 (IBM, Chicago, U.S.A) were used to compare the means, in which a one-sample T-test was performed to determine the significance of the hemolytic activity of fish-derived antimicrobial peptides on fresh mammalian blood cells.

Ethical Consideration

Ethical approval and permission to conduct this study was obtained from the Mbarara University of Science and Technology Research Ethics Committee (22/11-18) and registered with the Uganda National Council for Science and Technology (HS449ES).

RESULTS

Study selection and Characteristics

In the present study, lyophilized Sephadex G-25 isolated antimicrobial peptides belonging to distinct peaks two peaks (Peak I and Peak II),²³ were utilised. Sephadex G-25 chromatographic resin was used and two prominent peaks (peaks I and II) were observed with absorbance measured at 280 nm. Only Peak I antimicrobial peptides demonstrated an antimicrobial activity with a MIC of 0.31 ± 0.16 and 1.99 ± 0.13 µg/ml on *E. coli* and *S. aureus*, respectively, whereas no antimicrobial activity was detected in peak II. The present study therefore, utilised

only peak I archived samples of our previous study.²³

Peptide Profiling


The SDS-PAGE peptide profiles of the previously extracted peptides from the skin mucus of African catfish are shown in Figure 2. Low molecular weight peptides (about 5 kDa) were observed as a clear band for the Sephadex G-25 purified extract unlike in the case of the 5 kDa-ultrafiltered C18 SPE elute.

Alkaline Phosphatase Activity and Electrical Conductivity

Throughout the culture process, the alkaline phosphatase content in the control group (0.01 M PBS), remained at a low level (below 5 iu/l), indicating the cell wall was intact (Figure 3a). In the antimicrobial peptide treated group, the alkaline phosphatase activity increased exponentially after an hour of treatment, as more alkaline phosphatase enzyme continues to leak out of the periplasm. Figure 3b showed the results of conductivity measurements.

Conductivity of the control group (0.01 M PBS) remained relatively stable at low level throughout the culture process. On the other hand, just after an hour of treatment with the antimicrobial peptides, the electrical conductivity was elevated significantly when compared to the control group, as the large volumes of the electrolyte continues to leak out of the cytoplasm during the culturing process.

FIGURE 2. A 15 (%) SDS-PAGE of peptides extracted from African catfish



The SDS-PAGE molecular marker lane (M), 5 kDa-ultrafiltered C18 SPE elute (1), C18 SPE and Sephadex G-25 isolated peptides (2). The extracted peptides were about 5 kDa.

Hemolytic activity of Antimicrobial Peptides

When the peptide extracts of *C. gariepinus* skin mucus were incubated with fresh rabbit blood cells, the percentage hemolysis increased with increase in peptide concentration (Figure 4). However, the increase in hemolytic activity was not significant even at the maximum concentration (100 µg/ml; $P > .05$) at a zero percent hemolysis test value. This signifies the safety of the fish-derived antimicrobial peptides (tested concentrations) on the Rabbit blood cells.

DISCUSSION


The present study demonstrated the capability of antimicrobial peptides extracted from *C. gariepinus* skin mucus in disrupting the bacterial cell wall and cell membrane. The previous study employed Solid Phase Extraction (SPE) of the antimicrobial peptides optimized the isolation of hydrophobic peptides. This is because solid phase of the SPE cartridge is a hydrophobic matrix that captures hydrophobic peptides that are later recovered by organic solvents after washing. Hydrophobicity is essential in bacterial cell membrane non-polar interactions.³¹ Increased hydrophobicity up to the optimal threshold leads to enhanced ability of the peptides to associate with cellular hydrophobic lipid bilayer,³² and most importantly, allows such peptide easily cross the membrane.^{33,34} Much as, the present study was able to explore the action of these peptides on bacterial cell envelope (Figure 4), archived samples were utilized. Such samples are subject to deterioration. To mitigate this limitation, only lyophilized samples stored at 4 °C with clear band profiles on the SDS-PAGE and detected antimicrobial activity were considered for this study.

To gain insight into the mode of action of peptides, the bacterial cell envelope lysis of the *E. coli* was investigated by detecting the content of the extracellular alkaline phosphatase and electrolytes. The choice was guided by the fact that cell envelope damage is the commonest killing mechanism of peptides with antimicrobial or host defence potentials.³⁵ Alkaline phosphatase (ALP) catalyzes the hydrolysis of phosphate esters in an alkaline environment, resulting in the formation of an organic radical and inorganic phosphate. In this study, a rapid increase in the alkaline phosphatase activity in the *E. coli* culture medium within an hour of peptide treatment was observed, this persisted till the 13th hour of treatment. This hourly trend monitoring is hinged on the fact that, antimicrobial peptides firstly targets the cell membrane,³⁶ and later interact electrostatically interact with negatively charged microbial cell membrane, prior to membrane destruction. Subsequently, the antimicrobial peptides may align parallel to the cell membrane like a carpet where it destroys the cell membrane in a 'detergent-like' manner,³⁷ or they penetrate the bilayer of the cell membrane as described in the pore model where their either form Toroidal pore/Wormhole³⁸ or aggregates into channel forming multimers.³⁹ Such sequence of events requires interval monitoring and an hour monitoring has been reported as ideal.⁴⁰

Given that, alkaline phosphatase is localised between the cell membranes and cell walls of the bacteria⁴¹ and and any disruption of the cell wall penetrability, leads to its discharge out of the bacteria. Therefore, *C. gariepinus* skin mucus peptides were capable of destroying the cell wall.


Similar increase in ALP activity, have previously reported for *Porphyra yezoensis* peptides⁴⁰ and Ornidazole (ORZ),⁴² Linalool against *Pseudomonas fluorescens*,⁴³ Dihydromyricetin against Food-Borne Bacteria⁴⁴ and in AIEgen-Peptide Conjugate⁴⁵ among others, signifying bacterial cell wall destruction⁴⁶ and their potential application as far as sourcing of new antimicrobial drug candidates in the era of antimicrobial drug resistance is concerned.

FIGURE 4. Percentage hemolysis of Rabbit blood cells by peptide extracts of *C. gariepinus*.



T-1% Triton X-100. The concentration of 1-100 µg/ml did not hemolyze the Rabbit blood cells. The data are expressed as Mean ± SEM

Besides, cell wall distraction increases the cell membrane permeability resulting to cytoplasmic outflow and hence, a rise in extracellular electrolytes.^{47,48} Monitoring of the electrical conductivity of the extracellular medium can therefore be employed as a measure of the bacterial cell membrane disruption.⁴⁹ In this study, within an hour of peptide treatment, the electrical conductivity of *E. coli* culture medium significantly increased suggesting the cell wall and cell membrane breakage amounting to cytoplasmic fluid outflow, cell collapse and death. The gradual increase in conductivity after an increase, is possibly due to the fact that majority of the cytoplasmic electrolytes are lost within the first hour of membrane shock. This is in line with studies on OVTp12 peptide on *E. coli* and *S. aureus*;⁵⁰ Metallic oxide powders on *Candida albicans* NBRC1060, *Saccharomyces cerevisiae* NBRC1950, *Aspergillus niger* NBRC4067 and *Rhizopus stolonifer* NBRC4781;⁵¹ *Porphyra yezoensis* peptides on *S. aureus*;⁴⁰ Black Paper Essential Oils (BPEO) on meat-borne *E. coli*⁵² and antibiotics like Penicillin G, Nalidixic acid, Rifampicin on *Escherichia coli* 745 and *Staphylococcus aureus* 9779.⁵³ Moreover, cell damage as well ignites transient pore formation and freight of peptides into the cell for intracellular target interactions.⁵⁴ Much as, it would be essential to explore the interactions of the isolated peptides with the intracellular targets, it was beyond the scope of this study to explore peptide-intercellular target

FIGURE 3: Effect of Antimicrobial Peptide Treatment on Cell Wall and Cell Membrane Integrity

(a) Alkaline phosphatase activity in the culture medium, (b) Electrical conductivity in culture medium. PBS=0.01 M Phosphate Buffer Solution. Both the Alkaline phosphatase activity and Electrical conductivity were measured each hour for 20 hours. The data are expressed as Mean \pm SEM.

interactions as dictated by the facility constrains.

Toxicity against mammalian cells is the chief drawback that impedes most peptides from penetrating the pharmaceutical market.^{55,56} To this effect, preliminary assessment of any potential lead biologic or their sources for hemo-compatibility is worthy. In this study, *C. gariepinus* peptide extracts was found to be non-hemolytic even at the highest concentration assayed (100 μ g/ml), signifying their potential safety towards mammalian red blood cells and hence, relevance in sourcing of novel antimicrobial drug leads. The maximum concentration assayed (100 μ g/ml), was empirically arrived at based on the previous studies.^{57,58} Non-hemotoxic findings have as well been demonstrated in the skin mucus extract of Striped Dwarf Catfish, *Mystus vittatus*;⁵⁷ and Marine Catfish, *Tachysurus Dussumieri*.⁵⁹

To the contrary, the mucus extracts of Spotted sea catfish, *Arius maculatus*;⁵⁸ and venomous fish such as Stonefish, *Synanceia verrucosa*;⁶⁰ Lionfish, *Pterois volitans*;^{61,62} Scorpionfish, *Scorpaena plumieri*;⁶³ Pufferfish, *Akifugu rubripes*⁶⁴ have been reported as hemotoxic and lethal. The biologics in the hemotoxic extracts penetrates the deeper hydrophobic core of the mammalian cell membrane, once their hydrophobicity exceeds the optimal threshold.⁶⁵ Therefore, the antimicrobial peptides of *C. gariepinus* were below the hydrophobicity threshold for the studied concentrations. However, due to facility limitations, we were unable to determine the optimal threshold hydrophobicity for the extracted peptides.

CONCLUSION

The present study for the first time unveiled the toxicity profiles and possible mode of action of the previously isolated African catfish antimicrobial peptides. The peptides demonstrated outstanding lytic activity on

the bacterial cell envelope. Furthermore, they were non-hemolytic to normal mammalian blood cells. The current study therefore, fronts the novel source of safe and efficacious antimicrobial drug leads of potential applications to food, medicinal and pharmaceutical industries. However, we recommend the search for potentially maximum dose that is safe to the host cells but still effective against bacteria.

Acknowledgement

Hedmon Okella recognises a Fellowship for his post-graduate studies by Pharm-BioTechnology and Traditional Medicine Centre (PHARMBIOTRAC), Mbarara University of Science and Technology (MUST).

REFERENCES

- Magnadóttir B. Immunological control of fish diseases. *Mar Biotechnol.* 2010;12(4):361-379. doi:10.1007/s10126-010-9279-x
- Austin B, McIntosh D. Natural antibacterial compounds on the surface of rainbow trout, *Salmo gairdneri* Richardson. *J Fish Dis.* 1988;11:275-277.
- Behringer DC, Karvonen A, Bojko J. Parasite avoidance behaviours in aquatic environments. *Philos Trans R Soc B Biol Sci.* 2018;373(1751). doi:10.1098/rstb.2017.0202
- Smith NC, Rise ML, Christian SL. A comparison of the innate and adaptive immune systems in cartilaginous fish, ray-finned fish, and lobe-finned fish. *Front Immunol.* 2019;10(October). doi:10.3389/fimmu.2019.02292
- Dalmo RA, Ingebrigtsen K, Bogwald J. Non-specific defence mechanisms in fish, with particular reference to the reticuloendothelial system (RES). *J Fish Dis.* 1997;20(4):241-273. doi:10.1046/j.1365-2761.1997.00302.x

6. Cipolari OC, de Oliveira Neto XA, Conceição K. Fish bioactive peptides: A systematic review focused on sting and skin. *Aquaculture*. 2020;515:734598. doi:10.1016/j.aquaculture.2019.734598
7. Dash S, Das SK, Samal J, Thatoi HN. Epidermal mucus, a major determinant in fish health: A review. *Iran J Vet Res*. 2018;19(2):72-81. doi:10.22099/ijvr.2018.4849
8. Välimaa AL, Mäkinen S, Mattila P, et al. Fish and fish side streams are valuable sources of high-value components. *Food Qual Saf*. 2019;3(4):209-226. doi:10.1093/fqsafe/fyz024
9. Rottmann RW, Durborow R. The Role of Stress in Fish Disease. *South Reg Aquac Cent*. 1992;(474).
10. Dadgostar P. Antimicrobial resistance: implications and costs. *Infect Drug Resist*. 2019;12:3903-3910. doi:10.2147/IDR.S234610
11. Aoki W, Ueda M. Characterization of antimicrobial peptides toward the development of novel antibiotics. *Pharmaceuticals*. 2013;6(8):1055-1081. doi:10.3390/ph6081055
12. Kumari S, Tyor AK, Bhatnagar A. Evaluation of the antibacterial activity of skin mucus of three carp species. *Int Aquat Res*. 2019;11(3):225-239. doi:10.1007/s40071-019-0231-z
13. Tyor AK, Kumari S. Biochemical characterization and antibacterial properties of fish skin mucus of fresh water fish, *Hypophthalmichthys nobilis*. *Int J Pharm Pharm Sci*. 2016;8(6):6-10.
14. Loganathan K, Muniyan M, Prakash AA, Raja PS, Prakash M. Studies on the role of mucus from *clarias batrachus* (Linn) against studies on the role of mucus from *clarias batrachus* (Linn) against selected microbes. *Int J Pharm Appl ISSN*. 2014;2(3):202-206.
15. Elavarasi K, Ranjini S, Rajagopal T, Rameshkumar G, Ponmanickam P. Bactericidal proteins of skin mucus and skin extracts from fresh water fishes, *Clarias batrachus* and *Tilapia mossambicus*. *Thai J Pharm Sci*. 2013;37(4):194-200.
16. Lirio GAC, De Leon JAA, Villafuerte AG. Antimicrobial activity of epidermal mucus from top aquaculture fish species against medically-important pathogens. *Walailak J Sci Technol*. 2019;16(5):329-340. doi:10.14456/vol16iss5ppCorrectedProof
17. Bhatnagar A, Kumari S, Tyor AK. Assessment of bactericidal role of epidermal mucus of *Heteropneustes fossilis* and *Clarias batrachus* (Asian cat fishes) against pathogenic microbial strains. *Aquac Fish*. October 2021. doi:10.1016/j.AAF.2021.08.010
18. Ramesh B. Assessment of Antimicrobial peptides from mucus of fish. *IntJCurrBiotechnol*. 2013;1(1):5-8.
19. Balasubramanian S, Baby RP, Arul PA, Prakash M, Senthilraja P, Gunasekaran G. Antimicrobial properties of skin mucus from four freshwater cultivable Fishes (*Catla catla*, *Hypophthalmichthys molitrix*, *Labeo rohita* and *Tenopharyngodon idella*). *African J Microbiol Res*. 2012;6(24):5110-5120. doi:10.5897/AJMR11.532
20. Adel M, Safari R, Soltanian S, Zorriehzaha MJ, Esteban MÁ. Antimicrobial activity and enzymes on skin mucus from male and female Caspian kutum (*Rutilus frisii kutum* Kamensky, 1901) specimens. *Slov Vet Res*. 2018;55(4):235-243. doi:10.26873/svr-440-2018
21. Mahadevan G, Mohan K, Vinoth J, Ravi V. Biotic potential of mucus extracts of giant mudskipper *Periophthalmodon schlosseri* (Pallas, 1770) from Pichavaram, southeast coast of India. *J Basic Appl Zool*. 2019;80(1). doi:10.1186/s41936-019-0084-4
22. Al-Rasheed A, Handool KO, Garba B, et al. Crude extracts of epidermal mucus and epidermis of climbing perch *Anabas testudineus* and its antibacterial and hemolytic activities. *Egypt J Aquat Res*. 2018;44(2):125-129. doi:10.1016/j.ejar.2018.06.002
23. Okella H, Ikiriza H, Ochwo S, et al. Identification of antimicrobial peptides isolated from the skin mucus of African Catfish, *Clarias gariepinus* (Burchell, 1822). *Front Microbiol*. 2021;12(December). doi:10.3389/fmicb.2021.794631
24. Rana S, Bajaj R, Mann B. Characterization of antimicrobial and antioxidative peptides synthesized by *L. rhamnosus* C6 fermentation of milk. *Int J Pept Res Ther*. 2018;24(2):309-321. doi:10.1007/s10989-017-9616-2
25. Laemmli U. Cleavage of structural protein during the assembly of the head of bacteriophage T4. *Nature*. 1970;227:680-685.
26. Wang X, Li X, Li Y. A modified Coomassie Brilliant Blue staining method at nanogram sensitivity compatible with proteomic analysis. *Biotechnol Lett*. 2007;29(10):1599-1603. doi:10.1007/s10529-007-9425-3
27. Zhao Z, Liu J, Weir MD, et al. Human periodontal ligament stem cells on calcium phosphate scaffold delivering platelet lysate to enhance bone regeneration. *RSC Adv*. 2019;9(70):41161-41172. doi:10.1039/c9ra08336g
28. Lee HJ, Choi GJ, Cho KY. Correlation of lipid peroxidation in *Botrytis cinerea* caused by dicarboximide fungicides with their fungicidal activity. *J Agric Food Chem*. 1998;46(2):737-741. doi:10.1021/jf970501c
29. Lin MC, Hui CF, Chen JY, Wu JL. Truncated antimicrobial peptides from marine organisms retain anticancer activity and antibacterial activity against multidrug-resistant *Staphylococcus aureus*. *Peptides*. 2013;44:139-148. doi:10.1016/j.peptides.2013.04.004
30. Gao W, Xing L, Qu P, et al. Identification of a novel cathelicidin antimicrobial peptide from ducks and determination of its functional activity and antibacterial mechanism. *Sci Rep*. 2015;5(July):1-12. doi:10.1038/srep17260
31. Hancock RE. Cationic peptides: Effectors in innate immunity and novel antimicrobials. *Lancet Infect Dis*. 2001;1(3):156-164. doi:10.1016/S1473-3099(01)00092-5
32. Sun C, Gu L, Hussain MA, et al. Characterization of the bioactivity and mechanism of bactenecin derivatives against food-pathogens. *Front Microbiol*. 2019;10(November):1-13. doi:10.3389/fmicb.2019.02593
33. Burton MG, Huang QM, Hossain MA, Wade JD, Clayto

- nAHA, Gee ML. Longtimescale interaction dynamics between a model antimicrobial peptide and giant unilamellar vesicles. *Langmuir*. 2013;29(47):14613-14621. doi:10.1021/la403083m
34. Bahar AA, Ren D. Antimicrobial peptides. *Pharmaceuticals*. 2013;6(12):1543-1575. doi:10.3390/ph6121543
35. Hollmann A, Martinez M, Maturana P, Semorile LC, Maffia PC. Antimicrobial peptides: interaction with model and biological membranes and synergism with chemical antibiotics. *Front Chem*. 2018;6(JUN):1-13. doi:10.3389/fchem.2018.00204
36. Epand RM, Vogel HJ. Diversity of antimicrobial peptides and their mechanisms of action. *Biochem Biophys Acta*. 1999;1462(1999):11-28.
37. Oren Z, Shai Y. Mode of action of linear amphipathic α -helical antimicrobial peptides. *Biopolymers*. 1998;47(6):451-463. doi:10.1002/(SICI)1097-0282(1998)47:6<451::AID-BIP4>3.0.CO;2-F
38. Matsuzaki K, Murase O, Fujii N, Miyajima K. Translocation of a channel-forming antimicrobial peptide, magainin 2, across lipid bilayers by forming a pore. *Biochemistry*. 1995;34(19):6521-6526. doi:0006-2960/95/0434-6521\$09.00/0
39. Lohner K, Prossnigg F. Biological activity and structural aspects of PGLa interaction with membrane mimetic systems. *Biochim Biophys Acta - Biomembr*. 2009;1788(8):1656-1666. doi:10.1016/j.bbmem.2009.05.012
40. Jiao K, Gao J, Zhou T, et al. Isolation and purification of a novel antimicrobial peptide from *Porphyra yezoensis*. *J Food Biochem*. 2019;43(7):1-9. doi:10.1111/jfbc.12864
41. Sebastian M, Ammerman JW. The alkaline phosphatase PhoX is more widely distributed in marine bacteria than the classical PhoA. *ISME J*. 2009;3(5):563-572. doi:10.1038/ismej.2009.10
42. Chen H, Yang H, Weir MD, et al. An antibacterial and injectable calcium phosphate scaffold delivering human periodontal ligament stem cells for bone tissue engineering. *RSC Adv*. 2020;10(66):40157-40170. doi:10.1039/d0ra06873j
43. Guo F, Chen Q, Liang Q, et al. Antimicrobial activity and proposed action mechanism of linalool against *Pseudomonas fluorescens*. *Front Microbiol*. 2021;12(January):1-11. doi:10.3389/fmicb.2021.562094
44. Xiao XN, Wang F, Yuan YT, Liu J, Liu YZ, Yi X. Antibacterial activity and mode of action of dihydromyricetin from *ampelopsis grossedentata* leaves against food-borne bacteria. *Molecules*. 2019;24(15). doi:10.3390/molecules24152831
45. Zhang X, Ren C, Hu F, et al. Detection of bacterial alkaline phosphatase activity by enzymatic in situ self-assembly of the AIEgen-peptide conjugate. *Anal Chem*. 2020;92(7):5185-5190. doi:10.1021/acs.analchem.9b05704
46. Robert É, Fillion M, Otis F, Voyer N, Auger M. Understanding how the antimicrobial peptide thanatin interacts with the lipid bilayer of cell walls using model membranes. *Biophys J*. 2014;106(2):85a. doi:10.1016/j.bpj.2013.11.546
47. Green AA, Weech AA, Michaelis L. Studies on permeability of membranes: vii. Conductivity of electrolytes within the membrane. *J Gen Physiol*. 1929;12(3):487-493. doi:10.1085/jgp.12.3.487
48. Wu Y, Bai J, Zhong K, et al. Antibacterial activity and membrane-disruptive mechanism of 3-p-trans-coumaroyl-2-hydroxyquinic acid, a novel phenolic compound from pine needles of *Cedrus deodara*, against *Staphylococcus aureus*. *Molecules*. 2016;21(8). doi:10.3390/molecules21081084
49. Molina-Guijarro JM, Macías Á, García C, et al. Irvallec inserts into the plasma membrane causing rapid loss of integrity and necrotic cell death in tumor cells. *PLoS One*. 2011;6(4). doi:10.1371/journal.pone.0019042
50. Ma B, Guo Y, Fu X, Jin Y. Identification and antimicrobial mechanisms of a novel peptide derived from egg white ovotransferrin hydrolysates. *Food Sci Technol*. 2020;131:109720. doi:10.1016/j.lwt.2020.109720
51. Sawai J, Yoshikawa T. Quantitative evaluation of antifungal activity of metallic oxide powders (MgO, CaO and ZnO) by an indirect conductimetric assay. *J Appl Microbiol*. 2004;96(4):803-809. doi:10.1111/j.1365-2672.2004.02234.x
52. Zhang J, Ye KP, Zhang X, Pan DD, Sun YY, Cao JX. Antibacterial activity and mechanism of action of black pepper essential oil on meat-borne *Escherichia coli*. *Front Microbiol*. 2017;7(JAN):1-10. doi:10.3389/fmicb.2016.02094
53. Sawai J, Doi R, Maekawa Y, Yoshikawa T, Kojima H. Short communication: Indirect conductimetric assay of antibacterial activities. *J Ind Microbiol Biotechnol*. 2002;29(5):296-298. doi:10.1038/sj.jim.7000314
54. Schmidtchen A, Pasupuleti M, Malmsten M. Effect of hydrophobic modifications in antimicrobial peptides. *Adv Colloid Interface Sci*. 2014;205:265-274. doi:10.1016/j.cis.2013.06.009
55. Conlon JM, Mechkarska M, Lukic ML, Flatt PR. Potential therapeutic applications of multifunctional host-defense peptides from frog skin as anti-cancer, anti-viral, immunomodulatory, and anti-diabetic agents. *Peptides*. 2014;57:67-77. doi:10.1016/j.peptides.2014.04.019
56. Shilpakala SR, Mohan KVK, Atreya CD. A peptide derived from phage display library exhibits antibacterial activity against *E. coli* and *Pseudomonas aeruginosa*. *PLoS One*. 2013;8(2):1-11. doi:10.1371/journal.pone.0056081
57. Jothi GEG, Deivasigamani B, Priyadarshini P, Rajasekar T. Haemolytic and antimicrobial efficacy of the epidermal mucus of the striped haemolytic and antimicrobial efficacy of the epidermal mucus of the striped dwarf catfish, *Mystus vittatus* (Bloch 1794) from Vellar Estuary, Parangipettai. *Pharm Biotechnol Microbiol*. 2013;2014(1):1-6.
58. Manivasagan P, Annamalai N, Ashokkumar S, Sampathkumar P. Studies on the proteinaceous gel secretion from the skin of the catfish, *Arius maculatus* (Thunberg, 1792). *African J Biotechnol*. 2009;8(24):7125-7129. doi:10.4314/ajb.v8i24.68807
59. Raja K, Jayakumar T, Sahayanathan GJ, et al. Evaluation of anticancer, antibacterial and haemolytic activities of cru-

- de mucus from marine catfish *Tachysurus Dussumieri*. *Int J Pharma Bio Sci.* 2020;10(2):38-45. doi:10.22376/ijpbs/lpr.2020.10.2.138-45
60. Maillaud C, Hoang-Oppermann T, Hoang-Oppermann V, Rigot H, Girardot S, Nour M. Is stonefish *Synanceia verrucosa* envenomation potentially lethal? *Toxicon.* 2020;184:78-82. doi:10.1016/j.toxicon.2020.05.019
61. Robertson A, Garcia AC, Flores Quintana HA, et al. Invasive lionfish (*Pterois volitans*): A potential human health threat for ciguatera fish poisoning in tropical waters. *Mar Drugs.* 2014;12(1):88-97. doi:10.3390/md12010088
62. Cohen AS, Olek AJ. An extract of lionfish (*Pterois volitans*) spine tissue contains acetylcholine and a toxin that affects neuromuscular transmission. *Toxicon.* 1989;27(12):1367-1376. doi:10.1016/0041-0101(89)90068-8
63. Andrich F, Carnielli JBT, Cassoli JS, et al. A potent vasoactive cytotoxin isolated from *Scorpaena plumieri* scorpionfish venom. *Toxicon.* 2010;56(4):487-496. doi:10.1016/j.toxicon.2010.05.003
64. Go HJ, Kim CH, Park JB, et al. Biochemical and molecular identification of a novel hepcidin type 2-like antimicrobial peptide in the skin mucus of the pufferfish *Takifugu pardalis*. *Fish Shellfish Immunol.* 2019;93:683-693. doi:10.1016/j.fsi.2019.08.017
65. Wood SJ, Park YA, Kanneganti NP, et al. Modified cysteine-deleted tachyplesin (CDT) analogs as linear antimicrobial peptides: Influence of chain length, positive charge, and hydrophobicity on antimicrobial and hemolytic activity. *Int J Pept Res Ther.* 2014;20(4):519-530. doi:10.1007/s10989-014-9419-7

Peer Reviewed**Competing Interests:** None declared.**Funding:** This study was supported by the World Bank and Government of Uganda through PharmBiotechnology and Traditional Medicine (PHARMBIOTRAC) Incubation Center and CamTech Uganda, Mbarara University of Science and Technology (ACE-INCUB/2020/07), Mbarara, Uganda together with the International Foundation for Science (A_6226-1), Stockholm, Sweden, through grants to Hedmon Okella**Received:** 28 October 2021; **Accepted:** 26 November 2021**Cite this article as** Okella H, Ajayi OC, Ikiriza H, Mtewa GA, Kaggwa B, Aber J, Kato DC, Engeu OP. Bacterial Cell Envelope Lysis and Hemotoxicity of Peptides Previously Isolated From African Catfish, *Clarias gariepinus*. *East Afr Sci J.* 2022;4(1):93-100. <https://doi.org/10.24248/easci.v4i1.66>

© Okella et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are properly cited. To view a copy of the license, visit <http://creativecommons.org/licenses/by/4.0/>. When linking to this article, please use the following permanent link: <https://doi.org/10.24248/easci.v4i1.66>
