

Traumatic Brain Injury in Africa: The Republic of Malawi as an example.

T. Kapapa

Neurochirurgische Klinik, Universitätsklinikum Ulm, Albert-Einstein-Allee 23,
89081 Ulm, Germany

Correspondence to: Dr. med. Thomas Kapapa, Email: <thomas.kapapa@hotmail.de>

Background: *Due to the high rate of mortality and morbidity, traumatic brain injury is of great significance throughout the world. Preventive concepts have managed to considerably reduce the incidence in many countries in the northern hemisphere, but such an approach either does not exist or has failed in the developing nations of the southern hemisphere, where in some is a lack of data on the current incidence of traumatic brain injury. This is a review of the literature reflecting the current status of traumatic brain injuries in Malawi.*

Methods: *Issues of epidemiology as well as actuality of the data have been addressed in a PubMed and Medline review for the Republic of Malawi.*

Results: *There are very good studies about injuries in Malawi. However, they are not recent in a developing society and country. Traumatic brain injury as a neurosurgical and neurocranial problem in Malawi is underrepresented in the recent literature.*

Conclusion: *There is an urgent need for epidemiological studies of neurological and neurosurgical diseases including traumatic brain injuries to forward structured development in Malawi.*

Key words: Medical education; Public health; Africa; Neurosurgery; TBI

Introduction

Traumatic brain injury is a global problem, both medically and socially. During times of peace and war, traumatic brain injuries are steadily increasing in number¹. Traumatic brain injury refers to damage involving the head, including the brain and/or its linings, with neurological dysfunction². In recent decades, considerable advances have been made in the management of patients suffering a traumatic brain injury. In 1996 and 2007, the Brain Trauma Foundation published its first ever guidelines on the management of traumatic brain injury^{3,4}. The introduction and standardization of treatments for traumatic brain injury has resulted in a significant improvement in therapeutic outcomes as concerns mortality, morbidity, functional outcomes, treatment duration and costs⁵⁻⁸. Nevertheless, there are still considerable differences across the globe in terms of how traumatic brain injury is treated. Such differences depend on the one hand on medical technology and equipment and on the other hand on the medical expertise available.

The relevance of diagnostic decisions and adequate therapy for traumatic brain injury is evident from the wide-ranging outcomes. There may be short-term and acute effects, such as headache, impaired consciousness and focal neurological deficits, as well as long-term effects such as impaired cognitive performance and delayed development, fatigue, drowsiness, nausea, emotional disorders, e.g. changes in character and sleep disorders⁹. The objective of treatment for traumatic brain injury must be to prevent long-term morbidity, particularly in developing nations. Traumatic brain injury is still one of the great challenges to medical science, especially in developing countries such as in Africa¹⁰.

This review aims to address traumatic brain injury and its management in a third-world country as Malawi. Further aim was to confront the medical and social impact of traumatic brain injury in this fast economical and medical developing country and its representation in the recent medical literature. Relevant literature from the last 15 years (MedLine, PubMed) has been reviewed and summarized.

Malawi is one of the smallest, poorest, and most densely populated nations in Sub-Saharan Africa. It has a surface area of roughly 120,000 km² and a population of about 15,263,000. Current life expectancy at birth is 51 years in women and 44 years in men. Infant mortality in children under 5 years of age is 110 per 1,000 live births. The mortality rate in the population aged between 15 and 60 years is 496 in women and 691 in men per 1,000 inhabitants. Gross national income per capita is 619 EUR¹¹. 65% of the population is living below the poverty line, surviving on less than 76 eurocents per day¹². In principle, everyone has the right to public healthcare provision.

Malawi's healthcare system encompasses 21 district hospitals, each located in its own regional health district. In addition, there are four central hospitals in four larger municipal districts. Together, the state-run hospitals manage almost 60% of the nation's healthcare needs. Further health services are provided by church missions and independent hospitals, thereby managing the remaining 40% of the healthcare needed¹³.

There are roughly 23 trained surgeons in Malawi covering the entire surgical spectrum. They are not stationed at the district hospitals, however. Most district hospitals have one doctor who, after completing his medical studies and basic practical training, is assigned to one of the said facilities. He assumes responsibility for the hospital's medical and commercial management. Clinical officers with four years of paramedical training support the work of this District Health Officer. Periodically, surgeons from the central hospitals visit the district hospitals to provide assistance with the clinical workload¹³. At present, two neurosurgeons are working in Malawi.

Epidemiology of injuries and surgical statistics

A study of the epidemiology of injuries conducted at Kamuzu Central Hospital (February to June 2008) revealed that of the 1474 patients counted, men were in the majority (75.7%). The age distribution was described as bimodal, with a peak at the age of < 5 years and between 26 and 30 years (Figure 1).




Figure 1. Age distribution of patients recorded in the Malawi Trauma Registry¹⁴.

A study at Kamuzu Central Hospital in 2008 found that road traffic accidents (43%) were the most common cause for the injuries (N=1474). Passenger cars were most commonly involved in such accidents (19%), followed by attacks and assaults (24%), then falls (13.5%). Traumatic

brain injuries were reported at a rate of 7.5%. The most frequent means of reaching the hospital were private vehicle (43.8%), ambulance (15.4%), walking (14.5%) and public transport (7.8%). Depending on the method used for reaching the hospital, the pre-hospital phase varied between 3 hours and 21 minutes¹⁴.

Traumatic brain injury and its management in Malawi

Traumatic brain injury is one of the leading causes of death and morbidity worldwide¹⁵. Just as in Europe, road traffic accidents are one of the main causes for traumatic brain injury in Malawi². When compared on an international level, Malawi has fewer roads and means of transportation (*compared at a statistical level of approx. 100,000 vehicles*), but a very high number of road traffic accidents. In 1995, 82.9 accidents per 1000 vehicles were reported. Between 1987 and 1995, 126.8 to 138.7 deaths were reported following an accident per 10,000 vehicles. 55.6 to 73.5 individuals died at the site of the accident per 1000 vehicles¹⁶. According to the German Federal Statistical Office, 761 individuals were killed by a road traffic accident in March 2012 (80200)¹⁷. Roughly 25% of traumatic brain injuries in Germany are caused by road traffic accidents¹⁸. More recent figures are not available for Malawi. However, older figures reveal that there were about 5 acute traumatic brain injuries per 100 hospital admissions at the Queen Elizabeth Central Hospital between July and December 1995. In 1994, traumatic brain injury was the main cause of death amongst emergency admissions at the surgical department, ahead of burns or HIV-associated Kaposi's sarcoma¹⁹.

If the frequencies of surgical interventions in Malawi are compared, it becomes clear that neurocranial operations are extremely rare despite the fact that the number of traumatic brain injuries is high. Boreholes (no other neurosurgical procedures like craniotomy) are the most uncommonly procedure in Malawi (Number of operations = 48 696; 25 053 in district hospitals and 23 643 in central hospitals)¹³.

Even when neurosurgery itself is not the major medical issue in Malawi - beside greater problems like HIV, lung diseases or mortality in children - the following epidemiological questions thus arise from the discussion above, also with a view to improving a small medical speciality and the relevant medical care. What are the current figures and causes for traumatic brain injuries in the municipal and rural areas of Malawi? How are traumatic brain injuries distributed in Malawi in relation to age, sex and severity? What structure exists for managing traumatic brain injuries in Malawi (admissions, surgical treatment, and computed tomography)? What are the (treatment) outcomes following traumatic brain injury in Malawi?

The current incidence of traumatic brain injury in Malawi is unknown. There is an urgent need for new epidemiological research. With the growing prosperity and advancing logistical development of a country, the short and long-term effects of traumatic brain injury may not be underestimated.

References

1. Risdall JE, Menon DK. Traumatic brain injury. *Philos Trans R Soc Lond B Biol Sci.* 2011 Jan 27;366(1562):241-50. PubMed PMID: 21149359. Pubmed Central PMCID: 3013429. Epub 2010/12/15. eng.
2. Röhrer S, Woischneck D, Kapapa T. Epidemiological data on Head Trauma in Europe: Age and Sex. *European Orthopaedics and Traumatology.* 2012;3(2):127-134.
3. Bullock MR, Povlishock JT. Guidelines for the management of severe traumatic brain injury. Editor's Commentary. *J Neurotrauma.* 2007;24 Suppl 1:2 p preceding S1. PubMed PMID: 17511557. Epub 2007/05/22. eng.

4. Bullock R, Chesnut RM, Clifton G, Ghajar J, Marion DW, Narayan RK, et al. Guidelines for the management of severe head injury. Brain Trauma Foundation. Eur J Emerg Med. 1996 Jun;3(2):109-27. PubMed PMID: 9028756. Epub 1996/06/01. eng.
5. Arabi YM, Haddad S, Tamim HM, Al-Dawood A, Al-Qahtani S, Ferayan A, et al. Mortality reduction after implementing a clinical practice guidelines-based management protocol for severe traumatic brain injury. J Crit Care. 2010 Jun;25(2):190-5. PubMed PMID: 19592201. Epub 2009/07/14. eng.
6. Fakhry SM, Trask AL, Waller MA, Watts DD. Management of brain-injured patients by an evidence-based medicine protocol improves outcomes and decreases hospital charges. J Trauma. 2004 Mar;56(3):492-9; discussion 9-500. PubMed PMID: 15128118. Epub 2004/05/07. eng.
7. Hesdorffer DC, Ghajar J, Iacono L. Predictors of compliance with the evidence-based guidelines for traumatic brain injury care: a survey of United States trauma centers. J Trauma. 2002 Jun;52(6):1202-9. PubMed PMID: 12045655. Epub 2002/06/05. eng.
8. Vukic M, Negovetic L, Kovac D, Ghajar J, Glavic Z, Gopcevic A. The effect of implementation of guidelines for the management of severe head injury on patient treatment and outcome. Acta Neurochir (Wien). 1999;141(11):1203-8. PubMed PMID: 10592121. Epub 1999/12/11. eng.
9. Almasi SJ, Wilson JJ. An update on the diagnosis and management of concussion. WMJ. 2012 Feb;111(1):21-7; quiz 8. PubMed PMID: 22533212. Epub 2012/04/27. eng.
10. Puvanachandra P, Hyder AA. Traumatic brain injury in Latin America and the Caribbean: a call for research. Salud Publica Mex. 2008;50 Suppl 1:S3-5. PubMed PMID: 18373006. Epub 2008/05/09. eng.
11. World-Health-Organization. Country Statistics Malawi Lilongwe: World Health Organization; 2012 [cited 2012 07. May]. Available from: whomalawi@mw.afro.who.int.
12. van Amelsfoort JJ, van Leeuwen PA, Jiskoot P, Ratsma YE. Surgery in Malawi--the training of clinical officers. Trop Doct. 2010 Apr;40(2):74-6. PubMed PMID: 20305097. Epub 2010/03/23. eng.
13. Lavy C, Tindall A, Steinlechner C, Mkandawire N, Chimangeni S. Surgery in Malawi - a national survey of activity in rural and urban hospitals. Ann R Coll Surg Engl. 2007 Oct;89(7):722-4. PubMed PMID: 17959015. Pubmed Central PMCID: 2121267. Epub 2007/10/26. eng.
14. Samuel JC, Akinkuotu A, Villaveces A, Charles AG, Lee CN, Hoffman IF, et al. Epidemiology of injuries at a tertiary care center in Malawi. World J Surg. 2009 Sep;33(9):1836-41. PubMed PMID: 19597877. Pubmed Central PMCID: 3290404. Epub 2009/07/15. eng.
15. Maegele M, Engel D, Bouillon B, Lefering R, Fach H, Raum M, et al. Incidence and outcome of traumatic brain injury in an urban area in Western Europe over 10 years. Eur Surg Res. 2007;39(6):372-9. PubMed PMID: 17690556. Epub 2007/08/11. eng.
16. Olukoga A. Trends in road traffic crashes, casualties and fatalities in Malawi. Trop Doct. 2007 Jan;37(1):24-8. PubMed PMID: 17326882. Epub 2007/03/01. eng.
17. Statistisches-Bundesamt-der-Bundesrepublik-Deutschland. Gesundheitsberichterstattung des Bundes Bonn: Bundesministerium für Gesundheit; 2012 [cited 2012 28.05.2012].
18. Beynon C, Unterberg AW. Schweres Schädel-Hirn-Trauma. Unfallchirurg. 2011;114:713-23.
19. Adeloye A, Ssembatya-Lule GC. Aetiological and epidemiological aspects of acute head injury in Malawi. East Afr Med J. 1997 Dec;74(12):822-8. PubMed PMID: 9557432. Epub 1998/04/29. eng.