

PLANT LIFE FORM CLASSIFICATION AND DISTRIBUTION AT SUBA SEBETA FOREST, ETHIOPIA

Mistire Yifru¹, Tamrat Bekele^{2,*}, Ensermu Kelbessa² and Tigist Wondimu²

ABSTRACT: A study focusing on altitudinal distribution of life forms of vascular plants was conducted in Suba Sebeta Forest, located 40 km west of Addis Ababa in Oromia National Regional State. The objective of the study was to document the life form composition of Suba Sebeta Forest based on life form spectrum. In the study, a systematic sampling method was employed to collect vegetation data from 48 (20 m x 20 m) plots. These entire quadrats included 16 (5 m x 5 m each) subplots to incorporate all life form types. Vegetation classification was done based on Raunkiaer's major life form groups. R-Software (Version 2.15.1) Package was used to classify the life forms. The data were collected along altitudinal gradients, ranging from 2400 to 2870 m a.s.l. A total of 135 species belonging to 67 families were recorded from the study area; the results showed that Asteraceae was the dominant family accounting for 15.6% of the total record. Analysis of life form spectrum showed Phanerophytes as the dominant life form contributing 48 species (35.6%) followed by Hemicryptophytes (34 species, 25.18%), Chamaephytes (22 species, 16.30%), Therophytes (20 species, 14.81%) and Geophytes (11 species, 8.15%). Correspondence analysis between life form and environmental gradients showed no significant relationship between life forms and soil pH while altitudinal difference affects the life form distribution. These life form variabilities along altitudinal gradient lay the foundation for further studies for monitoring of changes relating to climate change.

Key words/phrases: Life form, Life form spectrum, Raunkiaer, Suba Sebeta Forest.

INTRODUCTION

The vegetation cover of an area has a definite structure and composition developed as a result of long-term interaction of biotic and abiotic factors (Peters, 1996). Several quantitative measures are employed to describe the structure of plant communities with much ecological precision, qualitative characters such as species richness, life form spectrum, and vertical disposition of species. Raunkiaer (1934) described communities of different climatic zones or phytoclimatic zones of the earth on the basis of life form spectrum or biological spectrum. Any change in the life form composition

¹ Kotebe University College, Addis Ababa, Ethiopia. E-mail: ymistire@yahoo.com

² Department of Plant Biology and Biodiversity Management, Addis Ababa University, P.O. Box 3434, Addis Ababa, Ethiopia. E-mail: tambek07@yahoo.com; ensermuk2002@yahoo.co.uk; twtigitw@gmail.com

away from its phytoclimatic zone is considered as an indicator of alteration in vegetation either due to biotic or edaphic factors or both. However, recently, the natural structural complexities of plant communities are being affected mostly due to reckless anthropogenic activities.

Excessive utilization of forest resources and over grazing have resulted in changes in the life form composition of plant communities (Reddy *et al.*, 2002; Verma and Shukla, 1993). Life form classification is a powerful instrument employed for several purposes like land management and climate control. It also provides a way of summarizing our knowledge of understanding on life form pattern. Life form composition of vegetation to a certain extent is an indicator of the climate, and is also helpful to compare geographical distribution and to identify widely distributed plant communities (Muhammad, 2012). Plant life forms differ in every climate zone on the basis of altitude. Long-lived life forms, such as Phanerophytes, dominate in tropical zones and Hemicryptophytes dominate in arctic or alpine zones, whereas short-lived life forms such as Therophytes are most frequent in deserts. Raunkiaer (1934) reported the existence of three types of life forms with respect to the different climate zones of the earth: Phanerophytic in tropics, Therophytic in deserts and Hemicryptophytic in cold temperate zone. Cain and Castro (1959) and Shimwell (1971) also reported Hemicryptophytes as indicators of temperate zone climate while Therophytes and Geophytes are indicators of desert climate and Mediterranean climate, respectively. Vegetation profile of the component species indicates the phyto-climatic condition of an area and is employed commonly in community structural description through profile diagrams (Unwin, 1989; Ashton and Hall, 1992; Chen, 1995; Pignatti, 1995; Sahunalu and Dhanmanonda, 1995; Visalakshi, 1995).

Similarities of plants in structure and form indicate their adaptive mechanism to a given area. Whenever such similar plants grow close together, they become strong competitors since they are adapted to use the environmental resources in the same manner. According to Mueller-Dombois and Ellenberg (1974), plants of the same life form growing together are likely to compete directly for the same space. Variation in life forms along altitudinal gradients has been used for a better interpretation of vegetation and species richness patterns in relation to environmental gradients. Life form diversity creates ecological opportunity and this positive feedback culminates in the high biotic diversity that is found in tropical forests. Variation of species composition along altitudinal gradient also correlates with ecological plasticity. The distribution with altitude of

every plant species gives an idea of its ecological plasticity. The life forms of plants include species, which have similarities in the complex of ecological conditions and thus are reflections of their habitat characteristics (Angelova and Tashev, 2005).

Several factors such as environment, periodic phenomena, and community development influence life forms. On the other hand, internal ecosystem functioning can be influenced by an already established life form. Solbrig (1993) described life forms as single character based functional groups, yet even at this level there are some similarities in life history and resource use that lend coherence to the categories. This is perhaps expected if life forms are the result of evolutionary forces that lead to ecological convergence (Bocher, 1977).

The vegetation resources of Ethiopia have been studied by different scholars (Logan, 1946; Pichi-Sermolli, 1957; von Breitenbach, 1962; 1963; Westphal, 1975; Chaffey, 1979; Friis *et al.*, 1982; Friis, 1986; Tewolde Berhan Gebre Egziabher, 1988; Friis and Mesfin Tadesse, 1990; EFAP, 1994; Teshome Soromessa and Sebsebe Demissew, 2002). Although, many studies have been done on the vegetation classification and different ecological parameters, researches that relate to plant life form are scanty if not none at all in Ethiopia. However, the current global climate change scenario necessitates study of life forms in every climatic zone. This study focuses on vegetation classification based on Raunkiaer life form spectrum along an altitudinal gradient in the Suba Sebeta Forest. In addition, the study aims at documenting the life form composition of Suba Sebeta Forest based on biological life form spectrum for a better understanding of the vegetation characteristic.

MATERIALS AND METHODS

Description of the study area

Suba Sebeta Forest is located 40 km southwest of Addis Ababa. Geographically, it lies between 38°31' and 38°35' E and 9°89' and 9°00' N in Oromia National Regional State. The forest is found in West Shewa Zone and is part of the central plateau with altitudinal range of 2200-3385 m a.s.l. (Afework Bekele, 1994) (Fig. 1). The forest is found on the southwestern slopes of Mt. Wochacha, which is a massive dead volcano mountain with an elevation of 3385 m a.s.l. The mountain sides are generally steep with ravines cut by streams and rivers. The southern base of the mountain is at 2200 m and flanks the Becho plains. The Suba Sebeta Forest is one of the remnant patches of dry Afromontane forest in Ethiopia. This type of forest

together with grassland and woodland once had formed a vegetation mosaic across the Ethiopian plateau. The forests and woodlands occur on the better drained soils of the mountains and sides of the valleys while the grassland occupies the heavy clay soils of the valley bottoms.




Fig. 1. Map of the study area.

Data collection

A reconnaissance survey of the vegetation was made in October 2013 in order to obtain an impression of the vegetation and topographic features. Actual field data were collected from October to November 2013. Vegetation and environmental data were collected in sample quadrats placed along transect lines, which were systematically laid. A total of 48 quadrats, at every 10 m altitudinal drop, were laid along transect lines between 2400–2870 m a.s.l. All woody plant species including trees and shrubs were recorded in 20 m × 20 m quadrats while herbaceous species were recorded in 16 sub quadrats (5 m × 5 m) that systematically cover the entire main quadrat.

Information on habit, habitat, vegetation type, the nature of perennial bodies was recorded so as to draw a life form spectrum, following the concept of Raunkiaer (1934). In the construction of life form spectra each species was assigned to a single life form.

Physiographic variables such as altitude and geographic coordinates were measured for each quadrat using GPS. Specimens of all vascular plant taxa were collected, pressed, dried and brought to the National Herbarium (ETH), Addis Ababa University, for identification. The specimens were dried in the dryer, kept in a deep freezer for 72 hours and identified referring to the volumes of Flora of Ethiopia and Eritrea as well as reference collections.

Soil sample

Soil samples were taken from five spots in each plot, from the four corners and from the centre. The weight of soil from the five spots was 250–300 g. Separate soil samples were taken from 0–15 cm depth of soil layers from each spot. The samples were mixed thoroughly, air dried and passed through a 2 mm mesh sieve to remove the stone pieces and the debris.

Data analysis

All the plant species recorded were grouped into different life forms based on criteria outlined by Raunkiaer (1934), and Mueller-Dombois and Ellenberg (1974). Also, a simple metrics was used to analyze proportions of life forms following Pharswan *et al.* (2010).

The percentage of life form was calculated using the following formula:

$$\% \text{ Life form} = \frac{\text{Number of species in any life form}}{\text{Total number of species of all life forms}} \times 100$$

Soil pH analysis

The combined soil sample was subjected to pH analysis which was performed in the eco-physiology laboratory of the Department of Plant Biology and Biodiversity Management, College of Natural and Computational Sciences, Addis Ababa University. Soil pH was measured by mixing 20 grams of soil sample and 20 ml of distilled water (1:1 ratio), and stirring for 30 minutes in a 100 ml beaker. The soil-water mixture was kept for 5 minutes and the reading was taken with a Jenway 3345 Digital pH meter.

RESULTS

Life form composition

A total of 135 vascular plant species were recorded in Suba Sebeta Forest belonging to five major life form groups. The percentage of species belonging to each life form category relative to the total number of species is presented in Fig. 2. The largest life form were Phanerophytes represented by 48 species (35.5%) followed by Hemicryptophytes (34 species, 25.2%), Chamaephytes (22 species, 16.3%), Therophytes (20 species, 14.8%) and Geophytes (11 species, 8.2%).

The 135 species recorded in the study belong to 67 families. The dominant families were Asteraceae (21 species, 15.6%), followed by Fabaceae (8 species, 5.9%), Lamiaceae (6 species, 4.4%), Apiaceae, Rubiaceae, Scrophularaceae each represented by 5 species each (3.7%).




Fig. 2. Number of species per life form in Suba Sebeta Forest.

The family Asteraceae (15.6%) is represented by all the five major life form groups (Table 1). The representative species in this family include *Tagetes minuta* – Therophyte, *Lactuca inermis* – Hemicryptophyte, *Sonchus bipontini* – Geophyte, *Solanecio gigas* – Chamaeophyte and *Vernonia wollastonii* – Phanerophyte. Fabaceae with 5.9% is the next diverse family represented by Therophyte – *Trifolium simensis*, Chamaeophyte – *Dolichos sericeus* and Phanerophyte – *Albizia schimperiana*. Hemichryptophytic and Geophytic life forms are not represented in Fabaceae.

Table 1. Dominant species in the five major life form groups and their percentage contribution in Suba Sebeta Forest.

Life form	Dominant species	%life form contribution	Growth form
Phanerophyte	<i>Bersama abyssinica</i>	43.8%	Tree
	<i>Dovyalis abyssinica</i>		Shrub
	<i>Dovyalis verrucosa</i>		Shrub
	<i>Jasminum abyssinicum</i>		Shrub
	<i>Juniperus procera</i>		Tree
	<i>Maytenus gracilipes</i>		Shrub
	<i>Maytenus undata</i>		Tree
	<i>Myrsine africana</i>		Shrub
	<i>Olea europaea L. subsp. cuspidata</i>		Tree
	<i>Olinia rochetiana</i>		Tree
	<i>Podocarpus falcatus</i>		Tree
	<i>Scolopia theifolia</i>		Shrub
	<i>Sideroxylon oxyacanthum</i>		Shrub
<i>Teclea nobilis</i>	Shrub		
Hemicryptophyte	<i>Adiantum poireti</i>	25%	Herb
	<i>Asplenium monanthes</i>		Herb
	<i>Cyperus tenuispica</i>		Herb
	<i>Geranium arabicum</i>		Herb
	<i>Kalanchoe petitiiana</i>		Herb
	<i>Mimulopsis olmsii</i>		Herb
	<i>Oplismenus shirtellus</i>		Herb
<i>Sanicula elata</i>	Herb		
Chamaephyte	<i>Achyranthes aspera</i>	15.6%	Herb
	<i>Hypoestes triflora</i>		Herb
	<i>Lobelia giberroa</i>		Herb
	<i>Mikaniopsis clematoides</i>		Herb
	<i>Solanecio gigas</i>		Herb
Therophyte	<i>Agrocharis melanantha</i>	12.5%	Herb
	<i>Cynoglossum coeruleum</i>		Herb
	<i>Galiuma parinoides</i>		Herb
	<i>Impatiens hochstteteri</i>		Herb
Geophyte	<i>Cynoglossum amplifolium</i>	3.1%	Herb

Life form similarity among plots

The cluster analyses resulted in five clusters (Fig. 3). The data matrix contains 48 quadrats and 135 species.




Fig. 3. Dendrogram outputs of the life form data obtained from hierarchical cluster analysis. The plot code and arrangement of plot along the dendrogram based on their cluster group.

C1: (Plots 1, 8, 9, 10, 11, 12)

C2: (Plots 2, 3, 7, 41, 42)

C3: (Plots 4, 22, 24, 28, 29, 32, 33, 37, 38, 39, 40, 43, 47)

C4: (Plots 5, 6, 13, 14, 15, 16, 17, 18, 19, 20, 21, 23, 25, 27, 31, 34, 36, 46)

C5: (Plots 26, 30, 35, 44, 45, 48)

Relation between altitude and soil pH

A negative correlation between soil pH and altitude was obtained. The soil pH ranged between 5.6 and 7.3 (Fig. 4). The lower pH (i.e., 5.6) indicating acidic soil was recorded at a high altitude (i.e., 2870 m a.s.l.). The highest pH (i.e., 7.3) indicating a more or less neutral soil was recorded in soils from the lower altitude (i.e., 2570 m a.s.l.).




Fig. 4. Relation between altitude and soil pH.

Endemic plants and their life form in Suba Sebeta Forest

Endemic plant species that were collected from Suba Sebeta Forest and their life forms contribution in the study area is given in Table 2.

Table 2. List of endemic plant species and their life form in Suba Sebeta Forest.

No.	Species name	Family	Life form	Growth form
1	<i>Cirsium schimperi</i>	Asteraceae	Geophyte	Herb
2	<i>Conyza nana</i>	Asteraceae	Hemicryptophyte	Herb
3	<i>Jasminum stans</i>	Oleaceae	Phanerophyte	Shrub
4	<i>Laggerato mentosa</i>	Asteraceae	Chamaephyte	Shrub
5	<i>Maytenus addat</i>	Celastraceae	Phanerophyte	Tree
6	<i>Mikaniopsis clematoides</i>	Asteraceae	Chamaephyte	Climber
7	<i>Millettia ferruginea</i>	Fabaceae	Phanerophyte	Tree
8	<i>Satureja paradoxa</i>	Lamiaceae	Hemicryptophyte	Herb
9	<i>Solanecio gigas</i>	Asteraceae	Chamaephyte	Herb
10	<i>Vernonia leopoldii</i>	Asteraceae	Chamaephyte	Shrub

DISCUSSION

In this study, out of the 135 species, 48 belonged to the Phanerophytic life form. The dominant Phanerophyte life form includes the two important Gymnosperms, *Juniperus procera* and *Podocarpus falcatus* which are characteristic species of the dry Afromontane forests. Tree and shrubby angiosperms like *Olea europaea* subsp. *cuspidata*, *Olinia rochetiana*, *Myrsine africana* and *Rosa abyssinica* also belong to this life form. The Hemicryptophytes also have great contribution to the life form spectrum. The most probable factor to this life form composition must be the humid nature of Suba Sebata Forest. Most Hemicryptophytes are herbaceous in habit that includes *Thymus schimperi*, *Veronica simensis* and *Rubia cordifolia*.

The Chamaephytic life form in Suba Sebata Forest is represented by the shrub *Hypericum quartinianum* and the herb *Mikaniopsis clematoides*, which carpeted the ground at higher altitudes. Other herbs belonging to the same group include *Achyrosperrum schimperi*, *Hypoestes triflora* and *Solanecio gigas*. Chamaephytes are indicators of alpine vegetation (Muhammad, 2012). In the life form composition, Therophytes and Geophytes have great contributions comprising 20 and 11 species each, respectively. Therophytes are annuals that include *Agrocharis melantha*, *Dichrocephala chrysanthemifolia* and *Cynoglossum coeruleum* which reproduce through seeds and complete their life cycle in a short period of time. Geophytes are represented by *Cynoglossum amplifolium*, *Arisaema enneaphyllum* and *Scadoxus multiflorus*.

Asteraceae is the largest family and genus in terms of endemism contributing 6 species (60%) that belong to Chamaephytic, Hemicryptophytic and Geophytic life forms. The remaining four families contain one species each and altogether covered 40% of the overall endemism. From these four families, three of them belong to Phanerophytic life forms and one is Hemicryptophytic life form. There are also species that are endemic to the Flora of Ethiopia and Eritrea. These near endemic species include *Kalanchoe petitiana*, *Thymus schimperi* and *Veronica simensis* that occurred in Hemicryptophytic life form while *Plectocephalus varians*, *Sideroxylon oxyacanthum* and *Erucastrum pachypodium* are found in Geophytic, Phanerophytic and Therophytic life forms, respectively.

The life form of plant species is usually a constant character (Mueller-Dombois and Ellenberg, 1974) and life forms are species-specific. But some species show different life forms when growing under very different

environmental conditions. Plants of the same life form growing together are likely to compete directly for the same space. The anthropogenic factors are so pervasive that there is scarcely any tropical forest in the world in which human activities have not left their mark on life form diversity (Denslow and Padoch, 1988; Goldammer, 1992). In this study, *Prunus africana* showed two different life form strategies: a Chamaephytic and Phanerophytic type. This result indicates that *Prunus africana* developed two life forms in the same forest to cope up with environmental influence that may be caused by human activity. The human activities in this species include removing the bark of the tree for traditional medicine, using the tree for timber production, as source of material for building local houses, fuel wood and making charcoal. This explanation is strongly supported by the works of Legesse Negash (2002).

The highest similarities of life form cluster along altitude were recorded between Phanerophytic and Hemichryptophyte life form. These life forms were similar due to their presence under similar soil pH range and altitude. Soil pH and altitude were correlated negatively and as the altitude increased pH value decreased or became slightly acidic. Great life form cluster differences were created between clusters three and cluster one which were dominated with Phanerophytic and Chamaephytic life forms, respectively. Chamaephytic life forms show increasing trend in the higher altitudes while Phanerophytes have a decreasing trend in higher altitudes. From this result, Chamaephytic life forms are better adapted in slightly acidic soil than Phanerophytes. The second difference in distribution was also observed between the Therophytic species which are mostly confined to lower altitudes and the Chamaephytic life forms that are abundant at higher altitudes. The third one was between Therophytes and Hemichryptophytes, which may be due to altitude, soil moisture and the soil pH differences.

The diversity and distribution of life forms is usually correlated with climatic heterogeneity (Cowling *et al.*, 1994) decreasing with increasing altitude (Montana and Valiente-Banuet, 1998; Pavón *et al.*, 2000). Results of the present study indicate that altitude is the main factor influencing life form composition. Most species in the present study are ephemerals that complete their life cycle just in a short period of time. The predominance of annuals at low altitudes can partly be explained by soil disturbance (Klimes, 2003). The more intense or the more frequent the disturbance is, the lower the proportion of long-lived life forms.

The results from Klimes (2003) confirm that Chamaephytes show an increasing trend towards higher altitudes. The work of Muhammad (2012) concluded that Chamaephytes are generally the most common life form both in high altitudes and high latitudes. At higher altitude precipitation is high resulting in leaching of the upper surface, which leads to increasing of soil acidity. With increasing altitude, temperature will typically decrease and the corresponding precipitation will increase (Barry, 1981; Ineson *et al.*, 1998; Shen *et al.*, 2013). Plants absorb dissolved nutrients in the soil water and the nutrient solubility depends largely on the pH value. Hence, the availability of elements varies at different pH levels.

CONCLUSION

This study contributes basic data on life form composition and life form spectrum of vascular plants at Suba Sebeta Forest. In order to provide baseline information for subsequent studies, a total of 135 species were identified and assigned to 67 families. The family Asteraceae was the most dominant followed by Fabaceae and Lamiaceae, respectively. The study of biological life form spectrum more accurately describes the vegetation physiognomy, since each plant is counted by its life form. The life form spectrum was dominated with Phanerophyte (35.56%) followed by Hemicryptophytes (25.18%), Chamaephytes (16.30%), Therophytes (14.81%) and Geophyte (8.15%). Altitudinal variation played a great role in life form distribution and composition in the study area. At higher altitudes, Chamaephytic life form was dominant while Hemicryptophytes were well distributed. Cluster analysis of the vegetation data of the forest produced five communities each of which have its own indicator and dominant life form type. The distribution of these plant life form communities in the forest was influenced by environmental factors and biotic stresses which are derivatives of the life form of plant and composition. While it is believed that this study is a pioneer attempt, further elaborated studies to correlate the species distribution with chemical and physical characteristics of soils and other environmental influences should be undertaken.

ACKNOWLEDGEMENTS

The first author would like to thank Addis Ababa University and the sponsoring institute, Kotebe University College. The role of Oromia Forest and Wildlife Enterprise in giving permission to work in Suba Sebeta Forest is highly appreciated. Analytical help from Prof. Zerihun Woldu and Ashenafi Tilahun is acknowledged. Staff members of the Department of Plant Biology and Biodiversity Management and the National Herbarium

(ETH), and Ethiopian Mapping Agency and Ethiopian Meteorological Service Agency are acknowledged for their help in various ways.

REFERENCES

- Afework Bekele (1994). **The Ecology of *Praomys alpines* (Ruppell, 1842) in the Menagesha State Forest, Ethiopia**. Ph.D. Dissertation, Addis Ababa University, Addis Ababa.
- Angelova, K. and Tashev, A. (2005). Complex analysis of the life forms of flowering plants in Mount Chepan and their vertical ranges of spread in altitude. *Trakia J. Sci.* **3**(6): 32–35.
- Ashton, P.S. and Hall, P. (1992). Comparison of structure among mixed dipterocarp forest of north-western Borneo. *J. Ecol.* **80**: 459–481.
- Barry, R.G. (1981). **Mountain Weather and Climate**. Methuen, London.
- Bocher, T.W. (1977). Convergence as an evolutionary process. *Bot. J. Linn. Soc.* **75**: 1–19.
- Cain, S.A. and Castro, G.M.D. (1959). **Manual of Vegetation Analysis**. Harper and Brothers Publication, New York.
- Chaffey, D.R. (1979). South-west Ethiopia forest inventory project: An inventory of forest at Munesa and Shashemene Project. Report 29, Ministry of Overseas Development Land Resource Division, London.
- Chen, W.L. (1995). Subtropical montane deciduous forests in southern China. In: **Vegetation Science in Forestry**, pp. 317–323 (Peet, R.K., Masuzawa, T., Yamada, I., Fujiwara, K. and Maycock, P.F., eds.). Kluwer Academic Publishers, Dordrecht.
- Cowling, R.M., Esler, K.J., Midgley, G.F. and Honing, M.A. (1994). Plant functional diversity, species diversity and climate in arid and semi-arid southern Africa. *J. Arid Environ.* **27**: 141–158.
- Denslow, J.S. and Padoch, C. (1988). **People of the Tropical Rain forest**. University of California Press, Berkeley, California.
- EFAP (1994). **The Challenge for Development**. Ethiopian Forestry Action Program (EFAP), Addis Ababa.
- Friis, I. (1986). The forest vegetation of Ethiopia. *Acta Univ. Ups. Symb. Bot. Ups.* **26**: 31–47.
- Friis, I. and Mesfin Tadesse (1990). The evergreen forests of northeast tropical Africa. In: *Mitt. Inst. Allg. Bot. Hamburg* **23a**: 249–263.
- Friis, I., Rasmussen, F.N. and Vollesen, K. (1982). Studies in the flora and vegetation of Southwest Ethiopia. *Opera Bot.* **63**: 8–70.
- Goldammer, J.G. (1992). **Tropical forests in transition - Ecology of Natural and Anthropogenic Disturbance Processes**. Birkhauser Verlag, Basel.
- Ineson, P., Taylor, T., Harrison, A.F., Poskitt, J., Benham, D.G., Tipping, E. and Woof, C. (1998). Effects of climate change on nitrogen dynamics in upland soils. 1. A transplant approach. *Glob. Chang. Biol.* **4**: 143–152.
- Klimes, L. (2003). Life forms and clonality of vascular plants along an altitudinal gradient in E. Ladakh (NW Himalayas). *Basic Appl. Ecol.* **4**: 317–328.
- Legesse Negash (2002). Review of research advances in some selected African trees with special reference to Ethiopia. *Ethiop. J. Biol. Sci.* **1**(1): 81–126.
- Logan, W.E.M. (1946). An introduction to the forests of central and southern Ethiopia. Imperial Forestry Insitute, University of Oxford, Oxford.

- Montana, C. and Valiente-Banuet, A.C. (1998). Floristic and life-form diversity along an altitudinal gradient in an intertropical semiarid Mexican region. *Southwest. Nat.* **43**(1): 25–39.
- Mueller-Dombois, D. and Ellenberg, H. (1974). **Aims and Methods of Vegetation Analysis**. John Wiley and Sons, New York.
- Muhammad, S.A. (2012). Life form and leaf size spectra of vegetation in Kotli Hills, Azad Jammu and Kashmir, Pakistan. *Greener J. Agri. Sci.* **2**(7): 345–350.
- Pavón, P.N., Hernández-Trejo, H. and Rico-Gray, V. (2000). Distribution of plant life forms along an altitudinal gradient in the semi-arid valley of Zapotitlán, Mexico. *J. Veg. Sci.* **11**: 39–42.
- Peters, C.M. (1996). **The Ecological and Management of Non-Timber Forest Resources**. World Bank Technical Paper 322, Washington.
- Pharswan, K., Mehta, J.P. and Subodh (2010). Floristic composition and biological spectrum of vegetation in Alpine Meadows of Kedarnath, Gharwal Himalaya. *Nature Sci.* **8**(7): 109–115.
- Pichi-Sermolli, R.E.G. (1957). Una carta geobotanica dell' Africa Orientale (Eritrea, Etiopia, Somalia). *Webbia* **13**: 15–132.
- Pignatti, S. (1995). Land use and human influences in the evergreen broad leaved forest regions of East Asia, the Mediterranean and Australia. In: **Vegetation Science in Forestry**, pp. 199–210 (Peet, R.K., Masuzawa, T., Yamada, I., Fujiwara, K. and Maycock, P.F., eds.). Kluwer Academic Publisher, Dordrecht.
- Raunkiaer, C. (1934). **The Life Forms of Plants and Statistical Plant Geography**. Clarendon Press, Oxford.
- Reddy, K.N., Jadhav, S.N., Reddy, C.S. and Raju, V.S. (2002). Life forms and biological spectrum of Marriguda Reserve Forest, Khammam district, Andhra Pradesh. *Ind. For.* **128**(7): 751–756.
- Sahunalu, P. and Dhanmanonda, P. (1995). Structure and dynamics of dry dipterocarp forest, Sakaerat. In: **Vegetation Science in Forestry**, pp. 465–494 (Peet, R.K., Masuzawa, T., Yamada, I., Fujiwara, K. and Maycock, P.F., eds.). Kluwer Academic Publisher, Dordrecht.
- Shen, C., Xiong, J., Zhang, H., Feng, Y., Lin, X., Li, X., Liang, W. and Chu, H. (2013). Soil pH drives the spatial distribution of bacterial communities along elevation on Changbai Mountain. *Soil Biol. Biochem.* **57**: 204–211.
- Shimwell, D.W. (1971). **Description and Classification of Vegetation**. Sidgewick and Jackson, London.
- Solbrig, O.T. (1993). Plant traits and adaptive strategies: their role in ecosystem function. In: **Biodiversity and Ecosystem Function**, pp. 97–116 (Schulze, E.D. and Mooney, H.A., eds). Springer-Verlag, Berlin.
- Teshome Soromessa and Sebsebe Demissew (2002). Some uses of plants by the Benna, Tsemay and Zeyise people, southern Ethiopia. *Ethiop. J. Nat. Res.* **4** (1): 107–122.
- Tewelde Berhan Gebre Egziabher (1988). Vegetation and environment of the mountains of Ethiopia: Implications for utilization and conservation. *Mt. Res. Dev.* **8**: 211–216.
- Unwin, G.L. (1989). Structure and composition of the abrupt rainforest boundary in the Herberton highland, North Queensland. *Aust. J. Bot.* **37**: 413–428.
- Verma, B.K. and Shukla, G. (1993). Life forms and biological spectrum of the flora of Jalaun District. *J. Indian Bot. Soc.* **72**: 67–68.
- Visalakshi, N. (1995). Vegetation analysis of two tropical dry evergreen forests in southern India. *Trop. Ecol.* **36**: 117–127.

- von Breitenbach, F. (1962). National forestry development planning: A feasibility and priority study on the examples of Ethiopia. *Ethiop. For. Rev.* **3/4**: 41–68.
- von Breitenbach, F. (1963). Forests and wood lands of Ethiopia, a geobotanical contribution to the knowledge of the principal plant communities of Ethiopia, with special regard to forestry. *Ethiop. For. Rev.* **1**: 5–16.
- Westphal, E. (1975). **Agricultural Systems in Ethiopia**. Center for Agricultural Publishing and Documentation, Wageningen.