

Effect of Altitude, Shade, and Processing Methods on Isotope and Biochemical Composition of Green Coffee Beans in Ethiopia

Adugnaw Mintesnot¹ and Nigussie Dechassa²

¹Jimma University, Department of Horticulture, P.O. Box 307, Jimma, Ethiopia

²Haramaya University, Department of Plant Sciences, P.O. Box 138, Dire Dawa, Ethiopia

*Corresponding author: E-mail: adugnawmintesnot@gmail.com

Abstract

Though Arabica coffee has evolved in Ethiopian natural forests, it is being produced without shade in many countries at varying altitudes. Thus, the objective of this study was to determine the composition, fingerprinting, and association of isotopes and biochemical composition as a function of altitude, shade, and processing methods. Carbon, nitrogen and oxygen Isotopes as well as %N and %C of the beans were determined using a continuous flow (CF) EA-IRMS, while caffeine, trigonelline and chlorogenic acid content were determined using HPLC/THERMO. Sucrose content was determined using Gas Chromatography. The study included washed and unwashed coffee samples from altitudes of 1150, 1545 and 1802 meters above sea level. For the determination of the aforementioned components of the bean, univariate analysis of variance, automatic regression modelling, and stepwise canonical discriminant functions were used. Percent N, $\delta^{15}\text{N}$, $\delta^{13}\text{C}$ and %C did not show significant result by altitude. For $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$, unshaded coffee showed higher mean value than the shaded counterpart. Shading disfavoured the composition of isotopic oxygen ($\delta^{18}\text{O}$) as the altitude increases. Shaded coffee at 1150 meters above sea level (m a.s.l.) showed the highest mean value while at 1802 m a.s.l. the lowest for both sucrose and trigonelline composition of green coffee beans. Washed coffee at 1802 m a.s.l. for 5-CQA, unwashed coffee at 1150 m a.s.l. for CFQA, unwashed coffee at 1545 m a.s.l. for TCGA, unwashed coffee at 1150 m a.s.l. for both sucrose and trigonelline showed the highest mean value while unwashed coffee at 1802 m a.s.l. for 5-CQA, TCGA, sucrose and trigonelline, and washed coffee at 1545 m a.s.l. for CFQA showed the least value. The highest mean value was obtained from unwashed-shaded coffee at 1545 m a.s.l. while unwashed- shaded coffee at 1802 was the lowest. The association of isotopes and biochemical composition of green coffee beans was found to be weak although for caffeine and percent N contributed a significant positive weight ($b = 6.604$, $P \leq 0.002$) and %C contributed a significant negative ($b = -0.388$, $P \leq 0.004$) weight for the model. Lowland coffees were well discriminated (94.4% of variation) by 4,5-Dicaffeoylquinic acids (4,5-DCQA) compared to coffees from midlands and highlands.

Keywords: Caffeine, Chlorogenic acids, Coffee quality profile, Coffee value chain, finger printing,

Introduction

Globalization of the coffee market increasingly concerns labelling of the origin of the produce. It is understood coffee quality has a strong association with its origin. Several attempts have been made to discriminate analytically the origin of green and roasted coffees. Signatures (finger printings) of various natures could be used for sorting the products of unique nature (Meinzer *et al.*, 1990). A review by Rodrigues *et al.* (2009) discloses different approaches of identifying origins of food stuffs for labelling. Analytical methods such as gas chromatography–mass spectrometry (Costa Freitas *et al.*, 2001), near infrared spectroscopy (Bertrand *et al.*, 2005), determination of organic compounds such as chlorogenic acids and fatty acid profiles (Martín *et al.*, 2001), tocopherols and triglycerides (González *et al.*, 2001) and stable isotope analysis of specific compounds extracted from the green coffee beans (Adugnaw, 2014) have used extensively with promising results.

Ethiopia, despite being the place of origin and diversification of Arabica coffee, has not yet fully exploited its genetic and environmental wealth discriminated in the world market (Petit, 2007). The country could play important roles as a supplier of some of the best coffees in the ‘global coffee value chain’, which could be accompanied by discrimination and promotional work that could identify it with unique inherent flavour. In this

regard, the approach using stable isotope ratios and biochemical composition for origin discrimination seem quite promising. Serra *et al.* (2005) inferred the continental origin (Africa, Asia and America) of coffee using the combination of the isotopic fingerprints of carbon, nitrogen and boron, which are reportedly used as integrated proxies for environmental conditions and agricultural practices. The authors were successful in identifying 88% of the intercontinental origin of the samples but not an intracontinental discrimination by means of principal component analysis. Variation in stable carbon isotope discrimination among genotypes grown under identical conditions has been used as a means of selecting for improved water use efficiency (WUE) and yield in C₃ crop species (Meinzer, *et al.*, 1990).

Environmental factors, such as altitude, shade and rainfall may contribute to the quality of the coffee beverage (Bote and Struik, 2011; Likassa Ebisa, 2014). However, further studies are needed to investigate additional environmental characteristics that affect coffee quality (Rodrigues *et al.*, 2009) reported that an evaluation of the caffeine content of beans from 99 progenies revealed intra- and inter-progeny variability. In 68 progenies from the Kaffa region they found caffeine values in the range 0.46-2.82%, and in 22 progenies from Illubabor region these values ranged from 0.42 to 2.90%. Bertrand *et al.*

(2006) reported a significant effect of Elevation on bean biochemical composition of the cultivar 'Caturra' where they found increased concentration of chlorogenic acid and caffeine with increasing elevation above 1100 m. The reverse trend was observed for trigonelline concentration in one situation and had no effect on another. Caffeine and fat concentrations increased with increasing elevation up to a certain level.

Natural diversity in Ethiopian specialty coffees may represent an important opportunity to add value to the economic and social development of the country as well as to the advancement of coffee science. Systematically establishing a relationship between geographic origin of coffee and its quality leading to the generation of reliable data is a priority area in Ethiopia (Adugnaw, 2014). However, studies on specialty coffee discrimination and authentication methods have been only recent phenomenon in Ethiopia. Such methods could complement the existing subjective organoleptic classifications methods. Thus, the objective of this study was to describe the composition, fingerprinting, and association of coffee bean isotopes and biochemical compounds as a function of altitude, shade and processing methods.

Materials and Methods

Site selection and experimental design

The study included washed coffee and unwashed coffee samples originating from south-western Ethiopia from three farms located at different altitudes (1150, 1545 and 1802 metres above sea level designated as lowland, midland and highland coffee growing areas, respectively (MoA, 2003). According to the data obtained from National Meteorological Services Agency (2010) the region is characterized by monomodal rainfall of about 1565 mm, maximum and minimum temperatures of 26.1 and 13.2 °C, respectively, relative humidity of 73.3%, sunshine hours of 5.4 and altitude ranging between 1150 and 1820 meters above sea level. The farms were purposely selected considering the altitudinal ranges and uniformity to management methods. The selected farms belonged to Coffee Plantation Development Enterprise. Nine sub coffee farms were selected from the three farms. From the three coffee farms, shaded and unshaded plots were identified and two types of processing methods were applied in three replications. Ripe coffee cherries were handpicked at their peak ripening phases during the 2010/11 crop season out of which washed and unwashed green coffee beans were carefully prepared without contamination.

Table 1. Description of study sites

| Farm | Region | Latitude | Longitude | Altitude (m a.s.l.) |
|---------------|------------------|-------------|--------------|---------------------|
| Bebeka (N= 3) | Benchmajil/ SNNP | 6°56.580' N | 35°30.607' E | 1150 |
| Kossa (N= 3) | Jimma/ Oromia | 7°57.223' N | 36°52.664' E | 1802 |
| Goma (N= 3) | Jimma/ Oromia | 7°55.253' N | 36°37.069' E | 1545 |

Laboratory analysis

Green coffee beans were subjected to freeze drying just before grinding to fine powder using a hand-held electrical Blade coffee grinder (Bosch MKM 6003 UC, Bean Container Capacity: 75g, Power: 180 Watt). Grinding was assumed to be sufficient when the powder escaped to the ceiling of the cap of the grinder, and it was immediately packed in a plastic cup with a tight stopper, and kept in a deep freezer until laboratory analysis.

Carbon, Nitrogen isotopes, and elements measurement

Carbon and Nitrogen Isotopes and %N and %C were measured using a continuous flow (CF) EA-IRMS, Sercon stable isotope mass spectrometers (UK) (OTSUKI, 1983). Finely ground green coffee bean powder (0.95 - 1.4 mg each) rolled in small tin capsules were loaded onto an auto sampler in a duplicate. The samples were purged by a helium (He) flow into a combustion tube and completely oxidized at a temperature of 1000 °C. A packed GC column removes impurities and separates N₂ and CO₂. A mass spectrometer ionizes gaseous molecules and separates the ions into a spectrum according to their mass-to-charge ratio (m/z), using electric and magnetic fields. The

relative abundances of the molecules of different m/z were then found by measuring the currents generated by these separated ion beams. A high vacuum keeps the analyzer pressure low enough (10-5 mbar) to reduce collisions between ions and background gas to an acceptable level. A permanent magnetic field was used (fixed B) and masses were selected by varying the tensions of the electric field V. A universal triple collector was used and B and V were kept constant for each element that has to be measured (Otsuki, 1983).

Oxygen ($\delta^{18}O$) isotope measurement (VSMOW)

Oxygen isotope measurement was done via a TC/EA (thermal conversion/elemental analyzer) coupled to an isotope ratio mass spectrometer (IRMS) (20-20, SerCon Ltd, Crewe, UK) (SerCon, 1983.). The green coffee bean powder was further dried overnight at 60-80 °C and stored in a desiccator after which each of a 1 mg samples was rolled in a silver cup in a duplicate wand loaded onto an auto sampler. The samples were pyrolyzed at 1400 °C in a molybdenum lined, aluminium oxide reduction tube filled with glassy carbon and topped with a glassy carbon crucible. The produced N₂ and

CO gases were separated via a 1 m gas chromatography (GC) column (E3030, Elemental Microanalysis Ltd., Okehampton, UK) at a temperature of 50 °C, helium (He) carrier gas pressure of 1.6 bar, Helium (He) flow retention time of about 250 s, and sample analysis time of 1000 s, and analyzed via isotope ratio mass spectrometry (IRMS) for $\delta^{18}\text{O}$. The internationally accepted reference values of $\delta^{18}\text{O}$ -KNO₃ for USGS32 (25.7±0.4 and USGS34 (-27.8±0.4, (Brand *et al.*, 2009) and $\delta^{18}\text{O}$ -NaNO₃ for USGS35 (56.8±0.3 (IAEA, 2004) were used to correct raw $\delta^{18}\text{O}$ values to $\delta^{18}\text{O}$ (‰) (Otsuki, 1983).

Caffeine, Trigonelline and Chlorogenic acids

The caffeine and chlorogenic acid contents were determined using HPLC/THERMO following the method of Alonso-Salces *et al.* (2009): 0.1 g freeze dried green coffee powder was weighed in an Erlenmeyer flask of 50 ml. 10 ml of MeOH/Acetic Acid (30:7.5:2.5) containing 2 mg/ml ascorbic acid was added and then placed in an ultrasonic bath for 15 minutes. The extract was filtered using Whatman filter papers No.2, and subsequently over a 0.45 micrometer PTFE filter after which 1 ml of the filtrate was taken in a vial and injected on HPLC/THERMO. The standard solutions of chlorogenic acid, caffeine, and caffeic acid were mixed each at 0.5/1/1.5 mg/ml in one mixture in methanol and each solution was injected twice for calibration. A calibration curve was made using the

standard concentration and area of sample and subsequently used to calculate the composition of the respective biochemical component using the area generated after the retention time. The detection was carried out at 278 nm for caffeine and trigonelline, and 324 nm for chlorogenic acids (CGAs). For the identification and quantitative analysis, a standard curve was prepared using standards of caffeine, trigonelline and Chlorogenic acids.

Sucrose measurement

Sucrose of the coffee beans was determined using GC VARIAN 3800 following the standard method. A sample of green coffee powder was freeze-dried and weighed (0.5 - 1 g) in 50 ml volumetric flask to which 30 ml distilled water plus 5 ml frozen Internal Standard Solution (ISS) (Phenyl-B-D-pyranoside) was added. It was placed at 60 °C for 30 minutes after which it was cooled. Next, 0.5 ml each of Carre I (15 g ZnSO₄ and 7.5 g Carre II (K₄FCICN)₆) was added to de-protein the sample. The distilled water was then filled to label of the mark on the 250 volumetric flasks and shaken well to homogenize the mixture. The solution was immediately filtered with Whatman filter papers, and subsequently 1 ml filtrate was taken in small bottles using glass Pasteur pipette and dried under nitrogen-drier using hollow needles to let nitrogen in to the bottle. To this dry extract was added 1 ml STOX (2.5 g hydroxylamine hydrochloride diluted with dry pyridine to 100 ml) under

hood and kept at 60 °C for 30 minutes and then cooled down. Then, 1 ml of HMDS (hexamethyldisilazan) was added and subsequently 0.1 ml TFA (trifluor acetic acid) was added before sedimentation for 60 minutes to get clear extract solution. From the clear extract solution, 1 ml was taken in a vial with rubber stopper to inject to GC VARIAN 3800.

Statistical analysis

Univariate Analysis of Variance and Tukey HSD method of mean separation was applied to study the significant difference. Moreover, multiple regression analyses were conducted to examine the relationship between coffee quality attributes and various potential predictors using SPSS 16 v2 software (SPSS Inc.2007). Stepwise canonical discriminant function and Stepwise Regression between isotopes and biochemical composition of green coffee beans were conducted to identify the best predictor for discriminating coffees of different altitudinal gradients.

Results and Discussion

Isotopic and Percent N and C composition of green coffee beans as a function of altitude, shade, and processing methods

This study showed that there were significant effects ($P \leq 0.05$) of altitude on $\delta^{15}\text{N}$ and $\delta^{13}\text{C}$. Shade also showed significant effect on $\delta^{13}\text{C}$, and

non-significant, otherwise ($p>0.05$) (Table 2-4 and Figure 1). With regard to $\delta^{15}\text{N}$, the mean value at 1802 meters above sea level was significantly lower than the mean value at both 1150 and 1545 meters above sea level. The trend showed that $\delta^{15}\text{N}$ concentration increased in the order of lowland, midland and highland samples (Figure 1). The mean value of $\delta^{15}\text{N}$ at 1802 meters above sea level was significantly lower as compared to the mean value at both 1150 and 1545 meters above sea level likely due to lower microbial deposition and hence discrimination during the decomposition process of N at higher altitudes. Most elements of biological interest have two or more stable isotopes, although one isotope is usually present in far greater abundance (Ehleringer and Rundel, 1989). While there is no evidence for biological fractionation of the isotopes, they may be potentially useful markers of ecosystem process studies.

One of the factors controlling the mineral content of plant material is the availability of plant nutrient in the nutrient medium. Moreover, the work of several researchers revealed that natural abundance $\delta^{15}\text{N}$ of plants reflects the net effect of a range of processes (Shibuya *et al.*, 2006). In this regard, the presence of multiple N-sources with distinct isotopic values, mycorrhizal associations, temporal and spatial variation in N availability, and changes in plant demand can all influence plant $\delta^{15}\text{N}$. In addition, the farm at higher altitude

may be planted with legume shade trees where appears and may supply N by atmospheric nitrogen fixation processes instead of discriminating against $\delta^{15}\text{N}$. Consequently, the $\delta^{15}\text{N}$ values of leguminous plants are often close to 0‰ (Ehleringer and Rundel, 1989). Ehleringer and Rundel (1989) showed that roughly 30% of the nitrogen fixed by a legume was transferred to the associated coffee trees. Snoeck, *et al.* (2000) also proposed certain amount of N transferred by legume to sole coffee trees *via* litter fall.

The mean value for $\delta^{13}\text{C}$ at 1150 meters above sea level was significantly lower than the mean value at 1545 and 1802 meters above sea level. There was a significant effect of shading on $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ levels of green coffee beans but not on %N and %C. For $\delta^{13}\text{C}$ mean value at 1150 meters above sea level was significantly lower as compared to the value at 1545 and 1802 m a.s.l. and the shaded coffee showed higher mean value most probably due to better temperature regulation for

photosynthetic process of higher altitudes and shade. In C_3 plants, discrimination against ^{13}C by the carboxylating enzyme, Rubisco ($\gg 27\%$), is linked to photosynthesis via the ratio of intercellular to atmospheric CO_2 concentrations (c_i/c_a) (Dawson *et al.* (2002)). This ratio reflects the relative magnitudes of net assimilation and stomatal conductance that relate to demand and supply of CO_2 , respectively. Carbon-13 data are thus a useful index for assessing intrinsic water use efficiency (A/g ; the ratio of carbon acquired to water vapor losses via stomatal conductance, g) and may even provide information on actual water use efficiency (the ratio of assimilation to transpiration) when the leaf-to-air vapor pressure difference is known (Ehleringer and Vogel, 1993). A few studies are now using C isotope data to investigate competition how the efficiency of resource use varied in the presence or absence of different neighbours (Williams *et al.* 1991). With regard to $\delta^{18}\text{O}$ and $\delta^{13}\text{C}$, unshaded coffee showed higher mean value than the shaded counterpart.

Table 2. Effect of altitude, shade and processing Methods on Stable Isotopes and %N and %C

| Factor | Level | %N | $\delta^{15}\text{N}$ | %C | $\delta^{13}\text{C}$ | $\delta^{18}\text{O}$ |
|----------|-----------|----------------------|-----------------------|-----------------------|------------------------|-----------------------|
| Altitude | 1150 (12) | 2.4±0.1 ^a | 5.9±0.3 ^a | 46.1±0.8 ^a | -27.6±0.2 ^b | 31.4±0.3 ^a |
| | 1545(12) | 2.4±0.1 ^a | 5.3±0.3 ^a | 46.8±0.8 ^a | -27.6±0.2 ^b | 32.1±0.3 ^a |
| | 1802(12) | 2.4±0.1 ^a | 4.4±0.3 ^b | 46.5±0.8 ^a | -26.7±0.2 ^a | 31.4±0.3 ^a |
| | P-value | 0.770 | 0.006 | 0.861 | <0.001 | 0.184 |
| Shade | Shaded | 2.4±0.0 | 5.0±0.3 | 46.3±0.7 | -27.6±0.1 | 31.1±0.2 |
| | Unshaded | 2.4±0.0 | 5.3±0.3 | 46.7±0.7 | -27.0±0.1 | 32.1±0.2 |
| | P-value | 0.904 | 0.419 | 0.712 | 0.005 | 0.005 |
| GM | | 2.4±0.0 | 5.2±0.2 | 46.5±0.5 | -27.3±0.1 | 31. ±0.2 |
| SD | | 0.2 | 1.2 | 2.5 | 0.7 | 1.7 |
| CV% | | 6.5 | 23.3 | 5.4 | -2.7 | 5.4 |




Figure 1. Isotopes and multi-element composition of green coffee beans as a function of altitude gradient

With regard to $\delta^{18}\text{O}$ unshaded coffee showed higher mean value as compared to shaded counterpart likely due to better accumulation of the heavy isotope at unshaded condition because of less humidity as compared to humid shaded condition. Oxygen isotope ratios have been most useful in tracing and describing water movement and as indicators of humidity regimes. Oxygen derived from CO_2 undergoes a complete exchange with the oxygen of the water in the plant during the synthesis of cellulose, and thus $\delta^{18}\text{O}$ of tissue water is the primary influence on the $\delta^{18}\text{O}$ of fixed oxygen in cellulose (Ehleringer and Rundel, 1989). The $\delta^{18}\text{O}$ generally varies with ambient humidity, which in turn reflects changes in water use (Meinzer, *et al.*, 1990). The $\delta^{18}\text{O}$ of leaf and tree ring cellulose are largely determined by the integrated leaf-to-air vapor pressure gradient during photosynthetic gas exchange (Ehleringer and Vogel, 1993). This leaf-air vapour pressure gradient changes with environmental

conditions (atmospheric humidity, soil moisture, air temperature) and plant response to these environmental changes (e.g., changes in water use, leaf temperature, and net assimilation). Measurement of the $\delta^{18}\text{O}$ composition of plant tissues may thus aid for the interpretation of differences in $\delta^{13}\text{C}$ among individual plants growing in the same location and among species in different environments.

The bi-factor interaction effect of Altitude * Shade, and Altitude * Processing methods were significant for $\delta^{18}\text{O}$ (Table s 3 & 4), and non-significant, otherwise. Shading disfavoured the composition of $\delta^{18}\text{O}$ in green coffee beans as the altitude increases while favouring at lower altitude. Unwashed coffee showed higher $\delta^{18}\text{O}$ composition of green coffee beans at higher altitudes but lower at low altitude. Both shading and washing could discriminate coffees of higher altitudes from that of lowlands. The tri-factor interactions resulted in significant effect of none of

the response variables (Table 3). Moreover, the determination of water use efficiency (WUE) is greatly improved by the simultaneous use of $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$ in plant tissues (Saurer *et al.* 1997). By considering concurrent variations $\delta^{13}\text{C}$ and $\delta^{18}\text{O}$, one can distinguish between biochemical and stomatal limitations to photosynthesis in response to a change in environmental conditions.

Shading disfavoured the composition of $\delta^{18}\text{O}$ in green coffee beans as the altitude increases while favouring at lower altitude. This indicates shading requirement reduces as altitude increases because of lower amplitude of temperature and higher relative humidity. Bote and Struik (2011) noted that shade triggers differences in physiological behaviour of the coffee plants, such as improved photosynthesis and increased leaf area index, resulting in better performance than possible in direct sun light. Shaded plants had greater biochemical and physiological potential for high dry matter production which would help them to maintain high coffee yields in the long term. Processing methods did not show any significant effect on any of the studied variables. Unwashed coffee showed higher $\delta^{18}\text{O}$ composition of green coffee beans at higher altitudes but lower at low altitude. Both shading and washing could discriminate coffees of higher altitudes from that of lowlands.

Rodrigues *et al.* (2009) observed differences on stable isotopic and elemental composition and associated

them by altitude and precipitation values associated with the different geographic locations. The isotopic compositions of carbon and nitrogen, which are the main elements of living organisms, are used as proxies for environmental parameters that characterize a certain crop or area (Meinzer *et al.*, 1990). Since different elements represent different characteristics of the environment (water cycle, plant physiology, agricultural practices, hydric stress, soil geology, etc.), the combination of these indicators may be used to unequivocally identify a certain ecosystem of origin. A noticeable feature of using stable isotope abundance data, in a multivariate approach to discriminate the origin of a product, is that each single isotopic signature is in fact the result of more than one unique phenomenon, and is thus very difficult to counterfeit for fraud purposes.

A study by Sera *et al.* (2005) showed that the small climatic differences among regions which do not have very marked differences may show little isotopic variation. Stable isotope ratios of carbon, nitrogen and boron have been shown to be good indicators of geographical-dependent parameters, and therefore to be useful tools to infer the region of production of green coffee (Rodrigues *et al.*, 2009; Adugnaw, 2014). An inherent limitation of isotope ratio techniques is due to the mismatching between national borders and climatic borders, delimiting different geographical regions; for this reason, coffees

produced in small adjacent countries with similar climatic conditions cannot be distinguished from one another on the basis of different isotope ratio values, while large countries with a

large variety of climatic areas may show samples with a range of isotope ratio values, displaying wide dispersion.

Table 3. Mean effect of bi-factor interactions on the composition of stable isotopes and %N & %C of green coffee beans

| Altitude * Shade | | %N | $\delta^{15}\text{N}$ | %C | $\delta^{13}\text{C}$ | $\delta^{18}\text{O}$ |
|------------------|----------|---------|-----------------------|----------|-----------------------|-----------------------|
| 1150 | Shaded | 2.3±0.1 | 5.4±0.4 | 45.9±1.2 | -27.9±0.2 | 32.2±0.4 |
| | Unshaded | 2.4±0.1 | 6.4±0.4 | 46.4±1.2 | -27.4±0.2 | 30.6±0.4 |
| 1545 | Shaded | 2.4±0.1 | 5.4±0.4 | 46.5±1.2 | -27.8±0.2 | 31.6±0.4 |
| | Unshaded | 2.4±0.1 | 5.2±0.4 | 47.1±1.2 | -27.3±0.2 | 32.5±0.4 |
| 1802 | Shaded | 2.4±0.1 | 4.3±0.4 | 46.5±1.2 | -27.1±0.2 | 29.4±0.4 |
| | Unshaded | 2.4±0.1 | 4.4±0.4 | 46.6±1.2 | -26.3±0.2 | 33.3±0.4 |
| P-value | | 0.89 | 0.442 | 0.967 | 0.676 | <0.001 |

Table 4. Mean effect of bi-factor interaction on the composition of stable isotopes and %N & %C of green coffee beans

| Altitude * Processing | | %N | $\delta^{15}\text{N}$ | %C | $\delta^{13}\text{C}$ | $\delta^{18}\text{O}$ |
|-----------------------|----------|---------|-----------------------|----------|-----------------------|-----------------------|
| 1150 | Washed | 2.3±0.1 | 5.9±0.4 | 45.7±1.2 | -27.4±0.2 | 31.7±0.4 |
| | Unwashed | 2.4±0.1 | 6.0±0.4 | 46.5±1.2 | -27.9±0.2 | 31.0±0.4 |
| 1545 | Washed | 2.3±0.1 | 5.7±0.4 | 46.2±1.2 | -27.6±0.2 | 31.0±0.4 |
| | Unwashed | 2.4±0.1 | 4.9±0.4 | 47.4±1.2 | -27.5±0.2 | 33.1±0.4 |
| 1802 | Washed | 2.4±0.1 | 4.7±0.4 | 45.9±1.2 | -26.5±0.2 | 31.2±0.4 |
| | Unwashed | 2.4±0.1 | 4.0±0.4 | 47.2±1.2 | -26.8±0.2 | 31.5±0.4 |
| P-value | | 0.875 | 0.552 | 0.977 | 0.365 | 0.01 |

Biochemical composition of green coffee beans as a function of altitude, shade and processing methods

Altitude had significant effect ($P \leq 0.05$) on 4-Caffeoylquinic acid (4-CQA), 5-Caffeoylquinic acid (5-CQA), Feruloylquinic acids (FQA), 3,4-Dicaffeoylquinic acids (3,4-DCQA), 4,5-Dicaffeoylquinic acids (4,5-DCQA), CFQA, Total Chlorogenic Acids (TCGA), sucrose, trigonelline composition of green coffee beans. However, altitude did not have a

significant effect ($P > 0.05$) on Caffeine and 3-CQA compositions. For 4-CQA and 5-CQA lowland samples showed the least mean value as compared to both midland and highland samples. On the other hand, for FQA, 4,5-DCQA, and TCGA, the highland sample showed the least mean value as compared to both lowland and midland samples.

The compositions of 3,4-DCQA, CFQA, and sucrose were the lowest for samples of coffee beans obtained from the lowlands compared to those

obtained from both midland and highlands. The effect of shade was significant for Caffeoylferuloylquinic acids (CFQA), and non-significant, otherwise. Unshaded coffee gave higher mean value of CFQA composition of green coffee beans as compared to shaded counterpart. Processing methods had significant effect on 3-CQA, 5-CQA, 4,5-DCQA, CFQA, TCGA and trigonelline composition, and non-significant, otherwise. Washed coffee showed higher mean value for 5-CQA, 4,5-DCQA, TCGA, and trigonelline composition of green coffee beans while unwashed coffee resulted in higher mean value for 3-CQA and CFQA composition of green coffee beans. As could be seen on Figure 2 the mean value for 3,4-DCQA, 4,5-DCQA, CFQA, TCGA, sucrose, and trigonelline decreased in the order of lowland, midland and highland samples. However, for caffeine, 4-CQA, and 5-CQA the value increased in the order of midland, highland and lowland where as for 3-CQA and FQA in the order of midland, lowland and highland samples.

The effect of Altitude*Shade interaction was significant for sucrose and trigonelline composition of green coffee beans, and non-significant, otherwise (Table 5). In this regard, shaded coffee at 1150 meters above sea level (m a.s.l.) showed the highest mean value while at 1802 m a.s.l. the lowest for both sucrose and trigonelline composition of green coffee beans. The effect of Altitude*Processing was significant

for 5-CQA, CFQA, TCGA, sucrose and trigonelline of composition of green coffee beans (Table 6). Washed coffee at 1802 m a.s.l. for 5-CQA, unwashed coffee at 1150 m a.s.l. for CFQA, unwashed coffee at 1545 m a.s.l. for TCGA, unwashed coffee at 1150 m a.s.l. for both sucrose and trigonelline showed the highest mean value while unwashed coffee at 1802 m a.s.l. for 5-CQA, TCGA, sucrose and trigonelline, and washed coffee at 1545 m a.s.l. for CFQA showed the least value. The study did not observe any significant effect of Shade* Processing methods. The Tri-Factor Interaction Effect of Altitude, Shade and Processing methods showed significant effect on 5-CQA composition of green coffee beans, and non-significant, otherwise (Table 7). In this regard, the highest mean value was obtained from unwashed coffee and shaded coffee at 1545 m a.s.l. while unwashed coffee and shaded coffee at 1802 was the lowest.

For 4-CQA and 5-CQA lowland samples showed the least mean value as compared to both midland and highland samples. The composition of 3,4-DCQA, CFQA, and sucrose was significantly the lowest at lowland compared to both midland and highland samples. On the other hand, for FQA, 4,5-DCQA, and TCGA, the highland sample showed the least mean value as compared to both lowland and midland samples. A study by Bertrand *et al.* (2006) is in complement with this result showing that elevation had a significant effect on bean biochemical composition,

with chlorogenic acid and fat concentrations increasing with increasing elevation for some varieties and little effect on the variation of

chlorogenic acid concentration but not on fat concentration.

Biochemical composition of green coffee beans as a function of altitude




Figure 2. Biochemical composition of green coffee beans as a function of altitude gradient

Table 5. Effect of Altitude, shade and processing methods on biochemical composition of green coffee beans

| | Caffeine | 3-CQA | 4-CQA | 5-CQA | FQA | 3,4-DCQA | 4,5-DCQA | CFQA | TCGA | sucrose | trigonelline |
|---------------------------|-----------------------|----------------------|----------------------|-----------------------|----------------------|----------------------|----------------------|----------------------|-----------------------|-----------------------|-----------------------|
| Altitude | | | | | | | | | | | |
| 1150 (120) | 13.6±0.2 ^a | 3.9±0.2 ^a | 5.7±0.1 ^b | 27.2±0.3 ^b | 4.5±0.2 ^a | 4.1±0.2 ^a | 8.1±0.3 ^a | 4.0±0.1 ^a | 57.4±0.6 ^a | 58.8±2.2 ^a | 87.6±0.3 ^a |
| 1545 (12) | 14.4±0.2 ^a | 4.0±0.2 ^a | 6.4±0.1 ^a | 28.9±0.3 ^a | 4.8±0.2 ^a | 2.9±0.2 ^b | 7.1±0.3 ^a | 3.2±0.1 ^b | 57.2±0.6 ^a | 50.1±2.2 ^b | 86.0±0.3 ^b |
| 1802 (12) | 13.9±0.2 ^a | 3.8±0.2 ^a | 6.0±0.1 ^a | 28.2±0.3 ^a | 4.2±0.2 ^b | 2.3±0.2 ^b | 5.8±0.3 ^b | 3.1±0.1 ^b | 53.2±0.6 ^b | 46.2±2.2 ^b | 81.6±0.3 ^c |
| P-value | 0.08 | 0.585 | 0.013 | 0.002 | 0.014 | <0.001 | <0.001 | <0.001 | <0.001 | 0.001 | <0.001 |
| Shade | | | | | | | | | | | |
| Shaded | 14.2±0.2 | 3.8±0.1 | 5.9±0.1 | 27.9±0.2 | 4.6±0.1 | 2.9±0.2 | 7.2±0.2 | 3.2±0.1 | 55.4±0.5 | 52.9±1.8 | 84.8±0.3 |
| Unshaded | 13.7±0.2 | 4.1±0.1 | 6.1±0.1 | 28.2±0.2 | 4.4±0.1 | 3.2±0.2 | 6.8±0.2 | 3.7±0.1 | 56.5±0.5 | 50.5±1.8 | 85.3±0.3 |
| P-value | 0.111 | 0.123 | 0.201 | 0.531 | 0.259 | 0.215 | 0.354 | 0.002 | 0.12 | 0.344 | 0.274 |
| Processing methods | | | | | | | | | | | |
| Washed | 14.2±0.2 | 3.7±0.1 | 6.0±0.1 | 28.5±0.2 | 4.6±0.1 | 3.1±0.2 | 8.1±0.2 | 2.8±0.1 | 56.7±0.5 | 49.2±1.8 | 85.9±0.3 |
| Unwashed | 13.7±0.2 | 4.1±0.1 | 6.0±0.1 | 27.6±0.2 | 4.3±0.1 | 3.1±0.2 | 5.9±0.2 | 4.1±0.1 | 55.2±0.5 | 54.2±1.8 | 84.2±0.3 |
| P-value | 0.111 | 0.029 | 0.867 | 0.025 | 0.085 | 0.945 | <0.001 | <0.001 | 0.029 | 0.064 | <0.001 |

Table 6. The Bi-Factor Interaction Effect of Altitude, Shade and Processing methods on Biochemical Composition of Green Coffee Beans

| | | Caffeine | 3-CQA | 4-CQA | 5-CQA | FQA | 3,4-DCQA | 4,5-DCQA | CFQA | TCGA | sucrose | trigonelline |
|----------------------------|-----------------|----------|---------|---------|----------|---------|----------|----------|---------|----------|----------|--------------|
| Altitude*Shade | | | | | | | | | | | | |
| 1150 | Shaded | 14.1±0.3 | 4.0±0.2 | 5.7±0.2 | 26.9±0.4 | 4.7±0.2 | 4.0±0.3 | 7.9±0.4 | 3.8±0.2 | 57.1±0.8 | 64.7±3.1 | 88.0±0.5 |
| | Unshaded | 13.1±0.3 | 3.9±0.2 | 5.7±0.2 | 27.4±0.4 | 4.2±0.2 | 4.2±0.3 | 8.2±0.4 | 4.2±0.2 | 57.7±0.8 | 53.0±3.1 | 87.3±0.5 |
| 1545 | Shaded | 14.6±0.3 | 3.8±0.2 | 6.3±0.2 | 28.8±0.4 | 4.9±0.2 | 2.7±0.3 | 7.4±0.4 | 2.9±0.2 | 56.7±0.8 | 48.4±3.1 | 85.0±0.5 |
| | Unshaded | 14.1±0.3 | 4.2±0.2 | 6.5±0.2 | 29.0±0.4 | 4.8±0.2 | 3.0±0.3 | 6.8±0.4 | 3.5±0.2 | 57.8±0.8 | 51.8±3.1 | 87.0±0.5 |
| 1802 | Shaded | 13.8±0.3 | 3.5±0.2 | 5.7±0.2 | 28.2±0.4 | 4.1±0.2 | 2.1±0.3 | 6.1±0.4 | 2.9±0.2 | 52.5±0.8 | 45.7±3.1 | 81.5±0.5 |
| | Unshaded | 13.9±0.3 | 4.1±0.2 | 6.2±0.2 | 28.2±0.4 | 4.2±0.2 | 2.5±0.3 | 5.5±0.4 | 3.3±0.2 | 53.9±0.8 | 46.7±3.1 | 81.6±0.5 |
| | P-value | 0.295 | 0.480 | 0.366 | 0.883 | 0.354 | 0.878 | 0.428 | 0.618 | 0.901 | 0.049 | 0.019 |
| 1150 | Washed coffee | 13.8±0.3 | 3.8±0.2 | 5.7±0.2 | 27.3±0.4 | 4.7±0.2 | 4.1±0.3 | 9.3±0.4 | 3.1±0.2 | 57.8±0.8 | 49.4±3.1 | 87.6±0.5 |
| Altitude*Processing | | | | | | | | | | | | |
| 1545 | Unwashed coffee | 13.4±0.3 | 4.1±0.2 | 5.8±0.2 | 27.0±0.4 | 4.2±0.2 | 4.1±0.3 | 6.8±0.4 | 5.0±0.2 | 57.0±0.8 | 68.2±3.1 | 87.7±0.5 |
| | Washed coffee | 14.4±0.3 | 3.8±0.2 | 6.3±0.2 | 28.6±0.4 | 4.8±0.2 | 2.7±0.3 | 7.8±0.4 | 2.4±0.2 | 56.4±0.8 | 50.3±3.1 | 84.3±0.5 |
| | Unwashed coffee | 14.3±0.3 | 4.3±0.2 | 6.4±0.2 | 29.1±0.4 | 4.9±0.2 | 3.0±0.3 | 6.5±0.4 | 4.0±0.2 | 58.1±0.8 | 49.9±3.1 | 87.6±0.5 |
| 1802 | Washed coffee | 14.3±0.3 | 3.6±0.2 | 6.0±0.2 | 29.5±0.4 | 4.5±0.2 | 2.5±0.3 | 7.2±0.4 | 2.8±0.2 | 55.9±0.8 | 48.0±3.1 | 85.8±0.5 |
| | Unwashed coffee | 13.4±0.3 | 4.0±0.2 | 5.9±0.2 | 26.8±0.4 | 3.9±0.2 | 2.2±0.3 | 4.5±0.4 | 3.3±0.2 | 50.5±0.8 | 44.4±3.1 | 77.3±0.5 |
| | P-value | 0.444 | 0.933 | 0.933 | 0.003 | 0.199 | 0.6 | 0.165 | 0.002 | 0.001 | 0.003 | <0.001 |

Table 7. The Tri-Factor Interaction Effect of Altitude, Shade and Processing on Biochemical Composition of Green Coffee Beans (mg/g dwb)

| Altitude | Shade | Processing methods | Caffeine | 3-CQA | 4-CQA | 5-CQA | FQA | 3,4-DCQA | 4,5-DCQA | CFQA | TCGA | sucrose | trigonelline |
|----------|----------|--------------------|----------|---------|---------|----------|---------|----------|----------|---------|----------|----------|--------------|
| 1150 | Shaded | Washed coffee | 14.2±0.5 | 3.8±0.3 | 5.7±0.3 | 26.6±0.6 | 5.0±0.3 | 4.1±0.4 | 9.2±0.6 | 3.0±0.2 | 57.2±1.1 | 50.8±4.4 | 87.8±0.7 |
| | | Unwashed coffee | 13.9±0.5 | 4.1±0.3 | 5.8±0.3 | 27.2±0.6 | 4.5±0.3 | 3.9±0.4 | 6.6±0.6 | 4.7±0.2 | 56.9±1.1 | 78.5±4.4 | 88.2±0.7 |
| | Unshaded | Washed coffee | 13.3±0.5 | 3.7±0.3 | 5.6±0.3 | 27.9±0.6 | 4.4±0.3 | 4.1±0.4 | 9.4±0.6 | 3.2±0.2 | 58.3±1.1 | 48.0±4.4 | 87.3±0.7 |
| | | Unwashed coffee | 12.9±0.5 | 4.1±0.3 | 5.7±0.3 | 26.8±0.6 | 4.0±0.3 | 4.2±0.4 | 7.0±0.6 | 5.2±0.2 | 57.1±1.1 | 57.9±4.4 | 87.2±0.7 |
| 1545 | Unshaded | Washed coffee | 14.6±0.5 | 3.5±0.3 | 6.2±0.3 | 29.5±0.6 | 4.5±0.3 | 2.2±0.4 | 7.8±0.6 | 1.9±0.2 | 55.6±1.1 | 46.5±4.4 | 83.1±0.7 |
| | | Unwashed coffee | 14.6±0.5 | 4.2±0.3 | 6.3±0.3 | 28.0±0.6 | 5.3±0.3 | 3.2±0.4 | 7.1±0.6 | 3.9±0.2 | 57.8±1.1 | 50.4±4.4 | 86.8±0.7 |
| | Shaded | Washed coffee | 14.2±0.5 | 4.0±0.3 | 6.4±0.3 | 27.7±0.6 | 5.0±0.3 | 3.2±0.4 | 7.7±0.6 | 3.0±0.2 | 57.2±1.1 | 54.1±4.4 | 85.5±0.7 |
| | | Unwashed coffee | 14.0±0.5 | 4.4±0.3 | 6.5±0.3 | 30.2±0.6 | 4.5±0.3 | 2.9±0.4 | 5.9±0.6 | 4.1±0.2 | 58.3±1.1 | 49.4±4.4 | 88.5±0.7 |
| 1802 | Unshaded | Washed coffee | 14.5±0.5 | 3.2±0.3 | 5.6±0.3 | 29.8±0.6 | 4.4±0.3 | 2.1±0.4 | 7.7±0.6 | 2.4±0.2 | 55.2±1.1 | 48.0±4.4 | 85.9±0.7 |
| | | Unwashed coffee | 13.1±0.5 | 3.9±0.3 | 5.8±0.3 | 26.5±0.6 | 3.9±0.3 | 2.1±0.4 | 4.5±0.6 | 3.3±0.2 | 49.9±1.1 | 43.3±4.4 | 77.2±0.7 |
| | Shaded | Washed coffee | 14.2±0.5 | 4.0±0.3 | 6.4±0.3 | 29.1±0.6 | 4.6±0.3 | 2.8±0.4 | 6.7±0.6 | 3.2±0.2 | 56.7±1.1 | 48.1±4.4 | 85.8±0.7 |
| | | Unwashed coffee | 13.7±0.5 | 4.1±0.3 | 6.1±0.3 | 27.2±0.6 | 3.8±0.3 | 2.3±0.4 | 4.4±0.6 | 3.3±0.2 | 51.2±1.1 | 45.4±4.4 | 77.4±0.7 |
| P-value | | | 0.714 | 0.721 | 0.799 | 0.007 | 0.299 | 0.41 | 0.397 | 0.108 | 0.96 | 0.293 | 0.881 |

Stepwise Regression between isotopes and biochemical composition of green coffee beans

A stepwise multiple regressions were conducted to evaluate whether stable isotopes and multi-elements were necessary to predict biochemical composition of green coffee beans. The regression analysis between stable isotopes ($\delta^{18}\text{O}$, $\delta^{15}\text{N}$, and $\delta^{13}\text{C}$), %N and %C, and biochemical compounds of green coffee beans showed significant contribution of the model for caffeine, 4,5-DCQA, TCGA, and sucrose composition of green coffee beans (Table 8) indicating significant positive regression weights. All the requested variables entered into the regression equation for all dependant variables at step 1 of the analysis. Thus, only one Model was employed. The model was significantly related to dependant variable $F(5, 30) = 3.147$, $p = 0.021$ for caffeine; $= 5.007$, $p = 0.002$ for 4,5-DCQA; $= 3.292$, $p = 0.017$ for

TCGA; $= 2.672$, $p = 0.041$ for sucrose (Table 9&10), and non-significant, otherwise. For caffeine, 4,5-DCQA, TCGA and sucrose about 34%, 46% 35% and 31% of the variation was accounted for by the model, respectively. For caffeine percent N contributed significant positive weight ($b = 6.604$, $p = 0.002$) and %C contributed significant negative ($b = -0.388$, $p = 0.004$), respectively) for the model. The study showed that all the studied stable isotopes did not contribute significant weight to the model. Moreover, a t-test showed that biochemical compounds in the list 3-CQA, 4-CQA, 5-CQA, FQA, and CFQA were not significantly influenced ($p > 0.05$) by any of the predictors (Table 9&10). The study investigated that there was significant contribution of $\delta^{15}\text{N}$ for variation of 4,5-DCQA ($P = 0.002$), 3,4-DCQA ($P = 0.043$), and TCGA ($P = 0.003$).

Table 8. The Model for regressing biochemical compounds on stable isotopes and %N & %C content of coffee beans

| Dependant variable | Model | R | R ² | Adjusted R ² | SE. of the Estimate | Model P-value (Regression) |
|--------------------|-------|-------|----------------|-------------------------|---------------------|----------------------------|
| Caffeine | 1 | 0.587 | 0.344 | 0.235 | 0.7926 | 0.021 |
| trigonelline | 1 | 0.474 | 0.225 | 0.096 | 3.7218 | 0.155 |
| Sucrose | 1 | 0.555 | 0.308 | 0.193 | 9.8744 | 0.041 |
| 3-CQA | 1 | 0.467 | 0.218 | 0.088 | 0.5349 | 0.171 |
| 3-CQA | 1 | 0.467 | 0.218 | 0.088 | 0.5349 | 0.171 |
| 4-CQA | 1 | 0.361 | 0.130 | -0.015 | 0.5272 | 0.494 |
| 5-CQA | 1 | 0.305 | 0.093 | -0.058 | 1.5686 | 0.689 |
| FQA | 1 | 0.454 | 0.206 | 0.0740 | 0.5917 | 0.202 |
| 4,5-DCQA | 1 | 0.674 | 0.455 | 0.364 | 0.7975 | 0.002 |
| 3,4-DCQA | 1 | 0.448 | 0.201 | 0.067 | 1.6497 | 0.218 |
| CFQA | 1 | 0.428 | 0.183 | 0.047 | 0.9391 | 0.272 |
| TCGA | 1 | 0.595 | 0.354 | 0.247 | 2.6674 | 0.017 |

Table 9. Stepwise Regression of caffeine, trigonelline, sucrose with stable isotopes and %N & %C content of coffee beans

| Dependant variable | Independent variables in the model | Unstandardized Coefficients B | Std. Error | Standardized Coefficients Beta | t | Sig. |
|-----------------------|------------------------------------|-------------------------------|------------|--------------------------------|--------|-------|
| Caffeine | (Constant) | 4.272 | 7.336 | | 0.582 | 0.565 |
| | %N | 6.604 | 1.958 | 1.134 | 3.373 | 0.002 |
| | $\delta^{15}\text{N}$ | -0.044 | 0.124 | -0.058 | -0.352 | 0.728 |
| | %C | -0.388 | 0.122 | -1.071 | -3.166 | 0.004 |
| | $\delta^{13}\text{C}$ | -0.280 | 0.204 | -0.231 | -1.373 | 0.180 |
| trigonelline | $\delta^{18}\text{O}$ | 0.144 | 0.083 | 0.270 | 1.736 | 0.093 |
| | (Constant) | 43.020 | 34.445 | | 1.249 | 0.221 |
| | %N | -6.669 | 9.194 | -0.265 | -0.725 | 0.474 |
| | $\delta^{15}\text{N}$ | 1.247 | 0.583 | 0.386 | 2.139 | 0.041 |
| | %C | 0.278 | 0.575 | .177 | 0.483 | 0.633 |
| Sucrose | $\delta^{13}\text{C}$ | -0.975 | 0.958 | -0.186 | -1.018 | 0.317 |
| | $\delta^{18}\text{O}$ | 0.378 | 0.389 | 0.165 | .972 | 0.339 |
| | (Constant) | -252.614 | 91.388 | | -2.764 | 0.010 |
| | %N | -9.396 | 24.394 | -0.133 | -0.385 | 0.703 |
| | $\delta^{15}\text{N}$ | 0.840 | 1.547 | 0.093 | 0.543 | 0.591 |
| %C | 1.576 | 1.526 | 0.359 | 1.033 | 0.310 | |
| $\delta^{13}\text{C}$ | -7.836 | 2.541 | -0.533 | -3.084 | 0.004 | |
| $\delta^{18}\text{O}$ | 1.117 | 1.032 | 0.173 | 1.082 | 0.288 | |

The stepwise Regression analysis showed significant positive influence of percent N and significant negative effect of percent C on green coffee bean caffeine composition but all the studied stable isotopes did not contribute significant weight to the model. Therefore, the weak association of stable isotopes and biochemical composition may require further investigation to be used as a tool for discriminating green coffee origins. The first function of the Canonical Discriminant Function completely separated coffees grown at 1150 m a.s.l. from those grown both at 1545 and 1802 m a.s.l. An experiment conducted in Central America by Bertrand *et al.* (2006) gave complementary result which discriminated the majority of the samples grown at high elevations from

the samples grown at the lowest elevations.

Most of the time for Arabica coffee (*Coffea arabica* L.) elevation and variety are important indicators of quality in world market. In conclusion, higher $\delta^{15}\text{N}$ and higher $\delta^{13}\text{C}$ values may be useful indicators of lowland and highland coffee samples, respectively. $\delta^{15}\text{N}$ was found to be a significant indicator of variation of 4,5-DCQA; 3,4-DCQA ; and TCGA. Similarly, higher $\delta^{18}\text{O}$ could discriminate highland unshaded coffees. The association of isotopes and biochemical composition of green coffee beans was found to be weak. In fact, for caffeine percent N contributed significant positive weight and %C significant negative weight, respectively for the model. This

implies the study revealed independency of the regressed variables each other. With the exception of caffeine $\delta^{18}\text{O}$, 3-CQA, %N, and %C most variables were useful predictors of altitudinal ranges of coffee growing sites. In fact, 4,5-

DCQA accounted for 94.4% of the variation composition of green coffee beans to classify lowland coffees from those at midland and highland grown coffees.

Table 10. Stepwise Regression of Chlorogenic acid families and stable isotopes and %N & %C content of coffee beans

| Dependant variable | Variables in the model | Unstandardized Coefficients B | Std. Error | Standardized Coefficients Beta | t | Sig. |
|--------------------|------------------------|-------------------------------|------------|--------------------------------|--------|-------|
| 3-CQA | %N | -1.007 | 1.321 | -.280 | -.762 | 0.452 |
| | $\delta^{15}\text{N}$ | .136 | .084 | .294 | 1.621 | 0.116 |
| | %C | .113 | .083 | .503 | 1.364 | 0.183 |
| | $\delta^{13}\text{C}$ | -.063 | .138 | -.084 | -.456 | 0.652 |
| | $\delta^{18}\text{O}$ | .091 | .056 | .278 | 1.636 | 0.112 |
| 4-CQA | %N | -.785 | 1.302 | -.233 | -.602 | 0.551 |
| | $\delta^{15}\text{N}$ | .105 | .083 | .244 | 1.274 | 0.212 |
| | %C | .066 | .081 | .317 | .813 | 0.422 |
| | $\delta^{13}\text{C}$ | .045 | .136 | .064 | .329 | 0.745 |
| | $\delta^{18}\text{O}$ | .077 | .055 | .250 | 1.394 | 0.173 |
| 5-CQA | %N | -1.457 | 3.875 | -.149 | -.376 | 0.710 |
| | $\delta^{15}\text{N}$ | -.079 | .246 | -.063 | -.320 | 0.751 |
| | %C | -.029 | .242 | -.048 | -.121 | 0.905 |
| | $\delta^{13}\text{C}$ | .463 | .404 | .227 | 1.147 | 0.260 |
| | $\delta^{18}\text{O}$ | .005 | .164 | .005 | .028 | 0.978 |
| FQA | %N | .810 | 1.462 | .205 | .554 | 0.584 |
| | $\delta^{15}\text{N}$ | .079 | .093 | .156 | .851 | 0.401 |
| | %C | -.032 | .091 | -.132 | -.355 | 0.725 |
| | $\delta^{13}\text{C}$ | -.267 | .152 | -.325 | -1.757 | 0.089 |
| | $\delta^{18}\text{O}$ | .109 | .062 | .302 | 1.761 | 0.088 |
| 4,5-DCQA | %N | -1.833 | 1.970 | -.285 | -.931 | 0.359 |
| | $\delta^{15}\text{N}$ | .420 | .125 | .509 | 3.358 | 0.002 |
| | %C | .211 | .123 | .529 | 1.715 | 0.097 |
| | $\delta^{13}\text{C}$ | -.418 | .205 | -.312 | -2.037 | 0.051 |
| | $\delta^{18}\text{O}$ | .075 | .083 | .127 | .895 | 0.378 |
| 3,4-DCQA | %N | -.434 | 4.075 | -.040 | -.106 | 0.916 |
| | $\delta^{15}\text{N}$ | .547 | .259 | .388 | 2.114 | 0.043 |
| | %C | -.084 | .255 | -.122 | -.328 | 0.745 |
| | $\delta^{13}\text{C}$ | -.095 | .424 | -.041 | -.223 | 0.825 |
| | $\delta^{18}\text{O}$ | -.033 | .172 | -.033 | -.191 | 0.850 |
| CFQA | %N | 1.294 | 2.320 | .209 | .558 | 0.581 |
| | $\delta^{15}\text{N}$ | .104 | .147 | .131 | .705 | 0.486 |
| | %C | .019 | .145 | .050 | .133 | 0.895 |
| | $\delta^{13}\text{C}$ | -.365 | .242 | -.283 | -1.510 | 0.141 |
| | $\delta^{18}\text{O}$ | .124 | .098 | .220 | 1.265 | 0.216 |

Canonical Discriminant Functions

A stepwise canonical discriminant function analysis of green coffee bean biochemical and stable isotopes and their respective element composition showed that 4,5-DCQA, TCGA, trigonelline, $\delta^{13}\text{C}$, 3,4-DCQA, $\delta^{15}\text{N}$, 4-CQA, sucrose, 5-CQA, FQA, CFQA in descending order of importance significantly ($P \leq 0.05$) contributed for discrimination of altitudinal ranges (Table 11). At each step, the variable that minimizes the overall Wilks' Lambda was entered. About 83.3% of original grouped cases correctly classified. Two out of four functions were sufficient to accommodate the

variation up to 100%. In fact, the first function accommodated 94.4% of the variation which is contributed by 4,5-DCQA composition of green coffee beans (Figure 3). The first function completely separated coffees grown at 1150 m a.s.l. from those grown both at 1545 and 1802 m a.s.l. About 5.6% of the variation was accounted for by function 2 which separated 75% each of coffees grown at 1802 and 1545 m a.s.l. on one hand and about 70% of coffees grown at 1150 from that of 1802 m a.s.l., on the other. Both 4,5-DCQA and 4-CQA composition of green coffee beans contributed for by the variations at function 2.

Table 11. Tests of Equality of Group Means

| Predictor | Wilks' Lambda | F | df1 | df2 | Sig. |
|-----------------------|---------------|--------|-----|-----|-------|
| 4,5-DCQA | 0.442 | 20.824 | 2 | 33 | 0.000 |
| TCGA | 0.596 | 11.18 | 2 | 33 | 0.000 |
| trigonelline | 0.558 | 13.069 | 2 | 33 | 0.000 |
| $\delta^{13}\text{C}$ | 0.637 | 9.415 | 2 | 33 | 0.001 |
| 3,4-DCQA | 0.707 | 6.85 | 2 | 33 | 0.003 |
| $\delta^{15}\text{N}$ | 0.717 | 6.506 | 2 | 33 | 0.004 |
| 4-CQA | 0.732 | 6.043 | 2 | 33 | 0.006 |
| Sucrose | 0.763 | 5.117 | 2 | 33 | 0.012 |
| 5-CQA | 0.784 | 4.533 | 2 | 33 | 0.018 |
| FQA | 0.793 | 4.32 | 2 | 33 | 0.022 |
| CFQA | 0.8 | 4.137 | 2 | 33 | 0.025 |
| Caffeine | 0.861 | 2.674 | 2 | 33 | 0.084 |
| $\delta^{18}\text{O}$ | 0.962 | 0.654 | 2 | 33 | 0.526 |
| 3-CQA | 0.969 | 0.52 | 2 | 33 | 0.599 |
| %N | 0.98 | 0.332 | 2 | 33 | 0.72 |
| %C | 0.988 | 0.195 | 2 | 33 | 0.824 |




Figure 3 Canonical Discriminant Functions

Table 12. Eigenvalue and Wilks' Lambda

| Function | Eigenvalue | % of Variance | Cumulative % | Canonical Correlation | Wilks' Lambda | Chi-square | df | Sig. |
|----------|--------------------|---------------|--------------|-----------------------|---------------|------------|----|------|
| 1 | 6.410 ^a | 94.4 | 94.4 | .930 | .098 | 73.227 | 8 | .000 |
| 2 | .380 ^a | 5.6 | 100.0 | .525 | .725 | 10.138 | 3 | .017 |

a. First 2 canonical discriminant functions were used in the analysis

Acknowledgements

The authors are grateful to the IUC-JU Soil Fertility Project for funding the research. We are especially thankful for the laboratory facilities of Ghent University, Belgium through IUC-JU Soil Fertility Project for providing all facilities required to analyse the coffee bean samples.

References

- Adugnaw Mintesnot. 2014. Association of Arabica Coffee Quality with Geographic Origins in Ethiopia. PhD Dissertation, Haramaya University, Ethiopia.
- Alonso-Salces, R. M., Serra, F., Reniero, F., Berger, A. and K. R. H. 2009. Botanical and Geographical Characterization of Green Coffee (*Coffea arabica* and

- Coffea canephora*): Chemometric Evaluation of Phenolic and Methylxanthine Contents. *J. Agric. Food Chem.* 2009, 57: 4224–4235.
- Bertrand, B., VAAST, P., ALPIZAR, E, ETIENNE, IH., FABRICE D., and CHARMETANT. P.2006. Comparison of bean biochemical composition and beverage quality of Arabica hybrids involving Sudanese-Ethiopian origins with traditional varieties at various elevations in Central America. *Tree Physiology* 26: 1239–1248.
- Bote, A. D. and Paul Struik, C. 2011. Effects of shade on growth, production and quality of coffee (*Coffea arabica*) in Ethiopia. *Journal of Horticulture and Forestry* Vol. 3(11): 336-341.
- Costa Freitas, AM, Parreira, C, Vilas-Boas, L. 2001. The use of an electronic aroma-sensing device to assess coffee differentiation-comparison with SPME gas chromatography-mass spectrometry aroma patterns. *J. Food Comp. Anal.* 14: 513-522.
- Dawson, T. E., Mambelli, S., Plamboeck, A. H., Templer, P. H., and Tu, K.P. 2002. STABLE ISOTOPES IN PLANT ECOLOGY. *Annu. Rev. Ecol. Syst.* 2002. 33:507–59.
- Ehleringer, J. R., Vogel, J. C. 1993. Historical aspects of stable isotopes in plant carbon and water relations. In: Ehleringer, J.R., Hall, A.E., Farquhar, G.D.E. (Eds.), *Stable isotopes and plant carbon–water relations*. Academic Press, San Diego, pp. 9–18.
- Ehleringer, J.R. and Rundel, P.W. 1989. Stable isotopes:History, Units, and Instrumentation. In: *Stable Isotopes in Ecological studies*, Ecological studies ed. by J.R. Ehleringer, , P.W. Rundel, and K.A. Nagy, volume 68:1-15.
- González-Rios, M, Suarez-Quiroz, ML, Boulanger, R, Barel, M, Guyot, B, Guirard, JP, Schrorr-Galindo, S. 2007. Impact of 'ecological' post-harvest processing on coffee aroma: II. Roasted coffee. *J. Food Comp. Anal.* 20: 297-307.
- Likassa Ebisa. 2014. Effect of Dominant Shade Trees on Coffee Production in Manasibu District, West Oromia, Ethiopia. *Sci. Technol. Arts Res. J.*, July-Sep, 3(3): 18-22.
- Martín, M, Pablos, F, González, A, Valdenebro, M, León-Camacho, M. 2001. Fatty acid profiles as discriminant parameters for coffee varieties differentiation. *Talanta*, 54: 291-297.
- Meinzer, F. C., Goldstein, G. and Grantz, D. A. 1990. Carbon Isotope Discrimination in Coffee Genotypes Grown under Limited Water Supply. *Plant Physiol.* 92, 130-135.
- Ministry of Agriculture (MoA). 2003. *Coffee botany, ecology, and nursery management* (Amharic version, 1995), Addis Ababa, Ethiopia. Pp. 183.
- Otsuki, A. 1983. Simultaneous measurements and determinations of stable carbon and nitrogen isotope ratios, and organic carbon and nitrogen contents in

- biological samples by coupling of a small quadruple mass spectrometer and modified carbon-nitrogen elemental analyzer, *Intl. J. Mass Spectrom. Ion Phys.* 48: 343-346.
- Petit, N. 2007. Ethiopia's Coffee Sector: A Bitter or Better Future? *J. Agrarian Change*, 7(2): 225–263.
- Rodrigues, C.I., Maia, R., Miranda, M., Ribeirinho, M., Nogueira, J.M.F., Ma'guas, C. 2009. Stable isotope analysis for green coffee bean: A possible method for geographic origin discrimination. *Journal of Food Composition and Analysis* 22: 463–471.
- Saurer, M, Aellen K, Siegwolf R. 1997. Correlating delta13C and delta18O in cellulose of trees. *Plant Cell Environ.* 20:1543–50
- Serra, F., Guillou, C.G., Reniero, F., Ballarin, L., Cantagallo, M.I., Wieser, M., Iyer, S.S., berger, Ka'roly H. and Vanhaecke, F. 2005. Determination of the geographical origin of green coffee by principal component analysis of carbon, nitrogen and boron stable isotope ratios. *Rapid Commun. Mass Spectrom.* 19: 2111–2115.
- Shibuya, E.K., Sarkis, J. E. S., Negrini-Neto, O., Martinelli, L. A. 2006. Carbon and nitrogen stable isotopes as indicative of geographical origin of marijuana samples seized in the city of Saõ Paulo (Brazil). *Forensic Science International* 167: 8–15.
- Snoeck,D., Zapata,F., and Domenach, Anne-Marie. 2000. Isotopic evidence of the transfer of nitrogen fixed by legumes to coffee trees. *Biotechnol. Agron. Soc. Environ.* 4 (2): 95–100.
- SPSS Inc.2007. SPSS Base 16.0 User's Guide. United States of America.
- Williams K, Richards JH, Caldwell MM. 1991. Effect of competition on stable carbon isotope ratios of two tussock grass species. *Oecologia* 88:148–51.