

ORIGINAL ARTICLE

# A novel *MEFV* gene mutation (A511V) in a Chilean FMF patient

Nejat Akar <sup>a,\*</sup>, Ayşenur Öztürk <sup>a</sup>, Çiğdem Arslan <sup>a</sup>, Ece Akar <sup>a</sup>,  
Silvia Castillo Taucher <sup>b,c</sup>, Cristobal Passalacqua <sup>b</sup>

<sup>a</sup> Faculty of Medicine, Pediatric Genetics Department, Ankara University, Ankara, Turkey

<sup>b</sup> Genetic Section, Medicine Department, Hospital Clínico Universidad de Chile, Chile

<sup>c</sup> Cytogetetics Section, Laboratorio Clinica Alemana, Chile

Received 1 October 2010; accepted 11 December 2010

## KEYWORDS

FMF;  
MEFV;  
Pyrin;  
Mutation;  
A511V

**Abstract** Familial Mediterranean fever (FMF) is an autosomal recessive disease which is characterized by recurrent fever and inflammation of serous membranes. A Chilean FMF patient was investigated for *MEFV* mutations. After DNA extraction, exons 3, 5, 10 and 3'UTR region of *MEFV* gene were analyzed by DNA sequencing while E148Q and R202Q mutations of exon 2 were detected by RFLP. A novel missense mutation, A511V (c.1532C>T, p.Ala511Val), was found in a heterozygous state in exon 5 of *MEFV* gene. Also, R202Q (c.605G>A, p.Arg202Gln) was detected in heterozygous state. R202Q was of clinical value in the diagnosis of FMF when combined with a disease causing mutation. In this patient, A511V was detected in compound heterozygous state with R202Q and this association may play an important role in FMF.

© 2011 Ain Shams University. Production and hosting by Elsevier B.V. All rights reserved.

*Abbreviations:* FMF, familial Mediterranean fever; *MEFV*, familial Mediterranean fever gene

\* Corresponding author. Address: Koru Mah., Uyum Sitesi 606, Sokak 18 Cayyolu, Ankara, Turkey. Tel.: +90 3125956348; fax: +90 3123201433.

E-mail addresses: akar@medicine.ankara.edu.tr, Nejat.Akar@medicine.ankara.edu.tr (N. Akar).

1110-8630 © 2011 Ain Shams University. Production and hosting by Elsevier B.V. All rights reserved.

Peer review under responsibility of Ain Shams University.

doi:10.1016/j.ejmhg.2011.02.010



Production and hosting by Elsevier

## 1. Introduction

Familial Mediterranean fever (FMF) is an autosomal recessive disease which is characterized by recurrent fever and inflammation of serous membranes (peritoneum, synovium and pleura [1]). FMF, having the most frequent occurrence between hereditary periodic diseases, was first identified as a clinic entity by Siegal in 1945 [2]. The most important complication of FMF is amyloidosis. Colchicine has been used in treatment since 1972 [3]. It reduced the frequency and severity of FMF attacks and prevented the development of amyloidosis [4–6].

FMF most commonly occurs in Mediterranean populations especially in Jews, Turks, Arabs and Armenians [1,7]. Although the disease is seen in other populations rarely, it was defined

in Greeks, Italians, Germans, Polishes, Australians, Cubans, Belgians, Brazilians [8], and also Egyptians [9]. The prevalence was reported as 1/256–1/500 in non-Ashkenazi Jews [10] and as 1/1073 in Turks [11]. On the other hand carrier frequency was estimated in North African Jews, Ashkenazi Jews, Armenians and Turks as 1/5, 1/11, 1/7 and 1/5, respectively [10,12,13]. No data are available regarding the proportion of *MEFV* mutation in Chilean FMF patients.

*MEFV* gene, responsible for FMF, in 16p13.3 [14], was cloned by International FMF Consortium and French FMF Consortium independently and indicated that it comprises 10 exons, including 3505 nucleotides and 15 kb length. The protein encoded by *MEFV* was called as pyrin/marenostrin which contains 781 amino acids [15,16]. Its expression in neutrophils and functions in inflammation were shown in studies [17].

After cloning of the gene, M694V, M680I and V726A mutations were the first to be determined in exon 10 [15,16]. To date, more than 180 mutations and/or polymorphisms related to FMF were identified in the *MEFV* gene [18].

A patient from Chile, diagnosed as FMF case according to the Tel-Hashomer criteria was investigated for *MEFV* mutations. The patient had periodic fever, abdominal pain lasting for several years.


## 2. Subjects and methods

Blood sample of the patient was transferred from Chile to Ankara/Turkey. DNA was extracted from peripheral blood lymphocytes according to standard procedures. Exons 3, 5, 10 and 3'UTR were analyzed by DNA sequencing whereas E148Q and R202Q mutations in exon 2 were detected by Restriction Fragment Length Polymorphism (RFLP) method. For polymerase chain reactions (PCR) of exon 2, 3, 5 and 10-3'UTR, F:5'-ATCTTGGGCCCTAAACGTGG-3' and R:5'-TCC-TTCAGGTCCGCAGATGC-3', F:5'-TCTGTGTAAGCAAC-TTGGGTTTG-3' and R:5'-TTGGGAAAATGAAGT AAGGCC-3', F:5'-TATCGCCTCCTGCTCTGGAATC-3' and R:5'-CACTGTGGG TCACCAAGACCAAG-3', F:5'-GAGGTGGACGTTGGAGACAA-3' and R:5'-TCCT CCTGAAATCCATGG-3' primers were used and 658, 480, 526, and 614 bp fragments were amplified, respectively. The amplicons of exon 2 were analyzed using Eco88I and PvuII


restriction endonucleases (Fermentas, Lithuania) for mutations E148Q and R202Q. For sequencing, PCR products were purified using a PCR purification kit (Metis, Turkey) and then analyzed by automated fluorescence DNA sequencing method (CEQ8000XL, Beckman Coulter, USA) [19,20].

## 3. Results

A new variant, the A511V (Alanine–Valine) missense mutation, caused by a C > T transition at nucleotide 1532 in exon 5, which results in an alanine–valine exchange at codon 511, was found in the heterozygous state by sequencing (Fig. 1), and also this mutation was confirmed by RFLP. Amplified 526 bp product digested with HhaI (Fermentas, Lithuania) at 37 °C and subjected to 3% agarose gel electrophoresis. After digestion, in sample of wild type there were 3 fragments (199, 165 and 162 bp) while 4 fragments in heterozygote sample (361, 199, 165 and 162 bp) (Fig. 2). On the other hand, R202Q mutation in exon 2, R314R in exon 3, E474E, Q476Q and D510D polymorphisms in exon 5 were detected in heterozygous state. But no mutation was found after sequencing of other exons and 3'UTR of the *MEFV* gene.



**Figure 2** PCR–RFLP of novel *MEFV* mutation A511V analyzed by agarose gel electrophoresis. PCR products 526 bp of exon 5 were digested by HhaI restriction endonuclease and band profiles were examined in 3% agarose gel. After digestion, 361, 199, 165 and 162 bp fragments were observed in heterozygosity whereas 199, 165 and 162 bp fragments in wild-type. M: X174 DNA-HaeIII Marker, 1–2: Uncut, 3–4: wild-type, 5–6: Heterozygous (these are the same sample).



**Figure 1** Electropherogram of the new A511V *MEFV* variant. (a) Wild-type (b) The C > T transition at nucleotide 1532 in exon 5 converts alanine to valine exchange in codon 511. Also 1530 C > T polymorphism is seen in heterozygous state (nt: nucleotide).

#### 4. Discussion

Molecular analysis of the *MEFV* gene revealed that a novel missense mutation A511V, located in exon 5 of the gene, was present in the heterozygous state. It causes amino acid replacement of alanine to valine. Polymorphisms and FMF causing mutations have been described in exon 5 suggesting the functional importance of this domain [21].

E474E (1422GA), Q476Q (1428AG), and D510D (1530TC) polymorphisms in exon 5 and R314R polymorphism in exon 3 were first defined by Bernot and colleagues (1998) and has been used as markers in haplotype analysis [22,23]. Cazeneuve et al. (1999) showed that these three polymorphisms were associated in cis position, in addition to R501R (1503CT) polymorphism [24]. Also, Aldea and his colleagues (2004) constituted haplotypes including 1422GA, 1428AG and 1530TC polymorphisms [21].

R202Q in exon 2 of *MEFV* gene was first identified by Bernot et al. (1998) and it has been proposed to be a common polymorphism [22]. Additionally, in FMF database, R202Q was indicated as a polymorphism in linkage disequilibrium with M694V [18]. In a study of 26 Greek FMF patients, R202Q homozygosity has been shown in 4 patients but was not detected in 60 healthy controls and statistically significant difference was observed between the two groups. R202Q gene alteration may be a mutation more than a polymorphism and because of the heterozygote controls, dose-dependent effect may occur in homozygous FMF patients [25]. In the following years, homozygosity of R202Q was reported to be associated with the disease and could be a mutation but has not been mentioned the relation with the M694V mutation [26]. Miyoshi et al. (2008) reported E148Q/R202Q compound heterozygosity in a Japanese patient which may be responsible for the disease [27]. Although, R202Q heterozygosity was detected in 163 healthy Japanese controls of 170, homozygosity could not be detected [28]. In our previous study, we found that there is one more haplotype, which was not in linkage disequilibrium with M694V. This result showed that R202Q might be associated with the disease at least in some FMF patients. R202Q homozygosity could not be detected in our control group but, high frequency of heterozygosity showed that it has no effect when it is in heterozygous state. However, when carrying R202Q in compound heterozygous with another disease causing mutation, the clinical spectrum appears [29]. So, this emphasizes importance of R202Q for diagnosis. Also our Chilean patient carries R202Q and A511V in compound heterozygote state. This association may create a defect on protein function and could play an important role in FMF progression.

We were not able to screen A511V mutation in Chilean population. Five hundred sixteen Turkish & Egyptian FMF patients were analyzed for exon 5 of *MEFV* gene by sequencing and A511V was not present. In conclusion, our results show that compound heterozygosity of R202Q and A511V may have a role in expression of the clinical findings. The relevance of the novel mutation with the disease should be confirmed by studying with FMF in large patient groups and by functional studies.

#### References

[1] Sohar E, Gafni J, Pras M, Heller H. Familial Mediterranean fever. A survey of 470 cases and review of the literature. *Am J Med* 1967;43(2):227–53.

[2] Siegal S. Benign paroxysmal peritonitis. *Ann Intern Med* 1945;23:1–21.

[3] Goldfinger SE. Colchicine for familial Mediterranean fever. *N Engl J Med* 1972;287(25):1302.

[4] Dinarello CA, Wolff SM, Goldfinger ES, Dale DC, Alling DW. Colchicine therapy for familial mediterranean fever. A double-blind trial. *N Engl J Med* 1974;291(18):934–7.

[5] Zemer D, Revach M, Pras M, Modan B, Schor S, Sohar E, Gafni J. A controlled trial of colchicine in preventing attacks of familial mediterranean fever. *N Engl J Med* 1974;291(18):932–4.

[6] Zemer D, Pras M, Sohar E, Modan M, Cabili S, Gafni J. Colchicine in the prevention and treatment of the amyloidosis of familial Mediterranean fever. *N Engl J Med* 1986;314(16):1001–5.

[7] Turkish FMF Study Group. Familial Mediterranean Fever (FMF) in Turkey results of a nationwide multicenter study. *Medicine* 2005;84(1):1–11.

[8] Eliakim M, Levy M, Ehrenfeld M. Laboratory examinations. In: Recurrent polyserositis (Familial Mediterranean Fever, periodic disease). Amsterdam, Elsevier, 1981.

[9] Ozturk A, Elsobky E, Elsayed S, Alhodhod M, Akar N. Mutational analysis of *MEFV* gene in Egyptian patients with familial Mediterranean fever. *Turk J Med Sci* 2009;39(2):229–34.

[10] Daniels M, Shohat T, Brenner-Ullman A, Shohat M. Familial Mediterranean fever: high gene frequency among the non-Ashkenazic and Ashkenazic Jewish populations in Israel. *Am J Med Genet* 1995;55(3):311–4.

[11] Ozen S, Karaaslan Y, Ozdemir O, Saatci U, Bakkaloglu A, Koroglu E, Tezcan S. Prevalence of juvenile chronic arthritis and familial Mediterranean fever in Turkey: a field study. *J Rheumatol* 1998;25(12):2445–9.

[12] Rogers DB, Shohat M, Petersen MG, Bickal J, Congleton J, Schwabe AD, Rotter JI. Familial Mediterranean fever in Armenians: autosomal recessive inheritance with high gene frequency. *Am J Med Genet* 1989;34(2):168–72.

[13] Yilmaz E, Ozen S, Balci B, Duzova A, Topaloğlu R, Besbas N, Saatci U, Bakkaloglu A, Ozguc M. Mutation frequency of familial Mediterranean fever and evidence for a high carrier rate in the Turkish population. *Eur J Hum Genet* 2001;9(7):553–5.

[14] Pras E, Aksentijevich I, Gruber L, Balow Jr JE, Prosen L, Dean M, Steinberg AD, Pras M, Kastner DL. Mapping of a gene causing familial Mediterranean fever to the short arm of chromosome 16. *N Engl J Med* 1992;326(23):1509–13.

[15] The French FMF Consortium. A candidate gene for familial Mediterranean fever. *Nat Genet* 1997;17(1):25–31.

[16] The International FMF Consortium. Ancient missense mutations in a new member of the RoRet gene family are likely to cause familial Mediterranean fever. The International FMF Consortium. *Cell* 1997;90(4):797–807.

[17] Kastner DL. Familial Mediterranean fever: the genetics of inflammation: review. *Hosp Pract (Minneapolis)* 1998;33(4):131–4, 139–140, 143–6 passim.

[18] <http://www.fmf.igh.cnrs.fr/ISSAID/infivers/search.php>.

[19] Akar N, Misiroglu M, Yalcinkaya F, Akar E, Cakar N, Tumer N, Akcakus M, Tastan H, Matzner Y. *MEFV* mutations in Turkish patients suffering from familial Mediterranean fever. *Hum Mutat* 2000;15(1):118–9.

[20] Akar N, Akar E, Ozel D, Tekin M, Ekim M, Yalcinkaya F. A note on mutation analysis in familial Mediterranean fever. *Pediatr Nephrol* 2003;18(2):196–7.

[21] Aldea A, Campistol JM, Arostegui IJ, Rius J, Maso M, Vives J, Yague J. A severe autosomal-dominant periodic inflammatory disorder with renal AA amyloidosis and colchicine resistance associated to the *MEFV* H478Y variant in a Spanish kindred: an unusual familial Mediterranean fever phenotype or another *MEFV* -associated periodic inflammatory disorder? *Am J Med Genet A* 2004;124A(1):67–73.

[22] Bernot A, da Silva C, Petit JL, Cruaud C, Caloustian C, Castet V, Ahmed-Arab M, Dross C, Dupont M, Cattan D, Smaoui N,

- Dode C, Pecheux C, Nedelec B, Medaxian J, Rozenbaum M, Rosner I, Delpech M, Grateau G, Demaille J, Weissenbach J, Touitou I. Non-founder mutations in the *MEFV* gene establish this gene as the cause of familial Mediterranean fever (FMF). *Hum Mol Genet* 1998;7(8):1317–25.
- [23] Aksentijevich I, Torosyan Y, Samuels J, Centola M, Pras E, Chae JJ, Oddoux C, Wood G, Azzaro MP, Palumbo G, Giustolisi R, Pras M, Ostrer H, Kastner DL. Mutation and haplotype studies of familial Mediterranean fever reveal new ancestral relationships and evidence for a high carrier frequency with reduced penetrance in the Ashkenazi Jewish population. *Am J Hum Genet* 1999;64(4):949–62.
- [24] Cazeneuve C, Sarkisian T, Pecheux C, Dervichian M, Nedelec B, Reinert P, Ayvazyan A, Kouyoumdjian JC, Ajrapetyan H, Delpech M, Goossens M, Dode C, Grateau G, Amselem S. *MEFV*-Gene analysis in armenian patients with Familial Mediterranean fever: diagnostic value and unfavorable renal prognosis of the M694V homozygous genotype-genetic and therapeutic implications. *Am J Hum Genet* 1999;65(1):88–97.
- [25] Ritis K, Giaglis S, Spathari N, Micheli A, Zonios D, Tzoanopoulos D, Deltas CC, Rafail S, Mean R, Papadopoulos V, Tzioufas AG, Moutsopoulos HM, Kartalis G. Non-isotopic RNase cleavage assay for mutation detection in *MEFV*, the gene responsible for familial Mediterranean fever, in a cohort of Greek patients. *Ann Rheum Dis* 2004;63(4):438–43.
- [26] Giaglis S, Papadopoulos V, Kambas K, Doumas M, Tsironidou V, Rafail S, Kartalis G, Speletas M, Ritis K. *MEFV* alterations and population genetics analysis in a large cohort of Greek patients with familial Mediterranean fever. *Clin Genet* 2007;71(5):458–67.
- [27] Miyoshi T, Yamashita K, Ohno T, Izumi T, Takaori-Kondo A, Sasada M, Uchiyama T. Familial Mediterranean fever gene as a possible modifier of Sweet syndrome with chronic myelogenous leukemia. *Acta Haematol* 2008;120(1):57–62.
- [28] Yamaguchi K, Ikeda K, Ihara K, Takada H, Kusuvara K, Hara T. Lack of association between E148Q *MEFV* variant and Kawasaki disease. *Hum Immunol* 2009;70(6):468–71.
- [29] Ozturk A, Ozcakar B, Ekim M, Akar N. Is *MEFV* gene Arg202Gln (605G > A) A disease-causing mutation? *Turk J Med Sci* 2008;38(3):205–8.