

Original article

Prevalence of food sensitization and helminth infection among primary school children in Ibadan, southwest Nigeria

Background: Food allergy is a problem of the western world, however, cases of food allergy have also been found in developing countries. In Nigeria, the prevalence of food allergy is scarce while the association of food sensitization with helminthiasis is unknown. **Objective:** This study was designed to determine the prevalence of sensitization to food among Nigerian school children and its relation to intestinal helminthiasis. **Methods:** A total of 128 participants were recruited from CAC Primary School, Sango, Ibadan, Nigeria. Skin Prick Test using freshly prepared food allergens was carried out. Stool samples were collected from them to screen for ova of parasites using concentration method. **Results:** A total of 26 (20.3%) out of 128 participants were SPT positive for various food allergens, while 16 (12.5%) had helminthic infection. There was however an equal distribution of helminthic infection among students who were food allergen SPT positive and negative. There was also no association between gender and susceptibility to food allergy. **Conclusion:** Sensitization to food is common in school children, and intestinal helminthiasis seems to have no influence on its prevalence.

Keywords: MPO, inflammatory response, neonates, sepsis.

Ayodele A. Adelokun and Olatunbosun G. Arinola*

Departments of
Chemical Pathology and
Immunology*,
University of Ibadan,
Nigeria.

Correspondence:
Prof. O.G. Arinola
E-mail: drarinolaog64
@ yahoo.com

INTRODUCTION

Food allergy constitutes a growing problem in the western world and its incidence seems to be on the increase.¹ Children experience food allergies more than adults as tolerance develops with increasing age. In developed countries, the prevalence of food allergy in children is 6-10%, while it is 3-4% in adults². In developing countries, such as Nigeria, report on prevalence of food allergy is scarce but study on prevalence of environmental allergy is extensive^{3,4,5,6}. In developed countries, almost all (90%) reported food allergic reactions are caused by cow's milk and dairy products, hen's egg, peanuts, nuts, gluten containing cereals (e.g. wheat, rye, barley), sesame, soybeans, mustard, fish, crustaceans and shellfish^{7,8,9}, hence they are referred to as "priority food allergens". Other less common allergenic foods include legumes, fruit juices (e.g. kiwifruit, apple, grape) and vegetables (e.g., celeriac, carrots and onions)¹⁰; but the types and prevalence of food allergy varies with geographical location.^{11,12,13,14,15}

The Centers for Disease Control and Prevention documented an 18% increase in the prevalence of reported food allergy in children in the United States of America (USA) within a decade, while a comparable observation was made within the same period in China.^{1,16} Foods causing most allergies in

Africa are egg white, peanut, milk, soy, seafoods, pineapple, vegetable oil and pork.^{17,18,19} Antigens present in food induce a systemic non-responsiveness termed 'oral-tolerance'²⁰. However, a failure to develop tolerance or a breakdown in tolerance in certain individuals results in excessive production of food-specific immunoglobulin E (IgE) antibodies. In these individuals, mast cell-bound IgE molecules are present in the gastrointestinal tract. Upon ingestion, food allergens bind to mast cell-bound IgE and cross-link the IgE molecules leading to the release of inflammatory mediators from the mast cell, causing smooth muscle contraction and vasodilatation.²¹

Human intestinal helminthiasis are rampant in the tropics because of favourable climatic, environmental and socio-cultural factors that permit their transmission for greater part of the year²². These factors predispose school-aged children to be at greater risk of helminthiasis.^{23,24} In Nigeria, commonly encountered helminthes are *Ascaris lumbricoides*, *Trichuris trichiura*, hookworm and schistosomes.^{25,26,27} Evidence exists that helminthes modulate the host immune system to guarantee their own survival²⁸ and also induce regulatory responses that dampen the activity of effector cells.²⁹ The modulatory effect of helminthes on the immune

system has led to the exploration of its use as therapy in autoimmune diseases and immune disorders.

Epidemiological studies have shown inverse associations between environmental allergen skin test reaction and various helminthic infections namely *Ascaris lumbricoides*,^{30,31} *Trichuris trichiura*,^{30,32} hookworm,²⁹ and *Schistosoma*.^{33,34} Also, previous studies focused on identifying co-morbid conditions of food allergy such as allergic rhinitis, atopic dermatitis, and asthma in developed countries.^{35,36,37,38} This present study is therefore designed to determine the prevalence of food sensitization in an emerging country – Nigeria and to find out the association between helminth infection, gender and food sensitization.

METHODS

Participants

A total of 128 pupils (60 boys and 68 girls; age range 5 to 13 years) were recruited from primary schools in Ibadan, Southwestern Nigeria. Ethical clearance was obtained from the State Ministry of Health (Oyo State) while permission was obtained from the State Universal Basic Education Board as well as Local Government Universal Basic Education Authority. Parental informed assent was obtained on behalf of each pupil.

Food allergens

The food allergens used were fresh cow's milk, egg, fish, crayfish, groundnut, soya, wheat, pineapple, vegetable and cassava obtained locally from the market. Allergens were aseptically prepared described as follows: Foods with hard consistency (fish, crayfish, groundnut, soya, wheat, vegetable and cassava) were oven-dried, ground and diluted in buffered saline at 1:3 w/v dilution³⁹. Fresh cow's milk and cooked egg white were used directly. All procedures were carried out under sterile conditions.

Skin Prick Test

Skin Prick Test (SPT) was carried out on all participants using the food allergens mentioned above as described by Arinola *et al*⁴⁰. The wheal diameter was measured with a millimeter ruler and wheal diameter greater than 3mm was considered as positive reaction.

Sample Collection and Analysis

Stool sample was collected into a universal container. The stool sample was microscopically examined for ova after concentration method as previously described by Arinola *et al*⁴¹.

Statistical Methods

Statistical analysis was done using statistical package SPSS version 17.0 and association between variables was determined by Chi-square (X^2) at 0.05 level of significance.

RESULTS

Out of the 128 participants, 26 (20.3%) were positive for the tested food allergens of which 16 were females and 10 males. Also, 16 (12.5%) students had helminthic infection, 11 of them were females and 5 were males. There were no gender differences in the prevalence of helminthic infection or food sensitization ($p > 0.05$). (Table 1).

As shown in table 2, a total of 14 (10.9%) students out of 128 had *Ascaris lumbricoides* infection, while 2 (1.6%) had hookworm infection. Five (8.3%) males had *Ascaris lumbricoides* infection while 9 (13.2%) females had *Ascaris lumbricoides* infection. Hookworm infection was seen in only 2 (2.9%) of the females.

In Table 3, from the 16 positive cases of helminth infection, 8 had positive SPT for the tested foods, while the remaining 8 had negative SPTs. There was also an equal distribution of 7 *Ascaris lumbricoides* to 1 hookworm infection in food allergy positive as well as food allergy negative students.

Sensitization to groundnut was the most prevalent food allergy, followed by milk, soybeans, wheat and cassava. There was however no sensitization to fish or vegetables in any of the students. Among the males, the most prevalent food allergens were groundnut, soybeans, wheat and crayfish; followed by milk, cassava, pineapple and egg. On the contrary, among the females, the most prevalent food allergens were groundnut and milk; none of the females was however susceptible to crayfish allergen; as in table 4.

Fourteen school children (6 males and 8 females) had single food allergy, while 12 school children (4 males and 8 females) had multiple food allergies (Table 5).

Table 1. Gender-based prevalence of food allergy and helminth infection among school children.

		Male (n = 60) (46.9%)	Female (n = 68) (53.1%)	Total (n = 128)	X²
Helminth infection	Positive	5 (8.3%)	11 (16.2%)	16 (12.5%)	0.181
	Negative	55 (91.7%)	57 (83.8%)	112 (87.5%)	
Food sensitization	Positive	10 (16.7%)	16 (23.5%)	26 (20.3%)	0.336
	Negative	50 (83.3%)	52 (76.5%)	102 (79.7%)	

Table 2. Types of helminths species in school children.

	Male (n = 60)	Female (n = 68)	Total (n = 128)
<i>Ascaris lumbricoides</i>	5 (8.3%)	9 (13.2%)	14 (10.9%)
Hookworm	0 (0%)	2 (2.9%)	2 (1.6%)
Total	5 (8.3%)	11 (16.2%)	16 (12.5%)

Table 3. Prevalence of food allergy among school children with different species of helminthic infections.

	HELMINTH POSITIVE n = 16		Total
	<i>Ascaris lumbricoides</i>	Hookworm	
Food allergy positive	7 (43.75%)	1 (6.25%)	8 (50%)
Food allergy negative	7 (43.75%)	1 (6.25%)	8 (50%)

Table 4. Prevalence of types of food allergies among both sexes in the school children.

	MALES (n = 60)	FEMALES (n = 68)	TOTAL (n = 128)
Milk	2 (1.6%)	5 (3.9%)	7 (5.5%)
Egg	2 (1.6%)	1 (0.8%)	3 (2.3%)
Fish	0 (0%)	0 (0%)	0 (0%)
Crayfish	3 (2.3%)	0 (0%)	3 (2.3%)
Groundnut	3 (2.3%)	5 (3.9%)	8 (6.3%)
Soybeans	3 (2.3%)	4 (3.1%)	7 (5.5%)
Wheat	3 (2.3%)	3 (2.3%)	6 (4.7%)
Pineapple	2 (1.6%)	3 (2.3%)	5 (3.9%)
Vegetables	0 (0%)	0 (0%)	0 (0%)
Cassava	2 (1.6%)	4 (3.1%)	6 (4.7%)

Table 5. Prevalence of combination types of food sensitization in relation to gender.

	Allergen	Males (n=60)	Females (n=68)	Total (n=128)
Single allergies	Milk	0 (0%)	2 (1.6%)	2 (1.6%)
	Egg	1 (0.8%)	1 (0.8%)	2 (1.6%)
	Crayfish	2 (1.6%)	0 (0%)	2 (1.6%)
	Groundnut	1 (0.8%)	1 (0.8%)	2 (1.6%)
	Soybeans	1 (0.8%)	2 (1.6%)	3 (2.3%)
	Wheat	0 (0%)	1 (0.8%)	1 (0.8%)
	Pineapple	1 (0.8%)	1 (0.8%)	2 (1.6%)
	Sub-total	6 (4.7%)	8 (6.3%)	14 (10.9%)
Multiple allergies	Milk + wheat	1 (0.8%)	0 (0%)	1 (0.8%)
	Milk + cassava	0 (0%)	1 (0.8%)	1 (0.8%)
	Milk + soya	0 (0%)	1 (0.8%)	1 (0.8%)
	Egg + cassava	1 (0.8%)	0 (0%)	1 (0.8%)
	Groundnut + pineapple	0 (0%)	1 (0.8%)	1 (0.8%)
	Groundnut + wheat	0 (0%)	1 (0.8%)	1 (0.8%)
	Groundnut + soya	1 (0.8%)	0 (0%)	1 (0.8%)
	Soya + cassava	0 (0%)	1 (0.8%)	1 (0.8%)
	Wheat + pineapple	0 (0%)	1 (0.8%)	1 (0.8%)
	Milk + groundnut + wheat	0 (0%)	1 (0.8%)	1 (0.8%)
	Groundnut +soya + wheat + cassava	0 (0%)	1 (0.8%)	1 (0.8%)
	Milk + crayfish + groundnut + pineapple + cassava	1 (0.8%)	0 (0%)	1 (0.8%)
		Sub-total	4 (3.1%)	8 (6.3%)
OVERALL TOTAL	OVERALL TOTAL	10 (7.8%)	16 (12.5%)	26 (20.3%)

DISCUSSION

Previous reports showed that the prevalence of food allergy varies from country to country, with a higher prevalence rate found in developed countries^{1,16,42,43}. Reports of food allergy in developing countries are scarce, however, environmental allergies have been studied^{3, 4,5,6,44}. The recent rise in prevalence of food allergy in western countries^{45,46} has necessitated our need to investigate such prevalence in tropical countries. Also, it has been noted that helminthic infections are widespread in tropical countries and the hygiene hypothesis suggests that due to the prevalence of helminthes in tropical countries, allergic diseases might be scarce.

From the present study, we observed that total prevalence of food allergy among the study population is 20.3%. Pawankar *et al*⁴⁷ reported an allergy prevalence of 20-30% in Africa, this was however not restricted to food allergy. In this study, the food with the highest prevalence of reactivity was groundnut (6.3%); followed by soya (5.5%) and milk (5.5%). Others were cassava (4.7%), wheat (4.7%), pineapple (4.7%), pineapple (3.9%), egg (2.3%) and crayfish (2.3%). These food allergens have also been reported in other parts of Africa^{17, 18, 19}. Previous reports in western countries showed that groundnut, milk, soya, egg, seafoods, sesame and wheat are 'priority food

allergens'^{48,49,50,51}. This is in agreement with our present findings.

Osborne *et al*⁵² reported prevalence rates of egg white (16.5%), peanut (8.5%) and cow milk (5.6%) among Australian children. This order of prevalence came in contrast to our observation; however, there was a comparable prevalence rate of cow's milk allergy. This lends support to the report of Ouahidi *et al*¹⁵ that the types and prevalence of food allergy vary with geographical location. Furthermore, in South Africa, Els *et al*⁵³, reported the following prevalence of food allergens: groundnut (9%), egg white (7%), wheat (4%), fish (4%), and milk (3%). These values are relatively higher than what we observed. This might be due to higher level of 'westernization' in South Africa compared to our study location.

Among the male school children, 3 (2.3%) of them each had a positive reactivity to groundnut, wheat, soybeans and crayfish; while 2 (1.6%) reacted positively to milk, egg, pineapple and cassava (Table 4). However, among the female school children, the highest prevalence of positivity was for groundnut and milk (3.9% each), followed by soybeans and cassava (3.1% each), wheat (2.3%), pineapple (2.3%) and egg (0.8%) (Table 4). None of the school children were sensitive to fish or vegetables. There was no gender difference in the prevalence of food sensitization among the school

children. This might be related to similar environmental factors impacting on the type of food sensitivity observed in them.

In this study, 12.5% of the school children had helminthic infection. This is comparable to the findings of Odu *et al*²⁷ who reported a prevalence of 15.7% among school-aged children. The intestinal helminthes observed in our study were *Ascaris lumbricoides* and hookworm. Various studies have demonstrated the presence of other intestinal helminthes – *Ascaris lumbricoides*, *Trichuris trichiura*, hookworm and schistosomes – across Nigeria^{25,26,27}. Other studies in rural areas across Nigeria, among school-aged children showed higher prevalence rates^{54,55}. Jimenez-Gonzalez *et al*⁵⁶ also reported a prevalence of 34.0% among inhabitants of a rural community in Mexico. This shows that rural communities have a higher prevalence of helminthic infections compared with urban communities. Our present study also shows that helminthic infection is not gender-specific as we found no significant difference in the prevalence between the sexes (male- compared with female-school children).

The percentage of present and absent sensitization to food allergens was comparable among the helminth infected children. It may therefore be hypothesized that immunoregulation by helminth infection did not affect response to food allergens. Various studies have reported interaction between helminthic infection and allergy^{28,57,58,59}. Helminths regulate host immune response, in a way that enhances their survival in the host and limits host tissue damage. However, despite strong T-helper 2-type (Th2) polarization of immune response by helminthic infection, chronic helminthic infections do not induce clinical symptoms of allergy. This paradoxical phenomenon was explained by the report of Maizels and Yazdanbakhsh⁶⁰ that chronic helminth infections protect against allergic diseases. It was also added that the hyporesponsiveness induced by helminthiasis is not only directed toward parasite antigens but seems to be extended to ‘bystander’ antigens, such as vaccine antigens or other pathogens⁶¹.

CONCLUSION

The study showed that both sensitization to food and helminth infection are prevalent among school children but helminth infection or gender have no influence on the prevalence of food sensitization.

REFERENCES

1. **BRANUM AM, LUKACS SL.** Food allergy among U.S. children: trends in prevalence and hospitalizations. National Center for Health Statistics Data Brief; 2008.
2. **HERZ U.** Immunological basis and management of food allergy. *J Pediatr Gastroenterol Nutr* 2008; 47:S54–S57
3. **ADANIJO AO, BANDELE EO.** Cockroach hypersensitivity in asthmatics in Lagos, Nigeria. *East Afr Med J* 2000; 77(11):622 – 6 PMID: 12862110
4. **IGE OM, ARINOLA OG, OLUWOLE O, FALADE AG, FALUSI AG, ADEREMI T, ET AL.** Atopy is associated with Asthma in adults living in rural and urban southwestern Nigeria. *J Asthma* 2011; 48(9):894 – 900.
5. **OGUNLADE OA, IGE OM, ARINOLA OG, ONADEKO BO.** Allergen-specific immunoglobulin E (IgE) antibodies and skin test reactivity in patients with asthma in Nigeria. *J Clin Immunol Immunopathol Res* 2012; 4(3): 25-8
6. **OLUWOLE O, ARINOLA OG, FALADE GA, IGE MO, FALUSI GA, ADEREMI T, ET AL.** Allergy sensitization and asthma among 13-14 year old school children in Nigeria. *Afr Health Sci* 2013; 13(1): 144 - 53
7. **VAN DER POEL L, CHEN J, PENAGOS M.** Food allergy epidemic - is it only a western phenomenon? *Curr Opin Allergy Clin Immunol* 2009; 22:121–6.
8. **SICHERER SH, LEUNG DYM.** Advances in allergic skin disease, anaphylaxis, and hypersensitivity reactions to foods, drugs and insects in 2010. *J Allergy Clin Immunol* 2011; 127:326–35.
9. **SPIČÁK V.** Changing face of allergy and allergic diseases. *Alergie* 2010; 12:83–6.
10. **BOYE JI.** Food allergies in developing and emerging economies: need for comprehensive data on prevalence rates. *Clin Transl Allergy* 2012; 2:25
11. **WESTRITSCHNIG K, SIBANDA E, THOMAS W, AUER H, ASP CK H, PITTNER G.** Analysis of the sensitization profile towards allergens in central Africa. *Clin Exp Allergy* 2003; 33:22–7.
12. **OH JW, PYUN BK, CHOUNG JT, AHN KM, KIM CH, SONG SW.** Epidemiological change of atopic dermatitis and food allergy in school-aged children in Korea between 1995 and 2000. *J Korean Med Sc* 2004; 19:716–23.
13. **LUNET N, FALC**

14. **LEUNG TF, YUNG E, WONG YS, LAM GWK, WONG GWK.** Parent-reported adverse food reactions in Hong Kong Chinese pre-schoolers: epidemiology, clinical spectrum and risk factors. *Pediatr Allergy Immunol* 2009; 20:339–46.
15. **QUAHIDI I, AARAB L, DUTAU G.** The effect of thermic and acid treatment on the allergenicity of peanut proteins among the population of the region of Fes-Meknes in Morocco. *Rev Francaise d'Allergol* 2010; 50(1):15–21.
16. **HU Y, CHEN J, LI H.** Comparison of food allergy prevalence among Chinese infants in Chongqing, 2009 versus 1999. *Pediatr Int* 2010 Oct;52(5):820–4
17. **ACHINEWU SC.** Food allergy and its clinical symptoms in Nigeria. *Food Nutr Bull* 1983, 5(3):18–9.
18. **OBENG BB, AMDAH AS, LARBI IA, YAZDANBAKSH M, VAN REE R, BOAKYE DA.** Food allergy in Ghanaian schoolchildren: data on sensitization and reported food allergy. *Int Arch Allergy Immunol* 2010; 155:63–73.
19. **GRAY C, KUNG ST.** Food allergy in South Africa: joining the food allergy epidemic? *Curr Allergy Clin Immunol* 2012; 25(1):24–9.
20. **BERIN MC, SAMPSON HA.** Mucosal immunology of food allergy. *Current Biology* 2013; 23, R389–R400 <http://dx.doi.org/10.1016/j.cub.2013.02.043>
21. **JANEWAY CA JR, TRAVERS P, WALPORT M, CAPUS JD.** *Immunobiology: The Immune System in Health and Disease*. 5th edition. New York: Garland Science; 2001. Effector mechanisms in allergic reactions. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK27112/>
22. **MORDI RM, NGWODO POA.** A study of blood and gastro-intestinal parasites in Edo State. *Afr J Biotechnol* 2007; 6 (19): 2201-7
23. **ALBONICO M, ALLEN H, GHITSULO L, ENGELS D, GABRIELLI AF, SAVIOLI L.** Controlling Soil Transmitted Helminthiasis in Pre-school –Age children through preventive chemotherapy. *PLOS Negl Trop Dis*, 2008; 2(3): e126.
24. **ODU NN, OKONKO IO, ERHI O.** Study of Neglected tropical diseases (NTDs): Gastro Intestinal Helminthes among school children in Port Harcourt, Rivers State, Nigeria. *Report and Opinion*, 2011; 3(9):6-16.
25. **ADEYEBA OA, AKINLABI AM.** Intestinal parasitic infections among school Children in a rural community, southwest Nigeria. *Nig J Parasitol* 2002; 23:11-8
26. **ARINOLA OG, YAQUB S AND RAHAMON SK.** Reduced serum IgE level in Nigerian children with Nigerian children with helminthiasis compared with protozoan infection: Implication on hygiene hypothesis. *Ann Biol Res* 2012; 3 (12):5754-7
27. **ODU NN, ELECHI VI, OKONKO IO.** Prevalence of Intestinal Helminthes Infection among Primary School Children in Urban and Semi-Urban Areas in Rivers State, Nigeria. *World Rural Observations* 2013; 5(1)
28. **SMITS HH, EVERTS B, HARTGERS FC, YAZDANBAKSH M.** Chronic Helminth Infections Protect Against Allergic Diseases by Active Regulatory Processes *Curr Allergy Asthma Rep* 2010 Jan; 10(1): 3–12.doi: 10.1007/s11882-009-0085-3
29. **FLOHR C, TUYEN LN, LEWIS S, QUINNEL R, MINH TT, LIEM HT, ET AL.** Poor sanitation and helminth infection protect against skin sensitization in Vietnamese children: a cross-sectional study. *J Allergy Clin Immunol* 2006; 118:1305–11.
30. **COOPER PJ, CHICO ME, RODRIGUES LG, ORDONEZ M, STRACHAN D, GRIFFIN GE, ET AL.** Reduced risk of atopy among school age children infected with geohelminth parasites in a rural area of the tropics. *J Allergy Clin Immunol* 2003; 111:995–1000.
31. **OBIHARA CG, BEYERS N, GIE RP, HOEKSTRA MO, FINCHAM JE, MARAIS BJ, ET AL.** Respiratory atopic disease, Ascaris-immunoglobulin E and tuberculin testing in urban South African children. *Clin Exp Allergy* 2006; 36:640–8.
32. **RODRIGUES LG, NEWLAND P, GUNHA SS.** Early infections with intestinal helminths reduce the risk of atopy later in childhood. *Clin Exp Allergy* 2008, 38:1769-77
33. **ARAUJO MI, DE CARVALHO EM.** Human schistosomiasis decreases immune responses to allergens and clinical manifestations of asthma. *Chem Immunol Allergy* 2006; 90:29–44.
34. **VAN DEN BIGGELAAR AH, VAN REE R, RODRIGUES LG, LELL B, DEELDER AM, KREMSNER PG, ET AL.** Decreased atopy in children infected with *Schistosoma haematobium*: a role for parasite-induced interleukin-10. *Lancet* 2000; 356:1723–7.
35. **EIGENMANN PA, CALZA AM.** Diagnosis of IgE-mediated food allergy among Swiss children with atopic dermatitis. *Pediatr Allergy Immunol* 2000; 11:95-100.
36. **BRANUM AM, LUKACS SL.** Food allergy among children in the United States. *Pediatrics* 2009; 124:1549-55.
37. **GRAY C.** Allergies in eczema. *Current Allergy Clin Immunol* 2011; 24(3):185-91.

38. **GRAY GL, LEVIN ME, ZAR HJ, PORTER PC, KHUMALO NP, VOLKWYN L ET AL.** Food allergy in South African children with atopic dermatitis. *Paediatr Allergy Immunol* 2014; 25:572-9.
39. **RANCE F, JUCHET A, BREMONT F, DUTAU G.** Correlations between skin prick tests using commercial extracts and fresh foods, specific IgE. and food challenges. *Allergy* 1997; 52: 1031-5.
40. **ARINOLA OG, AFDLABI KA, OLOPADE CO.** Immunological skin tests and hematological indices in Nigerian users of skin lightening creams. *EDOJ* 2011, 7(2):3
41. **ARINOLA GO, OLUWOLE O, OLADOKUN R, ADEDOKUN BO, OLOPADE OI, OLOPADE CO.** Intestinal helminthic infection increases serum levels of IL-2 and decreases serum TGF-beta levels in Nigerian asthmatic patients. *Open J Immunol* 2014; 4: 1 – 8 <http://dx.doi.org/10.4236/oji.2014.41001>
42. **RONA RJ, KEIL T, SUMMERS C, GISLASON D, ZUIDMEER L, SODERGREN E, ET AL.** The prevalence of food allergies: a meta-analysis. *J Allergy Clin Immunol* 2007, 120(3):638–46.
43. **ZUIDMEER L, GOLDBAHN K, RONA RJ, GISLASON D, MADSEN C, SUMMERS C ET AL.** The prevalence of plant food allergies: a systematic review. *J Allergy Clin Immunol* 2008; 121(5):1210–8.
44. **IGE OM, FALADE AG, ARINOLA OG.** Atopy is a risk-factor for adult asthma in urban community of Southwestern Nigeria. *Lung India* 2012, 29(2):114 – 9.
45. **HADLEY G.** Food allergies on the rise? Determining the prevalence of food allergies, and how quickly it is increasing, is the first step in tackling the problem. *EMBO Rep* 2006; 7(11): 1080–3. doi: 10.1038/sj.embor.7400846
46. **LINNEBERG A.** Increase in allergy: a global challenge. *Drugs Today (Barc)*. 2008; 44 (Suppl B):5–10.
47. **PAWANKAR R, BAENA-CAGNANI CE, BOUSQUET J, CRUZ AA, KALINER MA, LANIER BQ, ET AL.** State of World Allergy Report 2008: allergy and chronic respiratory diseases. *World Allergy Org J* 2008;1(Suppl):S4–S17
48. **GOLVER AF, NEVANTAUH H, MACDOUGALL CF, CANT AJ.** Severe food-allergic reactions in children across the UK and Ireland, 1998–2000. *Acta Paediatr* 2005; 94(6):689–95.
49. **UGUZ A, LACK G, PUMPHREY R, EWAN P, WARNER J, DICK J ET AL.** Allergic reactions in the community: a questionnaire survey of members of the anaphylaxis campaign. *Clin Exp Allergy*. 2005; 35(6):746–50.
50. **BOCK SA, MUNOZ-FURLONG A, SAMPSON HA.** Further fatalities caused by anaphylactic reactions to food, 2001–2006. *J Allergy Clin Immunol* 2007; 119(4):1016–8.
51. **JARVINEN KM, SIGHERER SH, SAMPSON HA, NOWAK-WEGRZYN A.** Use of multiple doses of epinephrine in food induced anaphylaxis in children. *J Allergy Clin Immunol* 2008; 122(1):133–8.
52. **OSBORNE NJ, KOPLIN JJ, MARTIN PE, GURRIN LG, LOWE AJ, MATHESON MC ET AL.** HealthNuts Investigators. Prevalence of challenge-proven IgE-mediated food allergy using population-based sampling and predetermined challenge criteria in infants. *J Allergy Clin Immunol* 2011 ; 127(3):668-76.e1-2. doi: 10.1016/j.jaci.2011.01.039
53. **ELS G, BOONZAAIER L, GREEN RJ.** Atopy in asthmatic children attending a tertiary hospital in Pretoria. *Curr Allergy Clin Immunol* 2010; 23:180-2.
54. **AWOLAJU BA, MORENIKEJI OA.** Prevalence and intensity of intestinal parasites in five communities in south-west Nigeria. *Afr J Biotechnol*, 2009; 8 (18):4542-6
55. **CHUKWUMA MC, EKEJINDU IM, AGBAKOBA NR, EZEAGWUNA DA, ANAGHALU IC, NWOBU DC.** The Prevalence and Risk Factors of Geohelminth Infections among Primary School Children in Ebenebe Town, Anambra State, Nigeria. *Middle-East J Sci Res*, 2009; 4(3): 211-5
56. **JIMENEZ-GONZALEZ DE, MARQUEZ-RODRIGUEZ K, RODRIGUEZ JM, GONZALES X, OXFORD J, SANCHEZ R ET AL.** Prevalence and risk factors associated with intestinal parasites in a rural community of central Mexico. *J Parasitol Vector Biol*, 2009; 1(2): 9-12
57. **COOPER PJ.** Interactions between helminth parasites and allergy *Curr Opin Allergy Clin Immunol* 2009; 9(1): 29–37. doi:10.1097/ACI.0b013e32831f44a6.
58. **VON MUTIUS E, VERGELLI D.** Farm living: effects on childhood asthma and allergy. *Nat Rev Immunol* 2010; 10(12):861–8.
59. **RUJENI N, TAYLOR DW, MUTAPI F.** Human Schistosome Infection and Allergic Sensitisation. *J Parasitol Res* 2012; Article ID 154743, doi:10.1155/2012/154743
60. **MAIZELS RM, YAZDANBAKHSH M:** T-cell regulation in helminth parasite infections: implications for inflammatory diseases. *Chem Immunol Allergy* 2008, 94:112–23.
61. **VAN RIET E, RETRA K, ADEBNIKA AA, JOL-VAN DER ZIJDE GM, UH HW, LELL B ET AL.** Cellular and humoral responses to tetanus vaccination in Gabonese children. *Vaccine* 2008, 26:3690–5.