

# Federal Polytechnic Ilaro

## Journal of Pure & Applied Sciences {FEPI- JOPAS}

Volume 3 Issue 2: December 2021, Edition



Published by:

The School of Pure and Applied Science

The Federal Polytechnic Ilaro, Ogun State, Nigeria.

<https://fepi-jopas.federalpolyilaro.edu.ng>

E-mail: [fepi.jopas@federalpolyilaro.edu.ng](mailto:fepi.jopas@federalpolyilaro.edu.ng)

ISSN: 2714-2531

## EDITORIAL TEAM

### A. Editorial Board

NAME	STATUS	AFFILIATION
R.T. Oyede (PhD)	Chairman	The Federal Polytechnic Ilaro, Nigeria
Prof. O.O. Ajani	Member	Covenant University Ota, Nigeria
Prof. H.O.D. Longe	Member	University of Lagos, Akoka, Nigeria
Prof. I.A. Ayodele	Member	University of Ibadan, Nigeria
C.E Ogunlade (PhD)	Member	Federal Polytechnic Ede, Nigeria
O. E. Oduntan(PhD)	Member	The Federal Polytechnic Ilaro, Nigeria

### B. Consulting Editors

NAME	AFFILIATION
Prof. H.O.D. Longe	University of Lagos, Akoka, Nigeria
R. A. Oloyo(PhD)	The Federal Polytechnic Ilaro, Nigeria
L.A. Azeez (Phd)	Osun State University, Oshogbo, Nigeria
A.O. Oyedeji (PhD)	The Federal Polytechnic Ilaro, Nigeria
O. O.Lanloye (PhD)	Bowen University Iwo, Nigeria, Kampala International University, Uganda.
N.O.A. Ilelaboye	The Federal Polytechnic Ilaro, Nigeria
E.O. Fatunmbi (PhD)	The Federal Polytechnic Ilaro, Nigeria
O.J. Adebawale (PhD)	The Federal Polytechnic Ilaro, Nigeria
Engr. I.A. Adeyanju (PhD)	The Federal University, Oye Ekiti, Nigeria
Engr. O.P. Aiyelabowo (PhD)	The Federal Polytechnic Ilaro, Nigeria

### C. Editor-in-Chief

NAME	AFFILIATION
Prof. O.O. Ajani	Covenant University Ota, Nigeria

### D. Managing Editor

NAME	AFFILIATION
O. E. Oduntan (PhD)	The Federal Polytechnic Ilaro, Nigeria

### E. Associate Editors

NAME	AFFILIATION
O.J. Adebawale(PhD)	The Federal Polytechnic Ilaro, Nigeria
O.J. Adegboye	The Federal Polytechnic Ilaro, Nigeria
O. Ogunyemi	The Federal Polytechnic Ilaro, Nigeria
T.O. Ogunseitan	The Federal Polytechnic Ilaro, Nigeria
E. Gabriel	The Federal Polytechnic Ilaro, Nigeria
A. O. Dawodu	The Federal Polytechnic Ilaro, Nigeria

#### Desk Officer:

Ayodele Emmanuel

The Department of Computer Science, The Federal Polytechnic, Ilaro. Ogun State.

#### Site Editors:

Dr. O. E. Oduntan

Omolewa Stephen

## FOREWORD

Compliment of the season to all our contributors, well-wishers and world of Academia in general. I respectfully appreciate and welcome you all to the volume 3 issue 2 of Federal Polytechnic – Journal of Pure and Applied Sciences (FEPI-JOPAS) which is a peer reviewed multi-disciplinary accredited Journal of International repute. It is imperative to re-affirm that FEPI-JOPAS publishes full length research work, short communications, critical reviews and other review articles. In this issue, readers will find a series of manuscripts of top-rated significance in pure and applied sciences, engineering and built environment. This issue is the last of its kind for 2021 calendar year which features findings from basic and applied researches of high societal impacts from the seasoned authors. These articles have been reviewed and packaged for wider readership through the collective efforts of our managing editor, publishing editors, our valuable reviewers and editorial board members.

In this particular issue, you will find that Ilelaboye and Jesusina evaluated the quality of biscuits and chin-chin made from okara enriched plantain-sorghum flour blends. Ojo and Ebisin utilized convolutional neural network for gender classification through facial analysis. Omotayo and Fafioye investigated antimalarial potential of ethyl acetate fraction of *Phyllanthus niruri* while Olubodun and Adetona examined landscaping as a strategy for combating air pollution in Lagos megacity. Buoye and Ojuawo provided imperative dataset on Covid-19 crisis management in Nigeria and Brazil. Obun-Andy and Banjo investigated effective communication as a tool for good governance in Nigeria. Yusuff and co-workers conducted a field survey on fish hatcheries in Yewa South and Yewa North Local Government of Ogun State. Akinlade and co-workers meticulously expatiated on the effect of aqueous blend of three herbs on haemato-biochemical indices of broiler chicken at starter phase. Ajeigbe, Sangosina, Ogunseitan, Lawal, & Yusuff analysed the Effects of Neem Leaves (*Azadirachta Indica*) and Cassava Peels on the Performance of West African Dwarf Goat. Abdussalam & Adewole in their paper carefully explained the Formulation of Natural Products Repellents for the Control of Cockroaches (*Periplaneta americana*). Elesin & Obafunmiso gave as Assessment of Public Toilets Facilities Provision and Management in Tertiary Institutions in Nigeria- An Overview of The Federal Polytechnic, Ilaro, Ogun State.

I would like to deeply appreciate and extend my profound gratitude to my co-editors, editorial board members, reviewers, members of FEPI-JOPAS, especially the Managing Editor, as well as all the contributing authors for making the production and publishing of this volume 3 issue 2 a reality. I will like to appreciate the authors in this issue for allowing their works to be subjected to our thorough and rigorous peer-review processes and for taking all the constructive criticism in good fate. The authors are solely responsible for the information, date and authenticity of data provided in their articles submitted for publication in the Federal Polytechnic Ilaro – Journal of Pure and Applied Sciences (FEPI-JOPAS). I am looking forward to receiving your manuscripts for the subsequent publications.

You can visit our website (<https://fepi-jopas.federalpolyilaro.edu.ng>) for more information, or contact us via e-mail us at [fepi.jopas@federalpolyilaro.edu.ng](mailto:fepi.jopas@federalpolyilaro.edu.ng).

Thank you and best regards.

Prof. Olayinka O. AJANI

**FEPI-JOPAS VOLUME 3 ISSUE 2 TABLE OF CONTENTS**

Serial No	Paper Title and Author(s)	Page
01.	<p align="center"><b>Physicochemical Evaluation and Pasting Properties of Flours, Biscuit And Chinchin Prepared From Okara Fortified Plantain – Sorghum Blends.</b></p> <p align="center">Ilelaboye N.O. and *Jesusina T.I. Department of Science Laboratory Technology, Federal Polytechnic, Ilaro, Ogun State Nigeria. <a href="mailto:titilayo.jesusina@federalpolyilaro.edu.ng">titilayo.jesusina@federalpolyilaro.edu.ng</a></p>	1-13
02.	<p align="center"><b>Framework Model of Facial Analysis for Gender Classification Using Convolutional Neural Network</b></p> <p align="center"><b>Ojo, A. I., &amp; Ebisin, A. F.</b></p> <p align="center"><i>Department, of Computer Science Ogun State Institute of Technology, Igbesa, Ogun State.</i></p> <p align="center">✉ <a href="mailto:ronkujoyme@yahoo.co.uk">ronkujoyme@yahoo.co.uk</a> , <a href="mailto:ebironke16@gmail.com">ebironke16@gmail.com</a></p>	14-26
03.	<p align="center"><b>A Survey of Fish Hatcheries in Yewa South and Yewa North Local Government Areas of Ogun State, Nigeria</b></p> <p align="center"><sup>1</sup>Yusuff, K. O. <sup>1</sup>Ibidapo-Obe, E. O. and <sup>1</sup>Sangosina, M. I. <sup>1</sup>Department of Agricultural Technology, School of Pure and Applied Sciences, The Federal Polytechnic, P.M.B. 50 Ilaro, Ogun state, Nigeria. Corresponding author : <a href="mailto:khadijah.yusuff@federalpolyilaro.edu.ng">khadijah.yusuff@federalpolyilaro.edu.ng</a>,</p>	27-33
04.	<p align="center"><b>Effect of Aqueous Blend of Three Herbs on Haematobiochemical Indices of Broiler Chicken at Starter Phase</b></p> <p align="center"><b>Akinlade, O. O.,<sup>1*</sup> Okusanya, P. O. and Okparavero, O. O.</b></p> <p align="center"><sup>1</sup>Department of Agricultural Technology, School of Pure and Applied Sciences, The Federal Polytechnic, P.M.B. 50 Ilaro, Ogun state, Nigeria. ✉ <a href="mailto:olamilekan.akinlade@federalpolyilaro.edu.ng">olamilekan.akinlade@federalpolyilaro.edu.ng</a></p>	34-39
05.	<p align="center"><b>Effective Communication as a Tool for Good Governance in Nigeria</b></p> <p align="center"><b>Obun-Andy, M. &amp; Banjo, A. O.</b></p> <p align="center"><i>Department of Mass Communication, Federal Polytechnic, Ilaro, Ogun State.</i> ✉ <a href="mailto:maria.obunandy@federalpolyilaro.edu.ng">maria.obunandy@federalpolyilaro.edu.ng</a></p>	40-44
06.	<p align="center"><b>In Vivo Antiplasmodial Effect of the Ethyl Acetate</b></p>	45-48

	<p align="center"><b>Fraction of Crude Extract of Phyllanthus Niruri.</b></p> <p align="center"><b>Omotayo, S. O., &amp; Fafioye, A. O.</b></p> <p align="center"><i>Department of Science Laboratory Technology Federal Polytechnic Ilaro, Ogun State.</i></p> <p align="center"><a href="mailto:olakunle.omotayo@federalpolyilaro.edu.ng">✉olakunle.omotayo@federalpolyilaro.edu.ng</a></p>	
07.	<p align="center"><b>Covid-19 Crisis Management in Nigeria and Brazil</b></p> <p align="center"><b>Buoye P. A* . and Ojuawo O. O.</b></p> <p align="center">Department of Computer Science , The Federal Polytechnic, Ilaro.</p> <p align="center">*<a href="mailto:adewuyi.buoye@federalpolyilaro.edu.ng">adewuyi.buoye@federalpolyilaro.edu.ng</a></p>	49-65
08.	<p align="center"><b>Assessment of Public Toilets Facilities Provision and Management in Tertiary Institutions in Nigeria- An Overview of The Federal Polytechnic, Ilaro, Ogun State.</b></p> <p align="center"><b>Elesin, O.G<sup>1</sup> and Obafunmiso, C.K<sup>2</sup></b></p> <p align="center"><sup>1</sup>Department of Urban and Regional Planning, The Federal Polytechnic, Ilaro. <a href="mailto:olanrewaju.elesin@federalpolyilaro.edu.ng">olanrewaju.elesin@federalpolyilaro.edu.ng</a>; <a href="mailto:princealesh4real@gmail.com">princealesh4real@gmail.com</a>.</p> <p align="center"><sup>2</sup>Department of Library and Information Science, The Federal Polytechnic, Ilaro.</p> <p align="center"><a href="mailto:christianah.obafunmiso@federalpolyilaro.edu.ng">christianah.obafunmiso@federalpolyilaro.edu.ng</a>; 08038559401;</p>	66-72
09.	<p align="center"><b>Effects Of Neem Leaves (<i>Azadirachta Indica</i>) and Cassava Peels On The Performance Of West Afr Ican Dware Goat</b></p> <p align="center"><b>Ajeigbe, O.M.. Sangosina,M.I. Ogunseitan, T. O. Lawal,R A. and Yusuff.K.O</b></p> <p align="center">Department of Computer Science, Gateway (ICT) Polytechnic Saapade-Remo, Ogun State.</p> <p align="center"><a href="mailto:moruf.sangosina@federalpolyilaro.edu.ng">✉moruf.sangosina@federalpolyilaro.edu.ng</a></p>	73-79
10.	<p align="center"><b>Formulation of Natural Products Repellents for the Control of Cockroaches (<i>Periplaneta americana</i>)</b></p>	80-83

***Abdussalam, F., & Adewole, A.***

*Department of Science Laboratory Technology*

*Federal Polytechnic Ilaro, Ogun State.*

✉ [olugbenga.adetona@federalpolyilaro.edu.ng](mailto:olugbenga.adetona@federalpolyilaro.edu.ng)

<https://fepi-jopas.federalpolyilaro.edu.ng>

Experimental

## In Vivo Antiplasmodial Effect of the Ethyl Acetate Fraction of Crude Extract of *Phyllanthus Niruri*.

Omotayo, S. O., & Fafioye, A. O.

Department of Science Laboratory Technology Federal Polytechnic Ilaro, Ogun State.

✉ [olakunle.omotayo@federalpolyilaro.edu.ng](mailto:olakunle.omotayo@federalpolyilaro.edu.ng)

### Abstract

This research was designed to study the effect of ethyl acetate fraction of *Phyllanthus niruri* on *Plasmodium berghei* infected mice. *P. niruri* is known to have antimalaria properties. Crude extract was fractionated with ethyl acetate. 50, 100, 200 and 400mg/kg body weight of the ethyl acetate fraction was dosed to albino mice infected with *Plasmodium berghei*. The Plasmodium count, packed cell volume and body weight were measured before, during and after the experiment. Result showed that all doses of the extract exhibited parasitemia inhibition at 88.88%, 85.23%, 79.68%, and 82.54 % which was not statistically significant ( $P < 0.05$ ) when compared with the positive control chloroquine of (90.37%). The Packed cell volume was observed to reduce across group after treatment when compared to before the treatment. However, weight gain was observed in treated groups. Result shows that the ethyl acetate fraction of *P. niruri* has a curative effect on *Plasmodium berghei* and can be utilized to treat malaria after further purification to isolate the chemical responsible for the activity.

**Keywords; Malaria, Ethyl acetate, Curative, Phyllanthus**

### INTRODUCTION

Malaria parasites have killed and sickened more people than any other eukaryotic illness (Chinsembu, 2015). Malaria caused around 411 000 deaths and 229 million infections worldwide in 2019, with children under the age of five continuing to be the most vulnerable population. (WHO, 2020). Despite the fact that malaria is curable, it continues to have a devastating impact on mankind. The poor and underprivileged are disproportionately afflicted by the disease since they have inadequate access to health care and cannot pay for the prescribed treatment in most nations. (WHO, 2020).

Artemisinin combination therapy has significantly aided in the reduction of malaria mortality. However, treatment failures and parasite resistance, as well as the fact that current antimalarial drugs are costly in disadvantaged areas, underscore the need for safe and affordable antimalarial therapies (Fairhurst & Dondorp, 2016).

Natural products play an important role in the treatment of malaria, as a source of lead compounds for the creation of new and effective antimalarial medications. (Boyom et al, 2011). Plant extractions are still used in Nigerian daily life to combat malaria and a range of other ailments, especially in areas where Western medications are unavailable.

The extensive and diverse flora of Nigeria is a potential source of powerful antiplasmodial natural compounds, and ethnopharmacological techniques to their study might be useful in the fight against malaria. In the exploration for new natural plant derived antimalaria, *in vitro* and *in vivo* antimalaria activity testing are required at the early stages.

*Phyllanthus niruri* is also known as kidney stone crusher (Micali et al., 2006). *P. niruri* is used to treat problems related to the gastrointestinal tracts (Karuna, Reddy, Baskar, & Saralakumari, 2009). *P. niruri* was also reported to display anticarcinogenic (Rajeshkumar et al., 2002), hepatoprotective (Amin, Alshawsh, Kassim, Ali, & Abdulla, 2013), and anti-inflammatory (Obidike, Salawu, Ndukuba, Okoli, & Osunkwo, 2010). *In vivo* and *in vitro* research have demonstrated that this herbal extract has antidiabetic (Okoli, Obidike, Ezike, Akah, & Salawu, 2011) and antioxidant effects (Colpo et al., 2014). In a bid to identify the active component of *P. niruri*, we investigated the effects of the ethyl acetate fraction of *P. niruri* plant extract on *Plasmodium berghei*-infected mice.

### MATERIALS AND METHOD

#### Plant Collection and Identification

*Phyllanthus niruri* whole plant were collected from Bosso, Nigeria and was identified by a botanist at the Department of Biological Sciences, F. U. T. Minna.



**Experimental Animals**

For this investigation, 18 albino mice weighing 24-30g purchased from the National Institute for Pharmaceutical Research and Development Abuja, Nigeria. They were fed a conventional meal and given water, and they were kept in standard circumstances.

**Extract Preparation**

The plant was thoroughly washed, dried, and ground into powder, and was soaked in 80% methanol for 24hours. This was filtered through Whatman filter paper grade 1. The filtrate was evaporated with a rotary evaporator and concentrated using a water bath. The crude extract was placed into a separating funnel, and dissolved in 50ml of distilled water. In order of polarity, n-hexane, chloroform, and Ethyl Acetate were used to partition the methanol extract. The Ethyl Acetate fraction was collected and used for this investigation.

**Parasite Inoculation**

*Plasmodium berghei* was received from the National Institute for Pharmaceutical Research and Development, Idu, Abuja, Nigeria, where it was kept alive in mice for a week. On day 0, each mouse was given 0.2ml of contaminated blood containing *P. berghei* parasitised red blood cells intraperitoneally. In addition, the freshly injected animals were checked on a daily basis for parasitemia.

**Evaluation of the Curative Effect**

Six groups of three mice were formed from the eighteen animals. To collect the blood, a mouse infected with *P. berghei* was anaesthetised with chloroform and slaughtered. The blood was diluted with normal saline to provide a volume of 0.2ml. Each of the eighteen mice was given 0.2ml diluted blood intraperitoneally. The extract was given orally once daily for six days after 72 hours of inoculation at dosage levels of 50, 100, 200, and 400mg/kgbw, respectively (D0, D1, D2, D3, D4 and D5). In the fifth group, a similar test with Chloroquine (5 mg/kgbw) acts as a positive control. The sixth group was not given any treatment and functioned as a control group. Thin films were produced from D0-D5 tail blood, treated with methanol, and stained for 20 minutes with 4 percent Giemsa (PH 7.2) before being studied under the microscope. On each slide, five fields were inspected, and the number of infected red blood cells (RBC) was counted and the mean was calculated. (Kabiru et al., 2013).

%Parasitemia Inhibition =

A= Basal Parasitemia count,  
B = Day 5 Parasitemia count

**Estimation of Packed Cell volume.**

By capillary action, blood was collected from each mouse's tail with a capillary tube. One end of the tube was plastestrine-sealed. After that, the closed capillary

tubes were placed on the hematocrit centrifuge and spun for five minutes at 1200 rev per min. The PCV was measured and recorded using a hematocrit reader.

**Data Analysis**


The data was analysed using Graph Pad Prism6. Data are presented as mean ± SD, One-way Anova were carried out to determine significant difference in parasitemia count across groups.

**RESULT AND DISCUSSION**


**Table 1: Curative test result of ethyl acetate fraction of *P. niruri*.**

Group	Treatment	Basal Parasitemia	DAY5 Parasitemia	% Parasitemia inhibition
1	50 mg/kg PnEtF	12.6 ±2.27	1.4 ±0.53*	88.88
2	100 mg/kg PnEtF	11.5 ±1.60	1.7 ±1.13*	85.23
3	200 mg/kg PnEtF	12.3 ±2.31	2.5 ±0.50*	79.68
4	400 mg/kg PnEtF	12.6 ±3.86	2.2 ±0.28*	82.54
5	5 mg/kg Chq	13.5 ±0.83	1.3 ±1.33*	90.37
6	NC	11.3 ±1.17	12.3 ±0.31	-8.85

\* There is significant difference at P < 0.05 Compared to Negative Control group




**Figure 1: Effect of ethyl acetate fraction of *P.niruri* on the parasitemia of infected mice.**



**Figure 2: Effect of ethyl acetate fraction of *P.niruri* on the weight.**

Data Presented as Mean  $\pm$  SD, n=3  
 \* There is significant difference at  $P < 0.05$  Between Day 0 and Day 5

Int(Infected not treated), Chq(Chloroquine)



**Figure 3: Effect of ethyl acetate fraction of *P.niruri* on Packed cell volume.**

Data Presented as Mean  $\pm$  SD, n=3  
 \* There is significant difference at  $P < 0.05$  Between Day 0 and Day 5

Int(Infected not treated), Chq(Chloroquine).

### DISCUSSION

The phytochemicals found in the crude extract of *P. niruri* in a literature study indicate that it has pharmacological activities (Kabiru et al.,2013). The mice infected with *plasmodium burgei* and dosed with the ethyl acetate fraction from the result in Fig 1 showed a strong curative effect when compared with the positive control group treated with chloroquine.

The effects of ethyl acetate extracts on percent PCV (percentage Packed Cell Volume) after therapy was found to decrease though not statistically significant, these corroborates the findings of (Kabiru et al., 2013 and Nardos and Makonnen, 2017). This is a common occurrence owing to the extract's action on infected blood cells, particularly red blood cells (RBCs). Days following therapy, the cells continue to progressively divide, and this situation is generally transient.

The impact of the ethyl acetate extracts on weight was also considered, and it was discovered that, as shown in figure 3, there was weight gain after treatment, and progressively increased days following treatment. The Initial weight loss might be due to the disease's (malaria) signs and symptoms, which include weight loss. The plant extract however during and after treatment improved on the weight.

### CONCLUSION

The ethyl acetate fraction of *P. niruri* has a curative effect on *Plasmodium burgei* and can be utilized to treat malaria after further purification to isolate the active compound responsible for the activity.

It is also recommended that the ethyl acetate fraction be subjected to further fractionation, such as vacuum liquid chromatography and column chromatography, in order to better exploit the active components in the fraction, increase yields, and serve as templates for the

development of therapeutically more effective antimalarial drugs.

## REFERENCES

- Amin, Z., Alshawsh, M., Kassim, M., Ali, H., & Abdulla, M.A. (2013). Gene expression profiling reveals underlying molecular mechanism of hepatoprotective effect of *Phyllanthus niruri* on thioacetamide-induced hepatotoxicity in Sprague Dawley rats. *BMC Complementary and Alternative Medicine*.13(1)160.
- Boyom, F., Fokou, P., Yamthe L., Mfopa, A., Kemgne, E., Mbacham, W., ... Rosenthal, P. (2011). Potent antiplasmodial extracts from Cameroonian Annonaceae. *Journal of Ethnopharmacology*, 134(3)717–724.
- Chinsembu, K. (2015). Plants as antimalarial agents in Sub-Saharan Africa. *Acta Tropica*. 152,32–48.
- Colpo, E., Vilanova, C., Pereira, R., Reetz, L., Oliveira, L., Farias, I., ... Rocha, J. (2014). "Antioxidant effects of *Phyllanthus niruri* tea on healthy subjects," *Asian Pacific Journal of Tropical Medicine*, 7(2)113–118.
- Fairhurst, R. & Dondorp, A. (2016). Artemisinin-resistant *Plasmodium falciparum* malaria. *Microbiology Spectrum*. 4(3).
- Kabiru, Y., Abdulkadir, A. Gbodi, T., Bello, U., Makun, H., Amah D., & Ogbadoyi, E. (2013). Evaluation of Haematological Changes in *Plasmodium-berghei*-infected Mice Administered with Aqueous Extract of *Phyllanthus amarus*. *Pakistan Journal of Biological Sciences*, 16 510-516.
- Karuna, R., Reddy, S., Baskar, R., & Saralakumari, D. (2009). Antioxidant potential of aqueous extract of *Phyllanthus amarus* in rats. *Indian Journal of Pharmacology*, 41(2)64–67.
- Micali, S., Sighinolfi, M., Celia, A., Stefani, S., Grande, M., Cicero, A., & Bianchi, G. (2006). Can *Phyllanthus niruri* affect the efficacy of extracorporeal shock wave lithotripsy for renal stones? A randomized, prospective, long-term study. *The Journal of Urology*. 176(3)1020–1022.
- Nardos, A. & Makonnen, E. (2017). In vivo antiplasmodial activity and toxicological assessment of hydroethanolic crude extract of *Ajuga remota*. *Malaria Journal*, 16(1)25.
- Obidike, I., Salawu, O., Ndukuba, M., Okoli, C., & Osunkwo, U. (2010). The anti-inflammatory and antinociceptive properties of the chloroform fraction from *Phyllanthus niruri* plant is mediated via the peripheral nervous system," *Journal of Dietary Supplements*. 7(4)341–350.
- Okoli, C., Obidike, I., Ezike, A., Akah, P., & Salawu, O. (2011). Studies on the possible mechanisms of antidiabetic activity of extract of aerial parts of *Phyllanthus niruri*, *Pharmaceutical Biology*. 49(3)248–255.
- Rajeshkumar, N., Joy, K., Kuttan, G., Ramsewak, R., Nair, M., & Kuttan, R. (2002). Antitumour and anticarcinogenic activity of *Phyllanthus amarus* extract, *Journal of Ethnopharmacology*, 81(1)17–22.
- World Health Organization, Fact sheet about malaria, WHO, Geneva, 2020.