

## THE EFFECTS OF NITROGEN RATES ON GROWTH AND YIELD OF FIVE VARIETIES OF SWAMP RICE (*Oryza sativa* Linn.) IN INI, SOUTH EASTERN NIGERIA.

N. E. NNEKE and B. A. NDON

(Received 20 May 2002; Revision accepted 5 August 2003)

### ABSTRACT

A field experiment was conducted in 1998 and 1999 in Ini local Government Area of Akwa Ibom State, Nigeria; mainly to determine the optimum nitrogen fertilizer rate for the area as well as select the rice varieties that responded optimally. A split plot design was used with five N rates {0, 50, 100, 150 and 200kg/ha} as the sub plot treatments and five rice varieties {Cisadane, Suakoko 8, IR 5, ITA 306 and MAS 2401} as the main plot treatments. Nitrogen was band applied as urea in two equal splits at 2 and 8 weeks after transplanting {WAT}, respectively. The results showed that the grain yield and growth parameters of the rice varieties were significantly { $P < 0.05$ } increased by the various N rates. In particular, the rate of 150 kg/ha was consistently superior to other rates in increasing yields. It produced the broadest leaf area and the highest number of spikes per panicle in 1999 and the highest number of tillers per plant and the heaviest grain yield in both years. Among the varieties, Suakoko 8 gave the highest significant { $P < 0.05$ } yield response to the applied N fertilizer, followed by IR 5 and MAS 2401 {no significant difference}, ITA 306 and Cisadane, in that order. Application of 150 kg/ha of N in the form of urea to Suakoko 8 variety is recommended for the area.

**KEY WORDS:** Nitrogen rates, rice varieties, effects, growth and yield.

### INTRODUCTION

Rice yields are generally low in Ini Local Government Area of Akwa Ibom State. This is attributable to the cultivation of low yielding varieties as well as poor soil fertility. Studies conducted by the Northern Akwa Ibom Swamp Resources Development Project NAKSRDP {1994}, indicated that seven different rice varieties are currently cultivated in Ini, the study area. This calls for a proper evaluation of the yield potentials of the various varieties.

Besides, for the past twenty years, wetland rice paddies in the study area have depended entirely on nutrients supplied from the soil and flood / irrigation water which are now becoming increasingly inadequate to sustain good growth and yield in a continuous cropping system. Opukiri *et al* {1991} observed that nitrogen is the most limiting plant nutrient for rice production in the tropics. This is because soil applied N is often lost through various transformation pathways such as volatilization, denitrification and leaching {De Data, 1987} and the fact that soils of Southern Nigeria are generally low in total N content, usually 0.02-1.16% {Moorman, 1980}. When applied in adequate amount, N encourages vegetative growth. It accounts for the plumpness of the grains and stimulate tillering and panicle formation {Foth, 1978}.

Opukiri *et al* (1991) have reported that with increase in the rate of applied N fertilizer, parameters such as number of spikelets per panicle, total dry matter and grain yield also increased significantly. However, beyond a certain rate considered as the optimum, no significant increases were observed.

Grist {1986} recommended 30-35 kg N/ha for the indica sub-species of rice and 100 kgN/ha for the japonica subspecies. Besides, Oyedokun and Sobulo {1979} reported that 30kgN/ha was the optimum N rate for Western Nigeria, while Sobulo {1982} recommended 50kgN/ha for the Savannah region. These blanket recommendations are considered inadequate considering the wide disparity in the soil and climatic conditions of the production areas {Grist, 1986} and the fact that rice varieties differ widely in their response to N fertilizer {Purselove, 1975}.

It is obvious that the optimum N rate for rice growing areas could well be determined by conducting experiments in specific areas where specific rice varieties are grown. For instance, Opukiri *et al* {1991} have recommended that to obtain satisfactory grain yields, fertilization with urea at 90kgN/ha was adequate for ITA 303 and FARO 11 under the climate and soil of Port Harcourt in River State.

Amidst these conflicting recommendations, this study was therefore, under taken in order to determine the optimum N fertilizer rate for Ini soils as well as select the varieties that responded optimally to the rate.

### MATERIALS AND METHODS

Field experiments were conducted in 1998 and 1999 in Ini Local Government Area {Latitudes 5° 03' to 5° 27' North and Longitudes 7° 39' to 7° 56' East} located at the northern part of Akwa Ibom State in Southeastern Nigeria {NAKSRDP, 1994}. The average annual rainfall in this area is about 2000mm and falls between March and November with peaks in bimodal pattern. The mean monthly sunshine is 3 hours 31 minutes and mean diurnal temperature is 30°C {Okoji, 1995}.

The treatments consisted of five swamp rice varieties {Cisadane, Suakoko 8, IR5, ITA 306 and MAS 2401} and five levels of N {0, 50, 100, 150 and 200kgN/ha}. The experiment was laid out in a split-plot design with the varieties in the main plot treatments and N levels in the sub plot treatments, respectively.

Soil Samples were taken at the depth of 0-15cm before each planting and analysed for both physical and chemical properties. Soil pH was determined by electrometric method {Bates, 1954}; particle size by the hydrometer method {Bouyoucos, 1951}; organic matter by the wet oxidation method of Walkley and Black {1934}; total N by Macro Kjeldahl method {Jackson, 1962}; and available phosphorus by the method of Bray P-1 {Bray and Kurtz, 1954}, K and Na were determined by flame analyzer and Mg and Ca by EDTA titration method {Jackson, 1962}. Exchange acidity was extracted with 1N KCl and the acidity determined by titrating with NaOH {ITA, 1979}.

Before planting, the seeds were dressed with Aldrex T in

order to check the incidence of brown spot disease which occurred commonly in the area. Transplanting was done in rows at 4 WAP {weeks after planting} at a spacing of 30cmx30cm. Two seedlings were planted per hill to achieve a plant density of 222,220 plants per hectare.

Nitrogen fertilizer was band applied as urea in two equal splits at 2 and 8 WAT. To prevent any possible nutrient drift, bunds of 30cm height were built around the various subplots. Weeding which combined hoe weeding and hand pulling, were done three times {4, 8 and 12 WAT} before harvest. Water level was maintained at a depth of about 15cm throughout the growing season except during weeding and fertilizer application (zero depth). Birds constituted the major pest of the crop from flowering and were controlled by erecting scare crows and employing the services of a "bird boy".

Twenty randomly chosen and tagged plants per subplot were used for the measurement of growth parameters. Measurement began at 4 WAT and continued every two weeks until grain harvest.

To obtain plant leaf area, the leaf length and the widest part of selected leaves were measured with a metre rule and the product obtained was multiplied by 0.75 (Hammer, 1980). The number of tillers per plant was obtained by counting the total number of tillers produced by each plant. Similarly, the number of spikes per panicle was obtained by counting the total number of spikes borne by each panicle.

At 75% browning, the panicles were harvested from a 1.5m x 0.9m area using a knife and thereafter threshed and sundried to about 14% moisture content (De Datta *et al.*, 1987) to obtain the grain yield.

Data collected were summarized and subjected to Analysis of Variance; while the Least Significant Difference Test ( $P < 0.05$ ) was used to compare treatment means (Gomez and Gomez, 1984).

**Table 1: Some physical and chemical characteristics of the soil used for the study**

Soil properties	Years	
	1998	1999
<b>Physical Characteristics</b>		
Sand (%)		
Silt (%)	13.60	13.50
Clay (%)	24.10	24.30
Textural Class	62.30	62.20
	Clay	Clay
<b>Chemical Characteristics</b>		
pH (H <sub>2</sub> O)		
Organic matter (%)		
Total N (%)	4.23	4.14
Available P (mg/kg)	3.32	3.35
Exchangeable bases	0.06	0.14
Ca (cmol/kg)	6.16	6.18
Mg		
K	2.50	2.55
Na	1.30	1.28
Exchange Acidity (cmol/kg)	0.09	0.09
ECEC (cmol/kg)	0.06	0.07
	3.32	3.35
	7.27	7.34

**Table 2: Effects of N rates (kg/ha) on the leaf area (cm<sup>2</sup>) of five swamp rice varieties**

N-rate	1998	1999	Mean	Variety*	1998	1999	Mean
0	38.61	42.21	40.41	C	38.83	47.40	43.12
50	43.59	47.43	45.51	S	51.23	48.94	50.09
100	48.73	51.70	50.22	R	50.31	53.61	51.96
150	54.60	55.47	55.04	T	51.13	52.88	52.01
200	56.45	56.49	56.47	M	48.48	50.47	49.48
LSD(0.05)	3.29	4.06	3.68	LSD(0.05)	2.78	4.61	3.70

\*C = Cisadane, S = Suakoko 8, R = IR 5, T = ITA 306, M = MAS 2401

**Table 3: Effects of N rates (kg/ha) on the number of tillers per plant of five swamp rice varieties.**

N - rate	1998	1999	Mean	Variety	1998	1999	Mean
0	5.05	5.43	5.24	C	5.52	5.68	5.60
50	5.72	6.25	5.99	S	8.28	8.65	8.47
100	6.79	7.30	7.05	R	6.78	6.91	
150	7.86	8.29	8.08	6.85			
200	8.14	8.58	8.36	T	6.82	7.15	
LSD(0.05)	0.57	0.61	0.59	6.99			
				M	6.15	7.47	6.81
				LSD(0.05)	0.64	0.67	0.66

## RESULTS

### Soil

The physical and chemical characteristics of the soil used for the study are shown in Table 1. The total N contents were 0.06% and 0.14%, respectively, for 1998 and 1999 and were considered too low for the optimum performance of rice.

Rice growth were generally higher in 1999, probably due to the residual or cumulative effect of applied N fertilizer in the previous year.

### Leaf area

Leaf area was found to increase with increasing rates of N application up to the maximum rate in both years (Table 2). In 1998, the increase was optimum at 150kgN/ha with a mean leaf area of 54.60cm<sup>2</sup>. A lower rate of 100kgN/ha was however, found to be optimum in 1999 with an average leaf area of 51.70cm<sup>2</sup>. When averaged over the two years, 150 kgN/ha was optimum with a mean leaf area of 55.04cm<sup>2</sup>. This translated to a percentage leaf area increase of 36.20% over the control.

Except Cisadane which produced the smallest leaf area, all other varieties did not differ significantly with respect to this parameter (Table 2).

### Number of tillers per plant.

Nitrogen application significantly enhanced the tillering ability of the varieties. The highest significant numbers of tillers per plant were 7.86 and 8.29 respectively, for 1998 and 1999. Averaged over the two years, the optimum number of tillers per plant (8.08) was obtained with 150kgN/ha as against 5.24 tillers per plant for the control plot. This was

equivalent to a percentage increase of 54.20% in number of tillers per plant over the zero N treatment (Table 3).

Among the varieties, Suakoko 8 produced the highest significant mean number of tillers per plant and Cisadane, the least. ITA 306, IR 5 and MAS 2401 were not significantly different from one another (Table 3).

#### Number of spikes per panicle:

Number of spikes per panicle was significantly affected by N application. In 1998, spike number per panicle increased by 28.39% by increasing N level from 0 to 150kgN/ha. Similarly, in 1999, it increased significantly by 19.67% with the application of 100kgN/ha compared to the control (Table 4). Averaged over the two years the number of spikes per panicle appreciated significantly by 28.15% when the N level was increased from 0 to 150kg/ha.

The pair of MAS 2401 and Suakoko 8 produced the highest mean number of spikes per panicle, followed by the pair of IR 5 and ITA 306; while Cisadane recorded the least number (Table 4).

#### Grain Yield

There were significant responses of the varieties to applied N in both years, despite the critically low soil N content in the first year (Table 5). Averaged over the two years, the response was optimum at 150 kgN/ha with a grain yield of 3.80t/ha, as against the yield of 2.33 t/ha obtained from the control. This translated to a percentage grain yield increase of 63.09%.

Suakoko 8 produced a significantly highest mean grain yield of 4.60 t/ha, followed by the pair of MAS 2401 and ITA 306 (not significantly different). IR 5 and Cisadane produced the lowest grain yields (Table 5).

#### DISCUSSION

The total N contents of 0.06% (1998) and 0.14% (1999) obtained from the experimental plot were considered low. However, they were in agreement with Moormann (1980) who reported that soils of Southern Nigeria have low N content of between 0.02 and 1.16%. The low N content underscored the need for N fertilization to enhance good growth and yield.

The increase in the total N content of the soil in 1999 and the generally better performance of the five rice varieties in the same year, could be attributed to the residual or cumulative effect of applied N in the previous year. The result is similar to that of Adepoju (1991), who reported of higher yields of maize in the late season than in the early season due to the cumulative effects of applied P in the early season.

The grain yield and the growth parameters of the rice varieties were significantly increased by N application. This observation was similar to those reported by Agboola and Oko (1976) and Opikiri *et al* (1991); that N application at various rates significantly increased the grain yield and growth parameters of rice varieties. The rates of 100 and 150kgN/ha produced the highest effects. The rate of 100kgN/ha gave the broadest leaf area and the highest number of spikes per panicle in 1998 only. Similarly, the rate of 150kgN/ha produced the broadest leaf area and the highest number of spikes per panicle in 1999 and the highest number of tillers per plant and the heaviest grain yield in both years. It is obvious that the rate of 150kgN/ha was superior to other rates in influencing growth and yield.

Among the varieties, Suakoko 8 gave the highest significant response to the applied N. Notably, it produced the highest significant grain yields in both 1998 and 1999.

Besides, the variety had other desirable qualities, it was stiff-strawed, did not lodge and produced long, plump grains.

**Table 4: Effects of N rates (kg/ha) on the number of spikes per panicle of five swamp rice varieties.**

N-rate	1998	1999	Mean	Variety	1998	1999	Mean
0	8.70	9.20	8.95	C	7.33	8.23	7.78
50	9.25	10.41	9.83	S	11.37	12.34	11.86
100	10.09	11.01	10.60	R	9.87	11.28	10.58
150	11.17	11.76	11.47	T	10.77	10.25	10.51
200	11.71	12.10	11.91	M	11.69	12.38	12.04
LSD (0.05)	0.54	0.78	0.66	LSD(0.05)	0.55	0.61	0.58

**Table 5: Effects of N rates (kg/ha) on the grain yield (t/ha) of five swamp rice varieties**

N-rate	1998	1999	Mean	Variety	1998	1999	Mean
0	2.35	2.31	2.33	C	2.10	1.63	1.87
50	2.70	2.68	2.69	S	4.53	4.66	4.60
100	3.17	3.25	3.21	R	2.88	3.01	2.95
150	3.81	3.78	3.80	T	3.15	3.23	3.19
200	3.82	3.83	3.83	M	3.21	3.33	3.27
LSD(0.05)	0.14	0.35	0.25	LSD(0.05)	0.17	0.33	0.25

Curiously, the yield of ITA 306 was relatively poor. This is inconsistent with the report by the National Cereals Research Institute (NCRI), as cited by IITA (1986), that the variety possesses superior yielding ability over varieties currently being cultivated in Nigeria. Its mean grain yield of 3.19t/ha is a considerable departure from the yield of 6.5-7.5t/ha reported by the source cited above. Be as it may, the variety had some desirable qualities, it was short statured, high tillering and short maturing. Other varieties namely, IR5 and MAS 2401 responded fairly well to the applied N but the later variety was long maturing and exceptionally tall, hence highly susceptible to lodging. In all, Cisadane variety responded poorly and was easily the least adaptable variety to the wetland ecology of the study area.

#### CONCLUSION

Rice growth and yield were found to have increased as the level of N increased, up to the maximum level. However, the highest significant increases were obtained with the rate of 150kgN/ha. This suggests that 150kgN/ha was the most suitable N rate for the tropical wetland soil of the study area for rice production.

Among the varieties, Suakoko 8 performed relatively better, especially in grain yield, than the other varieties. It is obvious that this variety was the most responsive to the applied N fertilizer.

#### REFERENCES

- Adepoju, C., 1991. The response of maize to P fertilization in an entisol. *Nig. Agric. J.* 26: 11-19.
- Agboola, A.A. and Oko, B.F.D., 1976. Effect of different sources of N on the yield of upland rice in tropical rainforest of Western Nigeria. *Nig. Agric. J.* 13 (3): 98-113.
- Bates, R.G., 1954. *Electrometric pH determinations*. John Wiley and Sons Inc., New York.
- Bouyoucos, G. J., 1951. Improved hydrometer method for making particle size analysis of soil. *Agron. J.* 54: 464-465.

- Bray, R. H. and Kurtz, L.T. , 1954. Determination of total organic and available forms of phosphorous in soils. *Soil Sci.* 59: 39 45.
- De Datta, S. K., 1987. Nitrogen transformation processes in relation to improved cultural practices for lowland rice. *Plant and soil* 100: 47 69.
- De Datta S.K., Fillery, I. R.P., Obcemea, W.N. and Evangelista, R. C., 1987. Floodwater properties, nitrogen utilization and nitrogen balance in a calcareous lowland rice soil. *Soil Sci. Soc. Am. J.* 51 (5): 1356.
- Foth, H.D., 1978. *Fundamentals of soil science* (6<sup>th</sup> ed.). John Wiley and sons, New York, 311 p.
- Gomez, K.A. and Gomez, A.A., 1984. *Statistical procedures for Agricultural Research* (2<sup>nd</sup> ed.) John Wiley, New York. Pp.199 202.
- Grist, D.H., 1986. *Rice* (6<sup>th</sup> ed.). Longman Group Limited., New York. Pp. 271 276.
- Hammer, G.L., 1980. Estimation of cassava leaf area by a simple, non-destructive field technique. *J. Aust. Inst. of Agric. Science* 46(1): 61 62.
- IITA, 1979. International Institute of Tropical Agriculture (IITA). *Selected Methods for Soil and Plant Analysis Manual Series No. 1.* IITA, Ibadan.
- IITA, 1986. International Institute of Tropical Agriculture (IITA). *IITA Annual Report and Research Highlights* IITA, Ibadan 50p.
- Jackson, M.L., 1962. *Soil Chemical Analysis.* Prentice Hall Inc., New Jersey.
- Moormann, F. R., 1980. Representative Toposequence in Soil in Southern Nigeria and their Pedology. *Characterization of Soils.* oxford Press, New York . 33p.
- NAKSRDP, 1994. Northern Akwa Ibom Swamp Resources Development Program Northern Akwa Ibom Swamp Resources Development Study. 2 Main Report.
- Okoji, M. A., 1995. Wetland Agriculture in Ini: Its Development and Sustainability. Paper presented at the third all African soil science Conference, University of Ibadan, Nigeria. Pp 1 16.
- Opukiri, S. B., Zuofa, K. and Douglass, D.C., 1991. The influence of urea fertilization on yield of rice on acid soil in Rivers State. *Nig. J. of Crop/Soil Science and Forestry* (1): 42 45.
- Oyedokun, J. B. and Sobulo, N., 1979. Responses of improved rice varieties to spacing and nitrogen in South-Western Nigeria. *Ghana J. Agric. Sci.* 10:121-127.
- Purseglove, J.W., 1975. *Tropical Crops Monocotyledons* (Vol. 1 and 2 combined). Longman Group, London. Pp. 161 197.
- Sobulo, R., 1982. Response of rice to NKP. Fe and Zn in Western Nigeria. *Nig. J. of Soil Science* 3:37- 42.
- Walkley, A. and Black, F. A., 1934. An examination of the different methods of determining soil organic matter and a proposed modification of the chromic acid titration method. *Soil Sci.* 37: 29 38.