

# RESPONSE OF MICROBIAL ENZYMES SYNTHESSES TO TOXICITY OF WEATHERED AND BIODEGRADED OILS.

L. O. ODOKUMA and G. C. OKPOKWASILI

(Received 17 September 2002; Revision accepted 16 December 2002)

## ABSTRACT

The toxicity of eight weathered and biodegraded Nigerian crude oils to the biosyntheses of four bacterial enzymes were examined. The constitutive intracellular enzyme, nitritase was the most sensitive to all toxicants. Two extracellular inducible enzymes tryptophanase and  $\alpha$ -glucosidase were less sensitive to the toxicants. These extracellular enzymes, however, were more sensitive than the intracellular inducible enzyme  $\beta$ -galactosidase. Inhibition of bacterial enzymes syntheses could be a useful, cheap, simple and rapid ecotoxicity assay for monitoring oil pollution in the Nigerian environment.

**Key words;** Sensitivity, Response, Microbial Enzymes, Toxicity, Weathered crude oil, Biodegraded crude oil

## INTRODUCTION

The environmental transformations of an organic chemical in many natural aquatic systems may be induced by physical, chemical and microbial factors within the system (Plimmer, 1978; Smith and Dragun, 1984; Okpokwasili and Odokuma, 1994; Okpokwasili and Nnubia, 1994). Chemically and biological, induced changes in the composition of a crude oil are known collectively as weathering (Atlas, 1981). Microbial degradation contributes significantly to the ultimate removal of oil from the environment (Atlas, 1981; Amanchukwu *et al.*, 1989). Some of the components of crude oils may be toxic to microorganisms. Short-chain- paraffins (>C10) generally are assumed to be toxic to microorganisms because of their water solubility and dual interactions with lipid membranes (Teh, 1974; Teh and Lee, 1974). Griffin and Calder (1977) showed that weathered oils produced water-soluble fractions (WSF) that were more toxic to *Serratia marino rubra* than the parent oils.

Many toxicity tests are accomplished by using microorganisms as test organisms (Williamson and Johnson, 1981; Wang, 1984; Giesy *et al.*, 1988; Jardim *et al.*, 1990; Xu and Schurri, 1990). Such bioassays must provide rapid and reliable data so they can be extrapolated to infer human risk of exposure as well as to trigger adequate waste management. These tests are based on the inhibition of some microbial vital function. Examples include inhibition of nitrite utilization by *Nitrobacter* (Williamson and Johnson, 1981; Wang, 1984). Microtox tests (Giesy *et al.*, 1988), Ames

Salmonella assay (Vandermeulen, 1986; Vandermeulen and Lee, 1986), SOS chromotest (Quillardet *et al.*, 1982) and carbon dioxide production (Jardim *et al.*, 1990). Others include the monitoring of internal pools of DNA and ATP (Jardim *et al.*, 1990) These techniques have rapidity and simplicity in achieving results as notable advantages.

Because enzymes are key catalysts of metabolic reactions in cells the inhibition of their activity or biosyntheses has been recently explored as a basis for ecotoxicity testing (Dutton *et al.*, 1990). Beta galactosidase, an enzyme that confers *Escherichia coli* with the ability to break down lactose to galactose and glucose is constitutive while the  $\beta$  galactosidase in *E. coli* is intracellular and induced by the presence of lactose. Tryptophanase, an inducible enzyme produced by *E. coli* in the presence of L-tryptophan, is extracellular. Both  $\beta$ -galactosidase and tryptophanase production are controlled by different operons in *E. coli* (Magasanik, 1988). Alpha-glucosidase, another inducible extracellular enzyme, is produced by *Bacillus licheniformis* in response to maltose (Magasanik, 1988). Studies on the effects of some toxicants on the biosyntheses of the three enzymes  $\beta$ -galactosidase, tryptophanase and  $\alpha$ -glycosidase have been initiated (Dutton *et al.*, 1990). Weathering involves the physical, chemical and biological degradation of materials (Plimmer, 1978). It involves physical factors such as the effect of wind, rain, temperature and the abrasive actions of rocks. It includes chemical factors such as hydrolysis, salinity, solution effect etc. The biological aspects include the degradative

L. O. ODOKUMA, Department of Microbiology, University of Port-Harcourt, Port-Harcourt, Nigeria  
G. C. OKPOKWASILI, Department of Microbiology, University of Port-Harcourt, Port-Harcourt, Nigeria

**Editor's Note:** This articles was earlier published in Global J. Pure & Applied Science 9 (2): 183-191 (2003). It is republished here because of errors in tables.

activities of micro and macroorganisms on materials. Thus weathering may be physical and chemical i.e. without biological input. It may also be biological. In this study however, weathered oils are regarded as oils that have been exposed mainly to physical and chemical forces, while biodegraded oils are those that have been exposed to physical, chemical and biological (microbial) forces.

Microbiological assays of chemicals are insufficient in deciding whether a particular chemical is toxic since the ecosystem contains both macro-and microorganisms. If the objective of the bioassay test is to ensure protection of all aquatic species then the inclusion of microorganisms is necessary as they clearly represent a greater biomass in natural systems. Their use also offers rapid, low cost and simple alternatives for achieving bioassay results, (Williamson and Johnson, 1981).

The objective of this study was to compare the inhibitory effects of weathered and biodegraded Nigerian crude oils on the biosyntheses of  $\beta$ -galactosidase, tryptophanase,  $\alpha$ -glucosidase and nitritase. The goal was to identify the operon-enzyme system most sensitive to these toxicants which will thus serve as a rapid and economic method for assessing the toxicity of spilled oil in the Nigeria environment.

## MATERIALS AND METHODS

### Test Bacteria

Toxicity assays were conducted with *Escherichia coli*, *Bacillus licheniformis* and *Nitrobacter* sp. *E. coli* and *Nitrobacter* sp. were both isolated from the New Calabar River in Rivers State Nigeria while *B. licheniformis* was isolated from the soil near the Microbiology laboratory in the University of Port Harcourt. To ensure adaptation to laboratory conditions all strains were maintained by continuous subculture every 48h into fresh media.

### Characterization and identification of Test Bacteria

Isolates were examined for colony, morphology and biochemical characteristics. For *E. coli* and *B. licheniformis*, the tests included Gram reaction, spore staining, oxidase test, oxidative/fermentative. (O/F) utilization of glucose, motility test, indole test, Voges-Proskauer, catalase test, urea utilization, aerobic and anaerobic growth, growth in 7% NaCl and at 55° to 60°C, which were carried out according to the methods described by Cruickshank *et*

*al.*(1975). For *Nitrobacter* sp., Gram reaction, colony morphology and sole utilization of nitrite as energy source, were used for its identification according to the method of Colwell and Zambruski (1972). Identification of bacteria to species level followed the scheme in Laskins and Lechevalier (1977).

### Chemical Reagents and Toxicants

All reagents employed in the study were of analytical grade and were purchased from Sigma Chemical Company, St. Louis, Missouri. Parent Crude oils assessed for toxicity were Antan Heavy (AH), Antan Medium (AM), Bonny Medium (BM) and Bonny light (BL). Others were Ughelli Quality Control Centre crude (UQCC), Trans Niger Pipeline (TNP), Brass River (BR) and Qua Iboe (Q1). All crude oils were provided by the Nigerian National Petroleum Corporation (NNPC), Port Harcourt, Nigeria.

### Weathered (physical and Chemical) Crude Oils

Weathered crude oils in this study consisted of a mixture of crude oil and sterile deionized water. There were no microorganisms (seed) from external sources.

### Biodegraded (Physical, Chemical and Biodegraded) Crude Oils.

This consisted of a mixture of crude oil with sterile deionized water into which microorganism (seed) from the New Calabar River had been introduced.

### Preparation of Toxicant Concentrations

Toxicant concentrations of 1000, 100, 10, 1.0 and 0.1 mg/L were prepared. For liquid toxicants an amount equivalent to one thousand milligrams of liquid toxicant was weighed and transferred into distilled water (1000ml of distilled water minus the equivalent volume of 1000mg of liquid toxicant). A similar procedure was repeated for other concentrations (100, 10, 1.0, 0.1). For example an amount equivalent to 100mg of the liquid toxicant was weighed and transferred into 100ml of distilled water minus the equivalent volume of 100mg or the liquid toxicant to produce 100mg/l of the toxicant.

Preliminary range finding test was undertaken. Organisms were exposed to logarithmic concentrations of toxicants. The least toxicant concentration that produced total inhibition of the enzyme system (or death of organism) was taken as 100% inhibition (the highest toxicant concentration). Also the highest toxicant

**TABLE 1: Effect Of Fresh (Parent) Crude Oils On The Inhibition of biosynthesis of four enzymes**

Crude oil	IC <sub>50</sub> (mg/L)			
	$\beta$ -galactosidase	Tryptophanase	$\alpha$ -glucosidase	Nitritase
Qua Iboc	NT	150	85,000	15.0
Brass	NT	140	NT	40.0
Rivers				
Bonny light	NT	400	NT	10.0
Bonny medium	80,000	8.0	NT	60.0
TNP	NT	2.0	NT	100
UQCC	NT	100	NT	60
Antan Heavy	1.0	0.8	8,000	3.5
Antan Medium	1.0	0.8	8,000	70

NT (Not toxic)= above 1,000000 mg/L

concentration that could produce no inhibition of enzyme synthesis was taken as the lowest toxicant concentration. Toxicant concentrations were then produced between these two limits.

Concentrations varied according to the toxicity of the toxicant To 50ml of distilled water, 0.1g of solid toxicant (or its equivalent in volume for liquid toxicants) was added. The volume was made up to 100 ml to produce a concentration of 1000 mg/L. Serial tenfold dilutions of this stock were produced till required minimum concentration (concentration that will have no effect on the enzyme biosynthesis) was achieved.

#### Toxicity Assay Protocol

The method of Dutton *et. al.* (1990) was employed for assays for the biosyntheses of  $\beta$ -galactosidase, tryptophanase and  $\alpha$ -glucosidase. They had the following basic steps in common: (1) cell growth (2) cell washing (3) cell exposure to toxicant; (4) induction of enzyme biosynthesis and (5) measurement of enzyme activity. Nitritase biosynthesis assay also followed the same steps except that the measurement of enzymatic activity was done indirectly by measuring the loss of substrate (nitrite) as in APHA (1985).

#### Cell Growth and preparation

*Escherichia coli* was grown in nutrient broth overnight at 37°C. Cells were diluted with medium to A<sub>550</sub> = 0.2. (Absorbance 0.2 at 550 nm). Medium without organisms was used as blank. Cells were allowed to grow to A<sub>550</sub> = 0.6 after which were washed thrice with 0.8% NaCl.

#### Exposure to Toxicant

Cell (0.1 ml) of washed cell suspension at A<sub>550</sub> = 0.6) were exposed to 0.9ml of toxicant and incubated for 30min.

#### Control

Control contained cell plus medium with no toxicant.

#### Enzyme Induction.

To 1ml each of the reaction mixture, 0.1 ml Isopropyl- $\beta$ -D-thiogalactoside (IPTG) 0.1%(w/v) (Sigma Chemical Company,) 0.5 – buffer and 0.5 ml growth medium were added. The mixture was incubated for 30 min.

#### Beta – galactosidase Measurement.

Since  $\beta$ -galactosidase is intracellular, it was necessary to lyse the cells. This was performed by the addition of 0.1 ml 10% Sodium

Dodecyl Sulphate (SDS) 10% (w/v) to the reaction mixture. This was followed by the addition of 0.2 ml *O*-nitrophenyl-*D*-galactoside (ONPG) 0.4% (w/v) Sigma Chemical Company. The mixture was incubated until colour developed (15 min). The reaction was stopped with 1 ml of cold Na<sub>2</sub>CO<sub>3</sub> (1M) and the absorbance at 542nm using a blank consisting of all reagents minus IPTG was determined (Dutton *et al*, 1990).

#### Tryptophanase Biosynthesis Assay

- (i) L-tryptophan (250 mg/L) in KH<sub>2</sub>PO<sub>3</sub> (13.6 g/L) adjusted to pH -7.8
- (ii) Ehrlich's reagent consisting of 6 parts of *p*-dimethylaminobenzaldehyde (5%, w/v) Sigma Chemical Company, in 95% ethanol and 12 parts acid alcohol (16 ml conc. H<sub>2</sub>SO<sub>4</sub> in 200 ml 95% ethanol).

#### Cell growth and preparation

*Escherichia coli* was grown in casein hydrolysate without tryptophan (10g/l), yeast extract (5 g/L) and NaCl (10g/l) overnight at 37°C.

Cell preparation was as for β-galactosidase biosynthesis assay

#### Exposure to Toxicant

Cells (0.1 ml of washed cell suspension at A<sub>550</sub> = 0.6) were exposed to 0.9 ml toxicant and incubated for 30. min.

#### Enzyme induction

To the reaction mixture 0.4 ml buffer, 0.1 L-tryptophan and 0.5 ml nutrient broth were added. The mixture was incubated for 120 min.

#### Tryptophanase Measurement

One milliliter of Ehrlich's reagent was added to the reaction mixture and incubated for 15 min. Absorbance at 568nm using a blank consisting of all reagents minus L-tryptophan, was determined.

#### Alpha-glucosidase Biosynthesis Assay

Maltose (4% w/v) and autoclaved *p*-nitrophenyl β-D-glucoside (PNG) 0.5% (w/v) were used. This solution was filter-sterilized (0.22µm pore size) and stored at 4°C in the dark.

#### Cell Growth, Preparation and Enzyme Induction

*Bacillus licheniformis* – was grown in tryptone soya broth without dextrose overnight at 37°C. Cell preparation and

toxicant exposure were as in β-galactosidase biosynthesis assay.

To the reaction mixture 0.4 ml Z-buffer. (Na<sub>2</sub>HPO<sub>4</sub> 7H<sub>2</sub>O 16.1g/L NaH<sub>2</sub>PO<sub>4</sub> H<sub>2</sub>O 5.5. g/L KCL 0.75 g/L; and MgSO<sub>4</sub> 7H<sub>2</sub>O 0.25 g/L) 0.1 ml maltose and 0.5 fresh medium were added. The mixture was incubated for 60 min.

#### Alpha-glucosidase Measurement

*O*-nitrophenyl-*D*-galactoside (ONPG) (0.2 ml) was added and the mixture incubated for 60 min. Reaction was stopped by the addition of 1ml Na<sub>2</sub>CO<sub>3</sub>. The absorbance of the mixture at 420nm was measured. A solution consisting of all other reagents minus maltose was used as blank for the reading.

#### Nitritase Biosynthesis Assay

The reagents used were as in method for nitrite determination APHA (1985). The other steps growth, cell preparation, exposure to toxicant, enzyme induction and nitritase measurement were as reported earlier (Okpokwasili and Odokuma, 1994).

#### Toxicity And Analysis

The method employed was adopted from Dutton *et al* (1990). One hundred milliliters of serial dilutions of each fresh crude oils were placed in 250 ml cotton wool-plugged shake flasks. The flasks were incubated for 4 weeks. Enzyme biosynthesis assays for nitritase, tryptophanase β-galactosidase and α-glucosidase were carried out on each of the crude oil concentrations. Periodicity of assays was immediately after preparation of toxicant concentration (day 0) and thereafter once a week for four weeks. This served as base-line control data.

Water samples containing a mixed population of hydrocarbonoclastic microorganisms was obtained from the New Calabar River at a site located about 1 km southwest of the University of Port Harcourt, Nigeria

#### Seeding of crude oil samples

To 75 ml of each crude oil concentration, 25 ml of the New Calabar River water was added

All concentrations for the enzyme assay were prepared in triplicate. Preliminary range finding test was carried out where possible (because some toxicants did not inhibit the enzyme systems). To determine toxicant concentration causing between 10 and 100% inhibition, the degree of enzyme inhibition was determined by measuring absorbance with respect to control

**TABLE 2: Effect of weathering and biodegradation of eight crude oils on the biosynthesis of nitritase**

Crude oil	IC <sub>50</sub> (mg/l)									
	Weathered crude oil (weeks)					Biodegraded crude oil (weeks)				
	0	1	2	3	4	0	1	2	3	4
Bonny Medium	55	0.2	0.2	4.0	4.0	100.0	1.00	0.35	0.03	0.02
Bonny Light	40	50.0	6.0	0.075	0.06	40	0.30	0.09	0.07	0.10
Brass River	120	0.2	0.5	0.50	30.0	40	1.0	0.50	0.095	0.0075
Qua Iboe	105	0.9	0.11	0.15	0.08	118	9.0	0.70	0.11	0.01
UQCC	50	0.10	0.40	0.09	0.08	60	5.5	0.10	0.10	0.01
TNP	200	0.09	0.10	0.50	0.09	300	0.85	0.40	0.02	0.03
Antan Medium	70	0.4	5.0	0.20	10	70	2.0	4.0	0.01	0.09
Antan heavy	0.2	0.07	0.10	0.05	0.008	6.0	0.08	0.095	0.10	0.005

NT (Not toxic) = above 1,000000 mg/l

(assigned 0% inhibition). Control contained no toxicant. For nitritase biosynthesis assay, the ratio of the absorbance in control to that of toxicant multiplied by 100 was taken as percent toxicant inhibition. For the three other enzyme systems, the ratio of the absorbance of the toxicant to that of the control multiplied by 100 was taken as the percent toxicant inhibition. The mean of triplicate samples was obtained and used to plot a graph of percent inhibition versus concentration of toxicant. The chemical concentration giving 50% inhibition (IC<sub>50</sub>) for weathered and biodegraded crudes was derived from the line of regression of the plot.

## RESULTS

The results of the effect of eight Nigerian parent crude oils on the inhibition of biosynthesis of four enzymes are presented in Table 1. The data revealed that nitritase was the most sensitive to the crude oils tested. This was followed by tryptophanase

and  $\alpha$ -glucosidase. Antan Heavy (AH) was the most toxic (IC<sub>50</sub> values were the least except in  $\alpha$ -glucosidase).

The effects of weathering and biodegradation of eight crude oils on the biosyntheses of four enzymes (nitritase, tryptophanase,  $\alpha$ -glucosidase and  $\beta$ -galactosidase) are presented in Tables 2-5. The data indicated that nitritase was the most sensitive followed by tryptophanase. (the IC<sub>50</sub> values of nitritase and tryptophanase were the least compared to the IC<sub>50</sub> of the other enzymes). Beta-galactosidase showed the least sensitivity (see Table 4). There was an increase in the toxicity of all crude from day 0 to the seventh day. By the 4th week, biodegraded crude become slightly more toxic to nitritase than weathered crude. An increase in the sensitivity of nitritase to both the biodegraded and weathered oils throughout the incubation period was observed. Results in Tables 3, 4 and 5 showed that weathered, crude oils were

**TABLE 3: Effect of weathering and biodegradation of eight crude oils on the biosynthesis of Tryptophanase**

Crude oil	IC <sub>50</sub> (mg/l)									
	Weathered crude oil (weeks)					Biodegraded crude oil (weeks)				
	0	1	2	3	4	0	1	2	3	4
Bonny Medium	10.0	6.0	4.5	7,500	NT	10.0	0.6	700.0	90.0	NT
Bonny Light	500	6.2	30.0	80.0	NT	400	1.100	NT	100	NT
Brass River	80	2.5	8.5	0.09	NT	70.0	14.0	NT	10.000	NT
Qua Iboe	130	80.0	1.0	400	NT	130	9.5	NT	5.0	NT
UQCC	100	4.0	600	250	NT	100	1.000	30.0	105	NT
TNP	2.0	6.5	20000	80.0	NT	2.0	7.000	200	9.0	NT
Antan Medium	3.0	5.0	NT	2.000	NT	0.5	70.0	3.0	7.0	1,000
Antan heavy	0.8	60.0	NT	60.0	10.000	1.1	10.0	5.0	10.0	5,000

NT (Not toxic)=above 1,000000 mg/l

slightly more toxic to  $\beta$ -galactosidase and tryptophanase than their biodegraded counterparts. This was obvious in the 4<sup>th</sup> week. However, for both enzymes there was a decrease in toxicity from day 0 to the 4<sup>th</sup> week of both weathered and biodegraded oils.

## DISCUSSION

Nitritase biosynthesis was the most sensitive to all toxicants. This was followed by tryptophanase and  $\alpha$ -glucosidase biosyntheses. Beta  $\beta$ -galactosidase biosyntheses showed the least sensitivity to these toxicants. There was no significant difference between the sensitivity of the enzymes to weathered crude oils and to biodegraded crude oils. Weathering is the physical, chemical and biological degradation of substances in *situ* (Plimmer, 1978). Therefore weathering includes the effects of radiation, wind, rocks, solution, hydrolysis, microorganisms and macroorganisms on

materials. In this study-weathered crude oils meant oils that were exposed to both physical and chemical factors. While biodegraded crude oils referred to crude oils that were exposed to physical chemical and biological (microbial degradation) factors. A number of factors such as solubility, substitution, chain length and concentration are responsible for the toxicity of an organic pollutant (Babich and Stotzky, 1977; Vandermeulen and Lee, 1986; Petibone and Cooney, 1988). Griffin and Calder (1977) have revealed that the addition of crude oil to water produces water soluble fractions (WSF). The production of WSF concentrations of between 17.5-26.0ug/L by three Nigerian crudes, Bonny medium Bonny light and Forcados blend has been reported (Imevbore *et al.*, 1987). The production of WSF by the crude in this study may be responsible for their toxicities. The relative insensitivity of  $\beta$ -galactosidase biosynthesis to most crude oils may be due to the inability of the WSF of these crudes to penetrate the outer membrane of *E. coli*

either due to low concentration of WSF or lack of permeases to transport the toxicant across the cell membrane. Therefore this enzyme would make a poor toxicity index. Similar results have been obtained by Dutton *et al.* (1990). Their studies revealed the relative insensitivity of  $\beta$ -galactosidase by *E. coli* to hydrophiles such as hydrothol. The higher sensitivity of nitritase biosynthesis may be due to the high permeability of the outer membrane of *Nitrobacter* sp to the WSF of the crude oils. The influx of the WSF probably inactivated the nitritase enzyme present within the cell. Nitritase would thus provide first choice as a toxicity index. The higher sensitivity of tryptophanase synthesis to crude oils than  $\beta$ -galactosidase biosynthesis in the same *E. coli* strain may be as a result of the extra-cellular nature of tryptophanase enzyme. This may result in the enzyme making easier contact with the WSF of the crude oil. A similar result was obtained by Dutton *et al.* (1990) who observed a higher sensitivity of

tryptophanase biosynthesis to Sodium dodecyl sulphate (SDS) and polychlorinated phenyls (PCP) than  $\beta$ -galactosidase biosynthesis. Gram positive bacteria, such as *Bacillus licheniformis*, do not possess an outer membrane (Koch and Schaechter, 1985) and thus may be expected to be more sensitive to the WSF crude oils than Gram negative bacterial such as *E. coli* and *Nitrobacter* sp. The data, however, showed that *B. licheniformis* was relatively insensitive to most crude oils. The production of spores by *B. licheniformis* may have been responsible for its insensitivity.

Antan medium and Antan heavy were the most toxic of the test crude oils (especially to the enzyme systems  $\beta$ -galactosidase tryptophanase and  $\alpha$ -glucosidase, see Table 1) The IC<sub>50</sub> for Antan heavy for  $\beta$ -galactosidase, tryptophanase,  $\alpha$ -glucosidase and Nitritase were 1.0, 0.8, 8000 and 3.5mg/L respectively. For Antan light the IC<sub>50</sub> for these enzyme systems were 1.0, 0.8, 8000 and 0.2mg/L respectively. This may have

**TABLE 4: Effect of weathering and biodegradation of eight crude oils on the biosynthesis of  $\beta$  galactosidase**

Crude oil,	IC <sub>50</sub> (mg/l)									
	Weathered crude oil (weeks)					Biodegraded crude oil (weeks)				
	0	1	2	3	4	0	1	2	3	4
Bonny Medium	NT	NT	NT	NT	NT	NT	NT	NT	10000	NT
Bonny Light	NT	1,000000	NT	80.0	NT	NT	NT	NT	NT	NT
Brass River	NT	7000	NT	NT	NT	NT	NT	NT	NT	NT
Qua Iboc	NT	10000	NT	NT	NT	NT	NT	NT	NT	NT
UQCC	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT
TNP	NT	NT	NT	NT	NT	NT	NT	NT	NT	NT
Antan Medium	4.5	20.0	NT	NT	NT	4.5	NT	NT	NT	NT
Antan heavy	1.1	2000	NT	NT	1,000000	4.0	NT	NT	NT	NT

NT (Not toxic)-above 1,000000 mg/l

**TABLE 5: Effect of weathering and biodegradation of eight crude Oils on the biosynthesis of  $\alpha$ -glucosidase**

Crude oil	IC <sub>50</sub> (mg/l)									
	Weathered crude oil (weeks)					Biodegraded crude oil (weeks)				
	90.0	1	2	3	4	0	1	2	3	4
Bonny Medium	30000	5,000	NT	NT	900000	07,0	NT	700.0	90.0	NT
Bonny Light	30000	0.6	NT	0.09	1,000	NT	NT	NT	0.85	NT
Brass River	40000	0.6	NT	NT	NT	NT	NT	NT	10,000	NT
Qua Iboe	NT	0.8	700000	NT	600000	NT	NT	NT	50000	NT
UQCC	NT	5,000	70000	NT	30.0	NT	NT	NT	NT	1,000
TNP	20,0	80,0	NT	NT	NT	NT	NT	10,000	NT	NT
Antan Medium	5000	0.07	NT	NT	NT	8000	700	10000	NT	NT
Antan heavy	0.025	0.50	NT	NT	7.7.0	60.0	3,000	600000	100	60.0

NT (Not toxic) = above 1000000 mg/l

been due to the production of a higher concentration of WSF rather than the presence of more toxic components in the WSFs. Bonny medium was more toxic to  $\beta$ -galactosidase and tryptophanase enzyme system than Bonny light. The IC<sub>50</sub> of Bonny medium to  $\beta$ -galactosidase and tryptophanase were 80,000 and 8.0mg/l respectively while the IC<sub>50</sub> of Bonny light to these enzymes systems were > 1000000 and 400 mg/l respectively. Bonny medium was more toxic than Bonny light. The observations differ from the reports of Imevboire *et al.* (1987). They observed that light crudes were more toxic to two shrimps (*Desmocariss trispinosa* and *Palaemonetes africanus*) than medium crudes. The results from the study may be due to the production of more toxic components in the WSF of heavier crude oils than a WSF concentration difference.

The effect of weathering and biodegradation on the biosyntheses of nitritase, tryptophanase and  $\beta$ -galactosidase is shown in Tables 2-5. The data showed that nitritase was the most sensitive, followed by tryptophanase. Beta-galactosidase showed the least sensitivity to both biodegraded and weathered oils. Explanation

adduced for the toxicity of the parent oils may also be responsible for these results. The decrease in toxicity of both biodegraded and weathered oils to these enzymes from day 0 to the 7<sup>th</sup> day may be attributed to the increase in the concentration of WSF. Griffin and Calder (1977) observed that weathered oils produced WSFs that were more toxic to *Serratia marinorubra* than the parent oils. They attributed this increased toxicity to the higher concentration of hydrocarbons in the WSF. By the 4<sup>th</sup> week, biodegraded oils became slightly more toxic to nitritase biosynthesis than weathered crudes. This may be attributed to the production of degradation products or metabolites by the hydrocarbonoclastic organisms that are more toxic than the parent oils to *Nitrobacter*. The toxicity of these products to *Nitrobacter* increased with incubation time which may have been responsible for the increase in sensitivity of nitritase with time for both biodegraded and weathered crude oils. Table 3, 4, and 5 revealed that weathered crudes were slightly more toxic to  $\beta$ -galactosidase and tryptophanase biosyntheses than biodegraded crude, especially by the end of the first and fourth weeks. This may be attributable to the high tolerance of *E. coli* to



the intermediate produced by the microorganism in the biodegraded samples. The high tolerance of *E. coli* to these intermediates may account for the decrease in sensitivity of biosyntheses of two enzymes from the second week to the fourth week for both biodegraded and weathered oils.

## CONCLUSION

Nitritase biosynthesis was the most sensitive to the toxicants, followed by tryptophanase and  $\alpha$ -glucosidase biosyntheses with similar sensitivities. Beta-galactosidase biosynthesis was the least sensitive to the toxicants. The medium inhibition concentration ( $IC_{50}$ ) for nitritase for weathered Bonny medium Bonny light, Brass River, Qua Iboe, UQCC, TNP, Antan medium and Antan heavy after 4 weeks were 4.0, 0.06, 30.0, 0.08, 0.08, 0.09, 10 and 0.008mg/L respectively. The  $IC_{50}$  for the same crude oils for tryptophanase and  $\beta$ -galactosidase for the same period were all greater than 1,000,000mg/l except Antan heavy for tryptophanase. The  $IC_{50}$  for  $\alpha$ -glucosidase for all crude oils for the same period were 900,000, 1000, >1000,000, 600,000, 30, >1000,000, >1000,00 and >7.0 mg/L, respectively.

The study shows that Nitritase enzyme inhibition will serve as a more sensitive indicator for the determination of toxicity levels of crude oils.

## REFERENCES

- Amanchukwu, S. C., Obafemi, A. and Okpokwasili, G.C., 1989. Petroleum hydrocarbon utilization and degradation by a palm wine yeast isolate. *Fems microbiology letters* 57: 151-154.
- American Public Health Association, 1985. Standard Methods For The Examination Of Water And Wastewater 16<sup>th</sup> edition.
- Atlas, R. M. 1981. Microbial degradation of petroleum hydrocarbons: An environmental perspective. *Microbiological Reviews* 45: 180-209.
- Babich, H and Stotzky, G. 1977. Sensitivity of various bacteria, including actinomycetes and fungi to cadmium and the influence of pH on sensitivity. *Applied and Environmental Microbiology* 33 (3): 681-695.
- Colwell, R. R and Zambruski, M. U. 1972. Methods in Aquatic Microbiology. Pp 1-453. University Park Press., Baltimore
- Cruickshank, R. Duguid J.P Marmion, B. P and Swain, R.H A. 1975 Medical Microbiology 12<sup>th</sup> edition. Pp 1-947. Churchill Livingstone, London
- Dutton, R. J., Bitton, G., Koopman, E. and Agami O. 1990 Effect of environmental toxicants on enzymes biosynthesis: A comparison of  $\beta$ -galactosidase,  $\alpha$ -glucosidase and tryptophanase. *Archives to Environmental Contamination and Toxicology* 1: 398
- Giesy, J. P., Graney, R. L. Newsted, J. P. Rosiu, C. J., Benda, A., Kreis R. G. and Horvath, F. J. 1988. Comparison of three sediment bioassay methods using Detroit River sediments. *Environment Toxicology Chemical*, 7: 483-498.
- Griffin, L. F. and Calder, J. A. 1977. Toxic effect of water-solution fractions of crude refined and weathered oils on the growth of a marine bacterium. *Applied and Environmental Microbiology* 33. 1092-1096.
- Imevbore, A. M. A., Adeyemi, S. M. and Afolabi, O. A., 1987. The toxicity of Nigeria crude oils to aquatic organisms. In Proceedings of the 1987 seminar on The Petroleum industry and the Nigerian Environment. Pp 171-174. Lagos :Nigeria National Petroleum Corporation.
- Jardim, W. F., Pasquini, C., Guimaraes, J. R. and De Fraia, L.C., 1990. Short-term toxicity test using *Escherichia coli*; monitoring CO<sub>2</sub> production by flow injection analysis. *Water Research* 24: 351-354
- Koch, A.L. and Schaechter, M. S., 1985. The world ways of *Escherichia. coli*: In: Demain, A.L. and Solomon N.A. (ed) *Biology of Industrial Microorganizations*, pp.1-25 Menlo Park: Benjamin Cummings.
- Laskin A. I. and Lechevalier H. A., (ed) 1977. pp 1-1054 West Palm Beach: CRC Press, Inc.
- Magasanik, B. 1988. Research on bacteria in the mainstream of biology. *Science* 243:5-1439.
- Okpokwasili, G. C. and Odokuma, L. O. 1994. Tolerance of *Nitrobacter* to toxicity of some Nigerian crude oils. *Bulletin of Environmental Contamination and Toxicology*, 52. 388-395
- Okpokwasili, G. C. and Nnubia, C. 1994. Effect of drilling fluids on marine bacteria from a Nigerian offshore oilfield. *Environmental Management*.
- Petibone, G. W. and Cooney, J. J., 1988. Toxicity of methyl tine to microbial populations in estuarine sediments. *Journal of Industrial Microbiology*, 2: 373-378.

- Plimmer, R. J., 1978. Degradation methodology: chemical and physical effects. The Proceedings of the workshop on Microbial Degradation of Pollutants in Marine Environments. Pp. 423-431 Gulf Breeze: Environmental protection Agency.
- Quillardet, P., Huisman, O., D'Ari, R. and Hofnung, M. 1982. SOS Chromotest, a direct assay of an SOS function in *Escherichia coli* K-12 to measure genotoxicity. *Proceeding of the National Academic of Science, USA* 79, 5975-5975
- Smith, L.R. and Dragun, J., 1984. Degradation of volatile chlorinated aliphatic priority pollutants in groundwater. *Environment International* 10: 291-298.
- Teh, J.S., 1974. Toxicity of short-chains fatty acids and alcohol's towards *Cladosporium resinae*. *Applied Microbiology*, 28: 840-844.
- Teh, J. S. and Lee, K. H., 1974. Effects of n-alkane on *Cladosporium resinae*. *Canadian Journal of Microbiology* 20: 971-976
- Vandermeulen, J. H. 1986. Altered grazing patterns of an experimental copepod-alga ecosystem exposed to naphthalene and Kuwait crude oil. *Bulletin of Environmental contamination and Toxicology* 36: 260-266
- Vandermeulen, J. H and Lee, R. W., 1986. Lack of mutagenic activity of crude and refined oils in the unicellular alga *Chlamydomonas reinhardtii*. *Bulletin of Environmental contamination and Toxicology*, 36: 250-2553.
- Wang, W. 1984. Time response of *Nitrobacter* to toxicity *Environment International*, 10: 21-26
- Williamson, K. J and Johnson, D, G. 1981. a bacteria bioassay for assessment of wastewater toxicity. *Water Research*, 15: 383-390
- Xu, H.H. and Schurri, K.M., 1990. Genotoxicity of 22 pesticides in microtitration SOS chromotest *Toxicity Assessment, 1-4*