

## SEDIMENTOLOGICAL AND RESERVOIR DESCRIPTION OF AFOWO OIL SAND DEPOSITS, SOUTH-WESTERN NIGERIA

A. AKINMOSIN<sup>+</sup>, O.S. OREDEIN and A.A. ODEWANDE

Earth Sciences Department, Olabisi Onabanjo University, Ago-Iwoye, Ogun State, Nigeria.

(Submitted: 30 May 2005; Accepted: 04 May 2006)

### Abstract

Particle size distribution and mineralogical analysis of Afowo oil sands were studied with the aim of characterizing the reservoir vis-à-vis bitumen saturation and recovery efficiency. Sedimentologically, transportation history of the sediments, their maturity and the energy regime of the depositional environment were equally considered. A total of 44 tar sand samples were collected at three (3) different localities in Okitipupa area of Ondo State for these studies. These localities are: Idiopopo, Idiobilayo and Igbotako.

Results of sedimentological and particle size distribution studies showed that on the average, these sands are medium grained, moderately sorted and mesokurtic. The grain morphologies are of low to high sphericity with shapes generally sub-angular to sub-rounded. However, Igbotako samples are poorly sorted with some quantities of fines content, while samples of Idiobilayo and Idiopopo are better sorted with insignificant quantities of fines. The tar sands are both chemically and mechanically stable. The quartz content is on the average about 95 percent. Minor mineralogical components are mica (biotite and muscovite) and heavy minerals (including zircon, tourmaline, sillimanite, staurolite e.t.c.).

There is a marked correlation between mineralogical as well as geochemical composition of the samples analyzed. Percentage composition of SiO<sub>2</sub> was found to be over 90 percent. Other oxides (e.g. MgO, K<sub>2</sub>O, FeO, etc.) were found to have very low proportion.

From the overall results, it can be deduced that Afowo sands might have been deposited in a moderate energy environment, possibly lagoonal. Moreover, considering reservoir quality, Idiobilayo and Idiopopo sands should be expected to have better reservoirs with higher bitumen saturation and good oil recovery efficiency due to their low content of fines particles.

### 1. Introduction

The Dahomey basin (Fig. 1) is a marginal pull-apart basin (Klemme, 1975) or Margin sag basin (Kingston *et al.*, 1983), which was initiated during the early Cretaceous separation of African and South American lithospheric plates.

Occurrence of seepage and tar sand deposits over the Okitipupa ridge in the Dahomey basin provided the initial impetus for oil exploration in Nigeria. From the turn of 19<sup>th</sup> century up till date, no less than over twenty groups comprising public and private ventures have shown degrees of interest.

These bituminous deposits outcrop along an east-west belt, which is approximately 120 km long and 4-6 km wide, extending from the boundary of Edo and Ondo to Ogun State, (Enu, 1987), see figure 2. A total reserve of heavy oil is estimated to be over 30 billion barrels (Adegoke *et al.*, 1980).

The occurrence of these deposits has been known since early last century, however, intense investigations commenced from mid 70's till now.

The pioneering efforts were initiated by the Geological Consultancy Unit of the University of Ife (now Obafemi Awolowo University). The geology of these deposits, oil saturation and reserve estimates as well as textural characteristics of the associated sands have been described (Adegoke *et*

*al.*, 1980; 1981, Enu, 1987). The physicochemical properties of the bitumen in relation to production and processing have been studied (Adegoke *et al.*, 1980; Oshinowo *et al.*, 1982; Ekweozor, 1985; Oluwole *et al.*, 1985). The origin of the bitumen has been considered (Coker, 1982; Ekweozor, 1985). Other relevant studies on the deposit include works done by Ako *et al.* (1983); Ekweozor (1986 and 1990); Ekweozor and Nwachukwu (1989); Enu (1987, 1990); Enu and Adegoke (1984). These works have highlighted relevant aspects of the geochemical and sedimentological characteristic of the deposit.

It is important to determine the reservoir characteristics (such as grain size parameters) in order to assist with future production procedures. This paper is based on detailed studies of 44 surface tar sand samples and will mainly emphasize the particle size distribution and mineralogical aspects.

### 2. Review of the stratigraphy of the Dahomey basin

The study area lies between latitudes 6°30' to 6°35' and longitudes 4°30' to 6°45' and falls within the eastern Dahomey Basin (Fig. 3). The work of Omatsola and Adegoke (1981) on the Cretaceous stratigraphy of the Dahomey basin has recognized




Figure 1: East-West geological section showing the Dahomey Basin and upper part of the Niger Delta (After Whiteman, 1982).




Figure 2: Geological map of southwestern Nigeria showing the Tar sands outcrop belt (Adapted from Enu, 1987)




Fig 3: Location Map of the Study area

three formations belonging to the Abeokuta group. These are: the Ise Formation, consisting essentially of continental sands, grits and siltstones, overlying the basement complex unconformably. Neocomian to Albian age has been assigned to this formation. Overlying the Ise Formation is the Afowo Formation, which consists of coarse to medium-grained sandstones with variable interbeds of shales, siltstones and clay. The sediments of this formation were deposited in a transitional to marginal marine environment. Turonian to Maastrichtian age has been assigned to this formation. The Araromi Formation consists essentially of sand, overlain by dark-grey shales and interbedded limestone and marls with occasional lignite bands. The formation conformably overlies the Afowo Formation and Maastrichtian to Paleocene age has been assigned (Omatsola and Adegoke, 1981).

Overlying the Abeokuta group conformably is the Imo group, which comprises of shale, limestone and marls. The two-lithostratigraphic units under this group are: Ewekoro and Akinbo Formations. Ewekoro Formation consists of thick fossiliferous limestone. Adegoke (1977) described the Ewekoro

Formation as consisting of shaly limestone 12.5m thick which tends to be sandy and was divided into three microfacies. Ogbe (1972) however, further modified this and proposed a fourth unit. The four microfacies making up this formation are described as follows: sandy biomicrite, shelly biomicrite, algal biosparite, and red phosphatic biomicrite. Ewekoro Formation is Paleocene in age and is associated with shallow marine environment due to abundance of coralline algae, gastropods, pelecypods, echinoid fragments and other skeletal debris. Akinbo Formation lies on the Ewekoro Formation and it comprises of shale, glauconitic rock bank, and gritty sand which is pure grey in colour and shows little clay. Lenses of limestone from Ewekoro Formation grades laterally into the Akinbo shale very close to the base. The base is characterized by the presence of a glauconitic band. The age of the formation is Paleocene to Eocene.

Overlying the Imo group is the Oshoshun Formation. It is a sequence of mostly pale greenish-grey laminated, phosphatic marls, light grey to white-purple clay with interbeds of sandstones. It also consists of claystone underlain by argillaceous

**Table 1:** Mean particle-size parameters of the three localities.

	LOCALITYA	REMARK	LOCALITYB	REMARK	LOCALITYC	REMARK
Number of samples	13		14		17	
Mode	1.56		1.58		2.29	
Mean	1.03	Medium Sand	0.80	Coarse Sand	1.43	Medium Sand
Standard deviation	0.92	Moderately Sorted	0.95	Moderately Sorted	1.11	Poorly Sorted
Skewness	0.02	Nearly Symmetrical	-0.71	Strongly Coarse Skewed	-0.04	Strongly Coarse Skewed
Kurtosis	1.04	Mesokurtic	-0.07	Mesokurtic	-0.04	Mesokurtic

**Table 2:** LOCALITY A

SAMPLE NO	QUARTZ	FELDSPAR	MICA	ACCESSORY MINERALS
A1	95.5	TR	1.5	3.0
A2	96.0	TR	1.5	2.5
B1	96.5	TR	1.0	2.5
B2	95.5	-	2.0	2.5
E3	94.0	-	2.0	4.0
F1	96.0	TR	1.5	3.5

**Table 3:** LOCALITY B

SAMPLE NO	QUARTZ	FELDSPAR	MICA	ACCESSORY MINERALS
N1	96.5	—	2.0	1.5
N2	97.5	—	2.0	0.5
O5	96.0	TR	2.0	2.0
O6	97.0	—	1.5	1.5
P3	95.0	TR	2.5	2.5

**Table 4:** LOCALITY C

SAMPLE NO	QUARTZ	FELDSPAR	MICA	ACCESSORY MINERALS
Q3	94.5	TR	2.0	3.5
S1	94.0	TR	1.5	4.5
T1	93.0	TR	2.0	5.0
T3	93.5	TR	2.0	4.5
U6	95.0	—	1.5	3.5

Note: TR- TRACES.

limestones with light grey shale at the bottom. There are inclusions of phosphatic and glauconitic materials in the lower part of the formation and the upper part is made up of medium to coarse-grained silty sandstone (Adegoke, 1969).

Eocene in age (Agagu, 1985). The sedimentation of the Oshoshun Formation was followed by a regression, which deposited the sandstone unit of

Ilaro Formation (Kogbe, 1976). The sequence represents mainly coarse sandy estuarine deltaic and continental beds, which show rapid lateral facies change.

The coastal plain sands are the youngest sedimentary unit in the eastern Dahomey basin. It probably overlies the Ilaro Formation unconformably, but convincing evidence as to this is lacking (Jones and




Fig 4A: Histogram for Particle size distribution in Location A




Fig 4B: Histogram for particle size distribution in Locality B




Fig 4C: Histogram for particle size distribution in Locality C




Fig 5B: Cummulative Frequency plot for Locality B




Fig 5C: Cummulative Frequency plot for Locality C




Fig. 5C: Cummulative Frequency plot for Locality C



Hockey, 1964). It consists of soft, poorly sorted clayey sand and pebbly sands. The age is from Oligocene to Recent.

### 3. Methodology

A total number of forty-four tar sand samples collected at outcrop points and along river channels in 3 different localities of tar sands area of Ondo state were used for this study. These localities include: Idiobilayo (13 samples), Idiopopo (12 samples) and Igbotako (12 samples).

After cleaning and drying, the samples 100g each were dry-sieved using vibrating sieving machine, [Endecott, London (E.F.L 2MK11)] for 15 minutes with sieves of the following mesh sizes: 2.00mm, 0.180mm, 0.125mm, 0.075mm which are equivalent to -1.00, 1.25, 2.50, 3.00 and 3.75 phi values respectively.

The individual weight, cumulative weight, as well as individual and cumulative weight percentage were determined. These were subsequently used to plot histogram and cumulative frequency curve for individual sample on an arithmetic and semi-logs graphs.

The parameters were computed by using moment statements: mean, mode, standard deviation, Kurtosis, and Skewness (using formulae by Folk and Ward, 1957). Besides the particle-size distribution, grain morphology, chemical and mineralogical

compositions were equally determined using standard procedures.

For the bitumen saturation analysis, ten(10)grams each of samples were weighed and put in a measuring cylinder into which toluene was poured and left for about 60 minutes. The samples were then washed and later decanted. This procedure was repeated until the samples were clean of bitumen. The washed samples were air dried and re-weighed. The new weights were noted, recorded and then subtracted from the initial weights. The difference in weight were converted to percentage(%) and recorded. An average for all the samples analysed for each facies type was taken as average saturation for the respective tar sand.

### 4. Results and Discussion

Mean and standard deviation of particle-size distribution of Afowo bituminous sands are presented in Table 1. The particle-size frequency-distribution curves of almost all the sediments analyzed indicate bimodal distributions. The curves show substantial variation from one another. The sediments show wide broad, moderately negatively skewed curves and is reflected in the histograms (Figs. 4 and 5) which indicate a wide range of relatively higher values for the standard deviation. The sediments in Idiobilayo and Idiopopo are better sorted than the sediments in Igbotako as it can be seen in Table 1. While sediments from Idiobilayo and Igbotako are medium grained,

## ELEMENTAL COMPOSITION OF SAMPLES FROM THE LOCALITIES.

Table 5: LOCALITY A

SAMPLENO	CaO	FeO	S <sub>1</sub> O <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	MgO	SO <sub>3</sub>	K <sub>2</sub> O
A1	-2.23	0.28	94.19	-5.56	2.49	0.01	0.09
A2	-2.15	0.36	93.25	-5.45	2.30	0.02	0.06
B1	-2.35	0.18	93.60	-5.40	2.60	0.04	0.08
B2	-2.11	0.33	93.16	-5.24	2.35	0.00	0.10
F1	-2.30	0.35	94.45	-4.30	2.60	0.03	0.09

Table 6: LOCALITY B

SAMPLE NO	CaO	FeO	S <sub>1</sub> O <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	MgO	SO <sub>3</sub>	K <sub>2</sub> O
N1	-1.95	0.09	93.37	-4.94	2.12	0.10	0.12
N2	-1.80	0.07	94.41	-4.76	2.31	0.09	0.41
O5	-1.87	0.06	96.42	-4.85	2.20	0.08	0.13
O6	-2.13	0.11	95.01	-4.90	2.41	0.12	0.14
P3	-2.16	0.16	96.45	-4.76	2.53	0.16	0.12

Table 7: LOCALITY C

SAMPLE NO	CaO	FeO	S <sub>1</sub> O <sub>2</sub>	Al <sub>2</sub> O <sub>3</sub>	MgO	SO <sub>3</sub>	K <sub>2</sub> O
Q3	-1.70	0.85	92.46	-2.99	2.25	0.12	0.03
S1	-1.15	0.90	93.50	-2.78	2.30	0.15	0.01
T1	-1.65	0.87	91.50	-2.20	2.45	0.17	0.04
T3	-1.95	0.30	94.24	-4.04	2.28	0.12	0.02
U6	-1.90	0.60	93.25	-3.05	2.38	0.16	0.03

Idiopopo sediments are however coarse grained all with variable quantities of fines.

The Afowo sands as observed in this study, are of the mean particle size that is coarse-medium and does not depend too much on standard deviation or percentage of fines (Figs. 6a-6c). Harison *et al.* (1981) in the cold Lake area showed that a strong correlation exists between mean particle size and weight (wt) % mud (<63¼ m). Coarse particles and low mud content allow for high bitumen content. A factor that could prove very important in the recovery of bitumen is the heterogeneity of a reservoir, because the withdrawal efficiency is strongly reduced if clusters of fines occur in a coarser matrix (Morrow, 1971; Takamura, 1982).

The reservoir is expected to be better off in the sediments of Idiobilayo and Idiopopo, which are coarse grained and better sorted than the sediments in Igbotako area.

Mineralogical composition of the sediments as identified under petrological microscope include quartz, mica and opaque minerals constituting 95%, 2% and 1% respectively. No feldspar was identified under the microscope, which can be indicative of mechanical maturity of the sediments, as shown on Tables 3, 4 & 5. The grain morphology shows low-high sphericity with shapes generally sub-angular to sub-rounded (Fig. 7).

Heavy mineral constituents of the sediments comprises of both opaque and non-opaque types. These minerals are of great importance in the study of provenance,

Minerals identified under the microscope are zircon, rutile, tourmaline, staurolite, sillimanite, kyanite, garnet and hornblende, (Figs. 7a and 7b). Average ZTR index calculated for some representative samples is greater than 75%.

Chemical composition of the samples as shown on Tables 5, 6 and 7, reveals SiO<sub>2</sub> as most abundant, averaging about 95% of the total oxides. This is followed by MgO, FeO and K<sub>2</sub>O in that order. As observed from the mineralogical composition, the proportion of these oxides is indicative of the chemical maturity of the sediments, which makes it possible to classify the sediments as quartz arenites. Moreover, with high proportion of SiO<sub>2</sub> and low proportion of K<sub>2</sub>O, the sediments can better be classified as arenaceous in nature (quartz arenite).

### 5. Summary and Conclusion

The bitumen sands of Afowo Formation as observed in this study can be characterized as being chemically and mechanically stable. The quartz content on the average is greater than 90% with little or no presence of feldspar. Minor components are mica (biotite and muscovite) and heavy minerals (mostly zircon and tourmaline).

The particle size of the sands varies irregularly. Mostly, the sands are coarse-medium grained and moderately sorted with no significant of fines-content. Except for few locations with poor sorting, bulk of the sediments are moderately sorted. This quality, coupled with insignificant fines content will in no doubt enhance both bitumen saturation and withdrawal efficiency positively.

### REFERENCES

- Adegoke, O.S., 1969. Eocene Stratigraphy of Southern Nigeria. Bulletin Bureau de Recherche Geologie ET Miners Memoirs. Vol. 69, pp. 22-24.
- Adegoke, O.S., 1977. Stratigraphy and paleontology of the Ewekoro Formation (Paleocene) of SW Nig. Bull. Am. Paleontol; vol.71, No. 295, 375pp.
- Adegoke, O.S., Ako, B.D. and Enu, E.I., 1980. Geotechnical investigations of the Ondo State bituminous sands. Vol. 1. Geology and reserve estimate. Rept. Geological Consulting Unit, Dept. of Geology, University of Ife. 257pp.
- Adegoke, O.S., Enu, E.I., Ajayi, T.R., Ako, B.D., Omatsola, M.E. and Afonja, A.A., 1981. Tar sands-A new energy raw material in Nigeria. Proc. Int. Symp. on new energy raw material. Karlovy Vary: 17-22.
- Agagu, O.K., 1985. A geological guide to bituminous sediments in Southwestern Nigeria. Unpubl. Rept. Dept. of Geology, University of Ibadan.
- Ako, B.D., Alabi, A.O., Adegoke, O.S. and Enu, E.I., 1983. Application of resistivity sounding in the exploration for Nigerian Tar Sand: Energy exploration and exploitation, Vol. 2, No. 2. 155-164.
- Coker S.J.L., 1982. Some aspects of the geology of the bituminous sands of parts of the Benin Basin. Nig. Min. and Geosc. Soc. 19, 121pp.
- Ekweozor, C.M., 1985. Nigerian Tar Sands. Bitumen chemical properties and their relation to origin, production and processing. Book of Abstracts: 21<sup>st</sup> Ann. Conf. Nig. Min. & Geosc. Soc., Jos Nigeria, 11<sup>th</sup>-15<sup>th</sup> March 1985.
- Ekweozor, C.M., 1986. Characteristics of the non-asphaltene products of mild chemical degradation of asphaltene. Org. Geochem. 10, pp. 1053-1058.
- Ekweozor, C.M., 1990. Geochemistry of oil sands of southwestern Nigeria In: B.D. Ako, E.I. Enu (editors). Occurrence, utilization and economic of Nigerian tar sands: A workshop held in Ogun State University Ago-Iwoye, Nigeria on 29-31 May, 1990. Published by the Nigerian Mining and Geosciences Society, Ibadan chapter. 50-62.
- Ekweozor, C.M. and Nwachukwu, J.I., 1989. The origin of Tar sands of southwestern Nigeria. NAPE Bull. 4, 80-94.
- Enu, E.I. 1987. The paleoenvironment of deposition of late Maastrichtian to Paleocene black shales in the eastern Dahomey basin, Nigeria Geologie en Mijnbouw. 66 pp. 15-20.
- Enu, E.I., 1990. Textural and occurrence of tar sands in Nigeria In: B.D. Ako, E.I. Enu (editors), Occurrence, utilization and economic of Nigerian tar sands: A workshop held in Ogun State University Ago-Iwoye, Nigeria on 29-31 May, 1990. Published by the Nigerian Mining and Geosciences Society, Ibadan chapter, 11-16.
- Enu, E.I. and Adegoke, O.S. 1984. Potential industrial mineral resources associated with the Nigeria tar sands. 27<sup>th</sup>






Fig. 6a: Mean particle size vs standard deviation for locality A.




Fig. 6b: Mean particle size vs standard deviation for locality B.




Fig. 6c: Mean particle size vs standard deviation for locality C




Fig. 7: Quartz and other mineralogical composition of the sediments showing the shapes of individual mineral grain.




Fig. 7a



**Mag X 100**  
Fig 7b: High quality of Zircon and Staurolite

Z = Zircon  
 K = Kyanite  
 Si = Sillimanite  
 S = Staurolite  
 O = Opaque mineral  
 T = Tourmaline  
 R = Rutile

- international geological congress Mosco. Book of abstract. Vol. Vii, pp. 345.
- Folk, R.L. and Ward, W.C. 1957. Brazo river bar: A study the significant of grain size paraeter. *Journal sed. Petrol.* 41, pp. 45-1058.
- Harrison, D.B., Glaister, R.P. and Nelson, H.W. 1981. Reservoir description of the Clearwater oil sand, Cold Lake, Alberta, Canada, in *The future of heavy crude and tar sands*: New York, McGraw-Hill, pp. 264-279.
- Jones, M.A and Hockey, R.O. 1964. *The Geology of Part Of South Western Nigeria*. Nig. Geol. Surv. Bull. No 31, 1-101.
- Kingston, D.R., Dishroon, C.P. and Williams, P.A. 1983. Global basin classification system. *American Association of Petroleum Geologists. Bulletin* 31, 87pp.
- Kogbe, C.A.. 1976. *Geology of Nigeria*. (Second revised edition). Published by Rock View (Nig.) 455pp. C.A Kogbe (editor).
- Morrow, N.R., 1971. Small-scale packing heterogeneities in porous sedimentary rocks: *Bulletin of the American Association of Petroleum Geologists*, vol. 55, pp. 515-520.
- Ogbe, F.G.A., 1972. Stratigraphy of Strata Exposed In Ewekoro Quarry, Southwestern Nigeria. In: *Africa Geology* pp. 305-322.
- Oluwole, F.F., Adegoke, O.S., Kehinde, L.O., Borishade, A.B., Ako, B.D., Ajayi, T.R., Enu, E.I., Nwachukwu, J.I., Coker, S.J.L., Wallace, D., Asubojo, O.I and Ogunsola, O. 1985. Chemical composition of bitumen extracts from Nigerian tar sands. In: *Proceedings of 3<sup>rd</sup> Intern. Confer. On heavy Crude and Tar sands*; Long Beach California; chap. 33, pp 373-379.
- Omatsola, M.E and Adegoke, O.S., 1981. Tectonic and Cretaceous Stratigraphy Of Dahomey Basin *Jour. Of Min. Geol.* Vol.54, pp. 65-87.
- Oshinowo, T., Ademodi, B. and Adediran, A.S., 1982. Bituminous tar sand of Nigeria. *Analysis of oils-Part 1. Jour. Nig. Soc. Chem. Engrs.* 1, 44-48.
- Takamura, K.. 1982. Microscopic structure of Athabasca oil sand: *Canadian Journal of Chemical Engineering*, vol. 60, pp. 538-545.
- Whiteman, A.J., 1982. *Nigeria: Its Petroleum Geology, Resources and Potential*, Vol. 2, Graham and Trotman, London.