

*Commentary*

## **Pale Europeans and Dark Africans share sun and common health problems: constancy in regional health differences and sunshine**

**Timo Töysä<sup>1</sup> and Osmo Hänninen<sup>2</sup>**

*Specialty General Practice, Rehabilitation Hospital Vetrea, Pohjolankatu 15, FI-74100 Iisalmi, and <sup>2</sup>Department of Physiology, University of Eastern Finland, FI-70211 Kuopio, Finland*

*CHD mortality, regional risk, sunshine, inflammation, silicon*

### **ABSTRACT**

Regional risk of cardiovascular mortality has only recently been added to the group of major risk factors, but its effective sub-factors are not fully understood. The aim of this study was to assess the CHD mortality in Finland and other Europe, its stability and association with capability in vitamin D synthesis. In a worldwide assay several causes besides geographical and climatological factors restrict vitamin D synthesis in skin and several factors affect the activation as also the inactivation of vitamin D. Sunshine anyhow is the constant source of vitamin D, while dietary habits have changed. Effects of vitamin D could be based on mineral, anti-inflammatory and structural factors. Different availability of sunshine has thus associated with the constancy in proportional difference in CHD mortality in Finland as an example country. Conclusion: Sunshine is associated with lower CHD mortality in Europe. Other additive mechanisms are suggested.

© Copyright 2014 African Association of Physiological Sciences -ISSN: 2315-9987. All rights reserved

### **INTRODUCTION**

Usually, between 50% and 90% of vitamin D in the body is coming from the production in the skin and the remainder is from the diet (Lips 2010). Finland and South-Africa are at different ends of the same time zone. Both depend on sun radiation in their vitamin/hormone D synthesis. Miyauchi et al. (2013) reported that latitude is associated with lower vitamin D synthesis in skin. It is possible to suggest which diseases are caused by the low Vitamin D3 levels by looking at the geographical distribution and incidence of the diseases. Most of them are much reduced at the equator and much higher in the extreme latitudes (Plotnikoff and Quigley 2003). They can be Vitamin D3 related. Beneficial effect of moderate sunshine can

be based on psychological well-being, too (Gannagé-Yared et al. 2001).

Previously people worked in rural occupations mainly outdoors and got more sun radiation. At present most people live in cities and are thus protected from sunshine by housing, vehicles used (also the dust generated) and clothing. During the past half century several studies have been performed and health behavior has changed in Finland, but still the difference in male coronary mortality is higher in Eastern than Western Finland. In South Africa significant numbers of people having dark skin have lower vitamin D levels than the light colored ones, while in whites no such difference exists. Dark pigmentation hinders the satisfactory vitamin D synthesis although it protects against sun burns. Now many descendants of African Americans have lower D blood levels (Hall et al. 2010). Unfortunately these are also global health problems as many people get less sun radiation, are also less physical active outdoors but who also have excessive and improper food intake. Big cities have also serious air pollution that hinders the sun radiation to reach people at street level. Furthermore the television watching helps people to escape sunshine during their leisure time.


Address for correspondence:  
Email [osmo.hanninen@uef.fi](mailto:osmo.hanninen@uef.fi)

Lips (2010) and Oren et al. (2010) have reported that Vitamin D deficiency was highly prevalent in India, China and Middle-East, less common in Japan, South-East Asia and North America and in Latin America, quite common in Africa, more common in Mediterranean area than in Nordic countries. Freisling et al. (2010) wrote that dietary intake of vitamin D was lower in Mediterranean area than in Norway, Sweden and Denmark.

### North Europe

The people of Eastern Finland, living in Northern Europe and higher hills have suffered more from male coronary heart disease and other diseases than people in the lower of western parts of the country since the 19<sup>th</sup> century (Kannisto, 1947). Regional mortality risk has been affirmed by Jousilahti et al. (1998) who wrote that CHD risk was about 40% higher in the north-eastern than in south-western Finland. Differences in major and minor risk factor levels explained only about 40% of the excess CHD risk in eastern Finland. This higher CHD mortality in eastern Finland (see Fig1) was quite constant in 1973-2000 (Pajunen et al. 2004).

In Fig1 we see that western Finland is located to the south from eastern Finland. The eastern Finland is positively associated with the number of rainy days (Fig2), i.e. with lower capability for synthesis of vitamin D.



**Fig1.** The oblique thick line divides as Pajunen et al. (2004) the Eastern Finland (with high CHD and general mortality) from the west.

**Fig 2.** Mean annual rainy days ( $\times 0.1$  mm/d) in 1981-2010, Finnish Meteorological institute (FMI). Added gross lines for 180 and 195 days.

In Finland 1951-72 the CHD mortality was lower in countryside than in urban areas (Fig3), which could be associated with the higher outdoor life especially in farming and forestry. Same time when number of people was 464.7 million (Statistics Finland 1978), the number of horses, used in farming and forestry, were reduced from 350,000 to 50,000 (Statistics Finland 1960 & 1978) and replaced to tractors, which got cabins, lumberjacks got helmets and outdoor work was in general reduced, all of these affecting vitamin D synthesis. Lamberg-Allardt et al. (1986) reported that vitamin D was trendlike higher in western than in eastern and northern Finland, trendlike higher in countryside than in urban areas

### South Europe


Residence in a low risk country (i.e. Mediterranean area, including Switzerland and Belgium) decreases the risk of cardiovascular mortality (including CHD) of smokers to the same level as by non-smokers in another country of Europe, if the other main risk factors are the same (Graham et al. 2007). In the southern Europe lower latitude is associated with lower number of annual number of rainy days than in other parts of Europe in general (van Engelen et al. 2014) increasing the capability for D-vitamin synthesis in the skin.

### South Africa


Studies show that the vitamin D-level is low in those Africans who have a lot of dark pigment in their skin. Some of them thus suffer even vitamin D-deficiency (Poopedi et al. 2009, Martineau et al. 2011).

Urbanization is also going on in South Africa which happened earlier in Europe. Kruger et al. (2011) have reported that it may increase the health problems in South Africa. Black women may have low bone mass due to low vitamin D status, low calcium intake, and high bone turnover. Furthermore the indoor living and television watching inhibit of course the natural sun radiation exposure.

In Africa there are also other reasons which can affect the satisfactory vitamin D synthesis in skin. Mining is important and the underground work limits sun radiation. Also mining dust in air prevents the satisfactory sun radiation. Also traffic and heating mean also dust pollution.



**Fig3.** Age adjusted CHD mortality of middle-aged men in rural and urban areas of Finland. (Valkonen and Martikainen 1990)



**Fig. 4.** The balance of vitamin/hormone D between the intakes in foods, supplements and the syntheses in skin, liver and kidneys and the storage and dilution in body lipids, adipose tissue especially (vitamin D is fat-soluble vitamin) and catabolism and final disposure by the liver into bile.

Deficiency of vitamin D means increases the risks of HIV and Tuberculosis (Martineau et al. 2011). The vitamin-D supplementation is motivated to prevent the problems of deficient vitamin levels in those whose dark skin may prevent the satisfactory endogenous D3 production.

## **Discussion**

It is generally known that Vitamin D works against rachitis by acting on mineral uptake. It has been suggested that vitamin D protects against several other chronic conditions like diabetes, and some cancers, all of which are prevalent or more prevalent among blacks than whites (Harris 2006). Vitamin D is needed for insulin secretion (Borisova et al. 2003), glucose tolerance and lipid metabolism. Insulin receptor gene promoter is sensitive to vitamin D (Maestro et al. 2003). Kositsawat et al. reported inverse correlation between serum 25(OH)D3 and plasma HbA1C. Recently in a prospective study vitamin D supplementation has decreased HbA1 level in African Americans (Green et al. 2014).

Experimentally vitamin D3 has shown anti-inflammatory action on the decrease of MDA and hs-CRP (Sharifi et al. 2014) and anti-infectious action (Li et al. 2014). Garcia et al. (2003) have reported that elevated PTH can increase LVH, hypertension and CVD death. Martins et al. (2014) wrote that experimentally vitamin D has decreased arterial stiffness and PTH level. Anyhow effects on CHD need further studies (Schnatz and Manson, 2014).

Vitamin D is lipophilic and is therefore distributed in adipose tissues (Fig4). Its turnover is slow (half-life approximately 2 mo). Its main transported metabolite, 25(OH)D3, shows a half-life of approximately 15 h while hormone 1,25(OH)2D has a half-life of approximately 15 h (Jones 2008).

The negative effect of human activity on air quality has been evident ever since the beginning of the industrialization in big cities. The traffic, heating and industries themselves release dust particles as also various chemicals released interact in air. Over 90% of European Union population lives in area where the recommendations of WHO air quality are exceeded, especially due to particles PM2.5 and 10 and ozon (O3) (EEA 2013). In Chinese megacities one can see sun shining from the cloudless sky above but one cannot properly see horizontally more than some 300 meters in the streets and you sense it your lungs (OH in Beijing personal experience).

In cities open swimming pools and beaches are rare and distant as also the parks of any significant size. Thus the possibilities to get sun radiation in any larger skin area are limited. People having dark skin in cities get less sun radiation than other Americans.

Obese people are overwhelmingly more deficient in lipid soluble Vitamin D (Botella-Carretero et al. 2007). Low levels of 25(OH)D are seen in fatty liver diseases independent on obesity (Hao et al., 2014, Nobili et al. 2014). Good part of Vitamin D stores of different components are located in adipose tissue and thus diluted by other lipids of the body. The mass of adipose tissues competes easily with the forms obtained in foods and produced in the skin, liver and kidneys. Obesity is also associated with low physical inactivity. Further possibility is too much tasty energy rich foods available, even selenium deficiency, too (Kim et al. 2012). There is discrepancy concerning vitamin D causality in obesity: Although 1,25(OH)2D inhibits adipogenesis in some mouse experiments, it can increase lipid accumulation in human subcutaneous preadipocytes (Nimitphong et al. 2012).

Active Vitamin D is metabolized by enzymes of the cytochrome P-450 family whose activity is modifiable. In general the first step, 25-hydroxylation to 25(OH)D, occurs in liver via CYP2R1 (Cheng et al. 2004) and then 1-hydroxylation in kidney via CYP27B1 (Omdahl et al. 2003). Degradation of 1,25(OH)2D and (and obviously of 25(OH)D occurs via cytochrome P450C24 (CYP24) (Wietrzyk 2007).

Alpha-tocopherol has been reported to regulate vitamin D3 25-hydroxylation (Velyky 2010). Mg-deficiency can potentiate peroxide stress by increasing lipid peroxidation and ATP-decrease (Manju and Nair 2006). Decreased ATP availability could decrease liver function and 25-hydroxylation.

Rude et al. (2006) wrote that reduction of Mg intake of 50% from nutrient requirement reduced 1,25(OH)2D formation, as well as increased release of the inflammatory cytokines TNF-alpha and IL-1beta. Matsuzaket al. (2013) reported that Mg deficiency induced downregulation of 1(OH)ase and upregulation of 24(OH)ase in the kidney, i.e. reduced 1,25(OH)2D synthesis and increased degradation (of 1,25(OH)2D and 25(OH)D). Wietrzyk (2007) wrote that isoflavonoids inhibit activity of CYP24, so inhibiting reduction of 25(OH)D and 1,25(OH)2D.

Noyola-Martínez et al. (2014) reported that INF- may contribute to calcitriol production while TNF- favors its catabolism in human trophoblasts. Lehmann and Maurer (2010) have reported that irradiated skin is able to produce calcitriol in substantial amounts, too. Li et al. (2008) observed 1,25(OH)<sub>2</sub>D synthesis and degradation in adipose tissue.

Availability of Vitamin D is globally not enough (Lips 2010). Serum levels of 25(OH)D reflect not only its availability, but effects of proactive and counteractive factors (Fig4.): Low levels of 25(OH)D in fatty liver diseases (Hao et al., 2014, Nobili et al. 2014) could reflect the activity of vitamin D 25-hydroxylase (Velyky et al. 2010). Relatively elevated CYP24 activity has reported to explain low vitamin D deficiency in gestational DM (Gho et al. 2013). On the other hand cytokines can control vitamin D status: INF- may contribute to 1,25(OH)<sub>2</sub>D production (via CYP27B1) while and TNF- favors its catabolism (via CYP24A1 gene expression) (Noyola-Martínez et al. 2014). Cf. Memon et al. (1993) reported that TNF-alpha and IL-6 can increase cholesterol synthesis more than 2-fold.

25(OH)D does not fully reflect the amount of 1,25(OH)<sub>2</sub>D because of the above mechanisms. The number of large scale vitamin D analyses are scanty. Dietary assessments forget sometimes the sunshine, which has been suggested to be the main route of vitamin D production (Lips 2010). Earlier the East-West difference in CHD mortality has been explained by significantly higher drinking-water silicon content and is structural characteristics in West Finland (Schwarz et al. 1977). Vitamin D can also increase collagen stability (Nagaoka et al. 2008), like silicon. Rabon et al. (1995) reported that vitamin D can increase more silicon absorption.

### Conclusions

It seems that the availability of Vitamin/hormone D<sub>3</sub> in humans is not necessarily optimal in South Africa, Europe and Finland as a special example. There are increasing data supporting that the Vitamin D is not only associated with mineral

uptake, but in immunological defense, diabetes and cardiovascular diseases and the same time one indicator of inflammation as cholesterol. Recently milk products have been supplemented with Vitamin D, why not the same in selected cases with food oils? Because the amount of active Vitamin D [1,25(OH)<sub>2</sub>D] is dependent on pro- and counteracting mechanisms, they should be monitored in epidemiological studies, too. Epidemiological studies are important not only for avoiding vitamin D deficiency, but toxicity, too. Because of deficiency of hard data concerning vitamin D effects on regional differences in CHD mortality in Europe, studies on silicon is suggested.

### References

- Air quality in Europe. European Environment Agency Report EEA 2013. 9/2013, 112 pp. ISSN 1725-9177.
- Borissova AM, Tankova T, Kirilov G, Dakovska L, Kovacheva R. The effect of vitamin D<sub>3</sub> on insulin secretion and peripheral insulin sensitivity in type 2 diabetic patients. *Int J Clin Pract* 2003; 57(4) 258-61.
- Botella-Carretero JJ, Alvarez-Blasco F, Villafruela JJ, Balsa JA, Vázquez C, Escobar-Morreale HF (2007). Vitamin D deficiency is associated with the metabolic syndrome in morbid obesity. *Clin Nutr.* 2007 Oct;26(5):573-80. Epub 2007 Jul 10.
- Cheng JB, Levine MA, Bell NH, Mangelsdorf DJ, Russell DW (2004). Genetic evidence that the human CYP2R1 enzyme is a key vitamin D 25-hydroxylase. *Proc Natl Acad Sci U S A.* 2004 May 18;101(20):7711-5. Epub 2004 May 5.
- Cho GJ, Hong SC, Oh MJ, Kim HJ (2013). Vitamin D deficiency in gestational diabetes mellitus and the role of the placenta. *Am J Obstet Gynecol.* 2013 Dec;209(6):560.e1-8. doi: 10.1016/j.ajog.2013.08.015. Epub 2013 Aug 14.
- EEA 2013. 9/2013, 112 pp. Air quality in Europe. European Environment Agency Report ISSN 1725-9177.
- van Engelen et al. (2014): European Climate Assessment & Dataset (ECAD). Map: <http://ecad.knmi.nl/utills/mapserver/climatology.php#bottom>, (accessed 28 April 2014) -->select: "Rain", "Wet days (RR >= 1 mm)", e.g. "1981-2010", "annual", Log scale "no". (van Engelen A et al, ECA&D Project Team, Royal Netherlands

- Meteorological Institute, P.O. Box 201, 3730 AE De Bilt, The Netherlands).
- Permission:  
[http://ecad.knmi.nl/documents/ECAD\\_datapolicy.pdf](http://ecad.knmi.nl/documents/ECAD_datapolicy.pdf)  
(free for scientific articles)
- Eng H, Mercer JB. Seasonal Variations in Mortality Caused by Cardiovascular Diseases in Norway and Ireland. *European Journal of Preventive Cardiology* April 1998 vol. 5 no. 2 89-95. doi: 10.1177/174182679800500204
- FMI (Finnish Meteorological Institute): <http://en.ilmatieteenlaitos.fi/normal-period> (Apr 4, 2014) --> "Precip. days ( $\times 0.1$  mm)ö, 1981-2010.
- Freisling H, Fahey MT, Moskal A, Ocké MC, Ferrari P, Jenab M, Norat T, Naska A, Welch AA, Navarro C, Schulz M, Wirfält E, Casagrande C, Amiano P, Ardanaz E, Parr C, Engeset D, Grioni S, Sera F, Bueno-de-Mesquita B, van der Schouw YT, Touvier M, Boutron-Ruault MC, Halkjaer J, Dahm CC, Khaw KT, Crowe F, Linseisen J, Kröger J, Huybrechts I, Deharveng G, Manjer J, Agren A, Trichopoulou A, Tsiotas K, Riboli E, Bingham S, Slimani N. Region-specific nutrient intake patterns exhibit a geographical gradient within and between European countries. *J Nutr.* 2010 Jul;140(7):1280-6. doi: 10.3945/jn.110.121152. Epub 2010 May 19.
- Gannagé-Yared MH, Tohmé A, Halaby G (2001). [Hypovitaminosis D: a major worldwide public health problem]. [Article in French]. *Presse Med.* 2001 Apr 7;30(13):653-8.
- Garcia de la Torre N, Wass JA, Turner HE. Parathyroid adenomas and cardiovascular risk. *Endocr Relat Cancer.* 2003 Jun;10(2):309-22.
- Graham I et al. *Eur J Cardiovasc Prev & Reh*, 14, Suppl 2, catalogue above on page E12 - E14 (2007).- <http://www.escardio.org/guidelines-surveys/esc-guidelines/Documents/PREV/guidelines-CVD-prevention-ES-FT-2007.pdf>
- Green RT, Gambhir KK, Nunlee-Bland G, Odonkor WA, Ganta VA (2014). Maintenance of long-term adequate levels of vitamin d lowers HbA1c in African American patients with type 2 diabetes. *Ethn Dis.* 2014 Summer;24(3):335-41.
- Hall LM, Kimlin MG, Aronov PA, Hammock BD, Slusser JR, Woodhouse LR, Stephensen CB. Vitamin D intake needed to maintain target serum 25-hydroxyvitamin D concentrations in participants with low sun exposure and dark skin pigmentation is substantially higher than current recommendations. *J Nutr.* 2010 Mar;140(3):542-50. doi: 10.3945/jn.109.115253. Epub 2010 Jan 6.
- Hao YP, Ma XJ, Luo YQ, Ni J, Dou JX, Hu YQ, Zhu JA, Bao YQ, Jia WP. Serum vitamin D is associated with non-alcoholic fatty liver disease in Chinese males with normal weight and liver enzymes. *Acta Pharmacol Sin.* 2014 Sep;35(9):1150-6. doi: 10.1038/aps.2014.48. Epub 2014 Aug 4.
- Harris SS. Vitamin D and African Americans. *J Nutr.* 136:1126-1129, 2006
- Jones G. Pharmacokinetics of vitamin D toxicity. *Am J Clin Nutr.* 2008 Aug;88(2):582S-586S.
- Jones G, Prosser DE, Kaufmann M. Cytochrome P450-mediated metabolism of vitamin D. *J Lipid Res.* 2014 Jan;55(1):13-31
- Jousilahti P, Vartiainen E, Tuomilehto J, Pekkanen J, Puska P. Role of the known risk factors in explaining the difference in the risk of coronary heart disease between eastern and southwestern Finland. *Ann Med* 30 (1998) 481-7. (ref in P. Pajunen, J. Torppa, M. Huohvanainen, V. Salomaa, V. Vartiainen. Sepelvaltimotautikuolleisuuden itä-länsiero kaventunut 30 vuoden seuranta-aikana miehillä. *Suomen Lääkärilehti* 51-52/2004 Vsk 59:5013-16.)
- Kannisto V. (1947) Kuolemansyyt vaestollisina tekijoina Suomessa. Vaitosk, Kansantaloudellinen yhdistys, Helsinki. (in *Karvonen* MJ. Prehistory of the North Karelia Project:15ô 18, in Puska P, Vartiainen E, Laatikainen T, Jousilahti P, Paavola M (editors). THE NORTH KARELIA PROJECT: FROM NORTH KARELIA TO NATIONAL ACTION. National Institute for Health and Welfare (THL), in collaboration with the North Karelia Project Foundation. ISBN 978-952-245-000-5 (nid.) ISBN 978-952-245-001-2 (PDF). Helsinki University Printing House. Helsinki 2009. TH
- Kim JE, Choi SI, Lee HR, Hwang IS, Lee YJ, An BS, Lee SH, Kim HJ, Kang BC, Hwang DY (2012). Selenium significantly inhibits adipocyte hypertrophy and abdominal fat accumulation in OLETF rats via induction of fatty acid -oxidation. *Biol Trace Elem Res.* 2012 Dec;150(1-3):360-70. doi: 10.1007/s12011-012-9519-1. Epub 2012 Oct 18.
- Kositsawat J, Freeman VL, Gerber BS, Geraci S. (2010). Association of A1C levels with vitamin D status in U.S. adults: data from the National Health and Nutrition Examination Survey. *Diabetes Care.* 2010 Jun;33(6):1236-8. doi: 10.2337/dc09-2150. Epub 2010 Mar 9.
- Kruger MC, Kruger IM, Wentzel-Viljoen E, Kruger A. Urbanization of black South African women may increase risk of low bone mass due to low vitamin D

- status, low calcium intake, and high bone turnover. *Nutr Res.* 2011; **31**(10): 748-58.
- Lamberg-Allard C, Ala-Houhal M, Ahola M, Parviainen MT, Räsänen L, Visakorpi J. Vitamin D Status of Children and Adolescents in Finland. *Ann Nutr Metab* 30:267-272 (1986). Study period September-December 1980.
- Lehmann B, Maurer M (2010). Vitamin D metabolism. *Dermatol Ther.* 2010 Jan-Feb;23(1):2-12. doi: 10.1111/j.1529-8019.2009.01286.x.
- Li P, Xu X, Cao E, Yu B, Li W, Fan M, Huang M, Shi L, Zeng R, Su X, Shi Y (2014). Vitamin D deficiency causes defective resistance to *Aspergillus fumigatus* in mice via aggravated and sustained inflammation. *PLoS One.* 2014 Jun 13;9(6):e99805. doi: 10.1371/journal.pone.0099805. eCollection 2014.
- Li J, Byrne ME, Chang E, Jiang Y, Donkin SS, Buhman KK, Burgess JR, Teegarden D (2008). 1 $\alpha$ ,25-Dihydroxyvitamin D hydroxylase in adipocytes. *J Steroid Biochem Mol Biol.* 2008 Nov;112(1-3):122-6. doi: 10.1016/j.jsbmb.2008.09.006. Epub 2008 Sep 17.
- Lips P (2010). Worldwide status of vitamin D nutrition. *J Steroid Biochem Mol Biol.* 2010 Jul;121(1-2):297-300. doi: 10.1016/j.jsbmb.2010.02.021. Epub 2010 Mar 1.
- Maestro B, Davila N, Carranza MC, Calle C. Identification of a vitamin D response element in the human insulin receptor gene promoter. *J Steroid Biochem Mol Biol*2003;84(2-3)223-30.
- Manju L, Nair RR. Magnesium deficiency augments myocardial response to reactive oxygen species. *Can J Physiol Pharmacol.* 2006 Jun;84(6):617-24.
- Martineau AR, Nhamoyebonde S, Oni T, Rangaka MX, Marais S, Bangani N, et al. Reciprocal seasonal variation in vitamin D status and tuberculosis notifications in Cape Town, South Africa. *Proc Natl Acad Sci U S A.* 2011; 108(47): 19013-7.
- Martins D, Meng YX, Tareen N, Artaza J, Lee JE, Farodolu C, Gibbons G, Norris K (2014). The Effect of Short Term Vitamin D Supplementation on the Inflammatory and Oxidative Mediators of Arterial Stiffness. *Health (Irvine Calif).* 2014 Jun;6(12):1503-1511.
- Matsuzaki H, Katsumata S, Kajita Y, Miwa M (2013). Magnesium deficiency regulates vitamin D metabolizing enzymes and type II sodium-phosphate cotransporter mRNA expression in rats. *Magnes Res.* 2013 Apr-Jun;26(2):83-6. doi: 10.1684/mrh.2013.0341.
- Memon RA, Grunfeld C, Moser AH, Feingold KR (1993). Tumor necrosis factor mediates the effects of endotoxin on cholesterol and triglyceride metabolism in mice. *Endocrinology.* 1993 May;132(5):2246-53.
- Miyauchi M, Hirai C, Nakajima H. The solar exposure time required for vitamin D<sub>3</sub> synthesis in the human body estimated by numerical simulation and observation in Japan. *J Nutr Sci Vitaminol (Tokyo).* 2013;59(4):257-63.
- Nagaoka H, Mochilda Y, Atsawasuwan P, Kaku M, Kondoh T, Yamauchi M. 1,25(OH)<sub>2</sub>D<sub>3</sub> regulates collagen quality in an osteoblastic cell culture system. *Biochem Biophys Res Commun.* 2008 Dec 12;377(2):674-8. doi: 10.1016/j.bbrc.2008.10.036. Epub 2008 Oct 18.
- Nimitphong H, Holick MF, Fried SK, Lee MJ (2012). 25-hydroxyvitamin D<sub>3</sub> and 1,25-dihydroxyvitamin D<sub>3</sub> promote the differentiation of human subcutaneous preadipocytes. *PLoS One.* 2012;7(12):e52171. doi: 10.1371/journal.pone.0052171. Epub 2012 Dec 18.
- Nobili V, Giorgio V, Liccardo D, Bedogni G, Morino G, Alisi A, Cianfarani S. Vitamin D levels and liver histological alterations in children with nonalcoholic fatty liver disease. *Eur J Endocrinol.* 2014 Mar 8;170(4):547-53. doi: 10.1530/EJE-13-0609. Print 2014 Apr.
- Noyola-Martínez N, Díaz L, Zaga-Clavellina V, Avila E, Halhali A, Larrea F, Barrera D (2014). Regulation of CYP27B1 and CYP24A1 gene expression by recombinant pro-inflammatory cytokines in cultured human trophoblasts. *J Steroid Biochem Mol Biol.* 2014 Oct;144 Pt A:106-9. doi: 10.1016/j.jsbmb.2013.12.007. Epub 2013 Dec 17.
- Omdahl JL, Bobrovnikova EV, Annalora A, Chen P, Serda R (2003). Expression, structure-function, and molecular modeling of vitamin D P450s. *J Cell Biochem.* 2003 Feb 1;88(2):356-62.
- Oren Y, Shapira Y, Agmon-Levin N, Kivity S, Zafrir Y, Altman A, Lerner A, Shoenfeld Y (2010). Vitamin D insufficiency in a sunny environment: a demographic and seasonal analysis. *Isr Med Assoc J.* 2010 Dec;12(12):751-6.
- Pajunen P, Torppa J, Huohvanainen M, Salomaa V, Vartiainen V. Sepelvaltimotautikuolleisuuden itälänsiero kaventunut 30 vuoden seuranta-aikana miehillä. *Suomen Lääkärilehti (Finnish Medical journal in Finnish)*51-52/2004 Vsk 59:5013-16.)
- Plotnikoff GA, Quigley JM. Prevalence of severe hypovitaminosis D in patients with persistent,

- nonspecific musculoskeletal pain. *Mayo Clin Proc.* 2003;78(12):1463-70.
- Poopedi MA, Norris SA, Pettifor JM. Factors influencing the vitamin D status of 10-year-old urban South African children. *Public Health Nutr.* 2011; **14**(2): 334-9.
- Rabon HW Jr, Roland DA Sr, Bryant MM, Smith RC, Barnes DG, Laurent SM (1995). Absorption of silicon and aluminum by hens fed sodium zeolite A with various levels of dietary cholecalciferol. *Poult Sci* 1995 Feb;74(2):352-9
- Rude RK, Gruber HE, Norton HJ, Wei LY, Frausto A, Kilburn J. Reduction of dietary magnesium by only 50% in the rat disrupts bone and mineral metabolism. *Osteoporos Int.* 2006 Jul;17(7):1022-32. Epub 2006 Apr 7.
- Schnatz PF, Manson JE (2014). Vitamin D and cardiovascular disease: an appraisal of the evidence. *Clin Chem.* 2014 Apr;60(4):600-9. doi: 10.1373/clinchem.2013.211037. Epub 2013 Nov 5.
- Sharifi N, Amani R, Hajiani E, Cheraghian B (2014). Does vitamin D improve liver enzymes, oxidative stress, and inflammatory biomarkers in adults with non-alcoholic fatty liver disease? A randomized clinical trial. *Endocrine.* 2014 Sep;47(1):70-80. doi: 10.1007/s12020-014-0336-5. Epub 2014 Jun 27.
- Schwarz K, Ricci BA, Punsar S, Karvonen MJ. 1977. Inverse relation of silicon in drinking water and atherosclerosis in Finland. *Lancet.* 1977 Mar 5;1(8010):538-9.
- Statistics Finland 1978 people: Tabl.13, p. 19)  
Statistics Finland 1960 horses: (Tabl. 82, p 88),  
Statistics Finland 1978 horses: (Tabl. 71)
- Valkonen T, Martikainen P. Development of mortality from ischaemic heart disease in subgroups of the population in Finland. *Sosiaalilääketieteellinen Aikakauslehti (J Soc Med)* 1990;27:273-88. (urban, countryside on page 278, Fig5). (Permission: Valkonen T June 6, 2014. Agreement: Salminen M (Editor-in-Chief). *J Soc Med* Aug 14, 2014).
- Velyky MM, Apukhovs'ka LI, Vasylevs'ka VM, Lotots'ka OIu, Besusiak AI, Khomenko AV. (2010). [Features of cholecalciferol hydroxylation in the liver of rats in conditions of D-hypervitaminosis and activity of alpha-tocopherol]. [Article in Ukrainian]. *Ukr Biokhim Zh.* 2010 Mar-Apr;82(2):67-74.
- Wietrzyk J (2007). [The influence of isoflavonoids on the antitumor activity of vitamin D3]. [Article in Polish]. *Postepy Hig Med Dosw (Online).* 2007;61:253-60.