

REVIEW OF POST- HARVEST DETERIORATION OF RUBBER SEEDS

CLARA LEYIBO IGELEKE. AND VICTOR IROGUE OMORUSI

ABSTRACT

Rubber seeds have potential use as a source of industrial oil as well as a protein supplement for use in livestock diets. This has necessitated large-scale collection and storage of rubber seeds for uses other than propagation. Like other traditional Nigerian grains and oilseeds, rubber seeds are prone to post-harvest deterioration. The main causes of deterioration of rubber seeds are excessive moisture, and pests, such as fungi, insects and rodents. To reduce losses, it is imperative that rubber seeds are dried to a moisture level of 7% or less, before storage. Only whole (unshelled) seeds should be considered for long-term bulk storage. Dried, insecticide treated rubber seeds kept in matter polypropylene bags under strict storage hygiene can be effectively stored for at least one year.

Keyword: Rubber seeds, post-harvest, Deterioration, Preservation and Storage.

INTRODUCTION

Annual production of seeds of para rubber (*Hevea brasiliensis* Muell Arg.) Nigeria is estimated at 43,000 tones (RRIN, 1985a). This quantity of seeds is largely wasted, as only a fraction is utilized in growing rootstocks for budding. However, studies have shown the potential of rubber seeds as a source of seed oil for industrial use, such as in the manufacture of alkyd resins for the paint industry (Aigbodion, 1991; Nair *et al.*, 1981) and in the formulation of printing ink, glazing putty, dermal fat-liquor, liquid soap and hair shampoo (RRIN, 1989b). The rubber seed oil, especially when refined, has been shown to have promise as a linoleic acid-rich vegetable oil (Gnathi *et al.*, 1990; Ukhum and Uwatse, 1988). The cake or meal derived from rubber seed after oil extraction is also potentially useable for livestock feeds and diet (RRIN, 1985b). So also is the undefeated rubber seed (Achibewhu, 1986). These findings have necessitated large-scale collection and storage of rubber seeds to ensure regular supply of raw material for industries that may have need for rubber seeds, since rubber seed production is seasonal occurring only once a year in Nigeria (July to October).

For the purpose of this paper, the term 'post-harvest' covers the period between the forcible discharge of rubber seeds from the rubber pods, and the crushing of seeds for industrial use. Much information is available on post-harvest deterioration of traditional Nigerian grains and oilseeds such as maize, groundnut, palm produce, cocoa and kolanuts (Eggin, 1961; Kuku and Adeniji, 1977; Oludemokun and McDonald, 1979). Similar information on rubber seed is lacking. This paper reports on causes of post-harvest deterioration of rubber seeds as well as a method of reducing losses from deterioration. Since much of deterioration occurs during storage, strategies for effective bulk storage of rubber seeds as adopted at the rubber Research Institution of Nigerian (RRIN), are highlighted.

Causes of post – harvest deterioration of rubber seeds

Post-harvest operations of rubber seeds include seed collection, transportation, drying, and storage. Improper methods of such operations, as well as inability to effectively control chemical and biological factors during the post-harvest period could result in deterioration in quality of rubber seeds. In Nigeria above operations are usually by picking the seeds indiscriminately irrespective of clonal types from the plantation floor by vendors. Seeds are

packed in jute sacks and then transported in trucks or in wheel barrows and sold to the rubber seed factories. Seeds are heat-dried in machines, thereafter are re-bagged in fresh jute bags and stored in warehouses for subsequently uses.

Chemical/Biochemical Factors:

Deterioration of seed quality after harvest may occur as a result of chemical reactions brought about by the activity of enzymes naturally present in seeds. Such dexterous changes in rubber seeds which have been observed after harvest include the increase in free fatty acids (Anon, 1978) and production of hydrocyanic glucoside, linamarin, which the seeds contain (Uzu *et al.* 1986; Webster and Baulkwill, 1989). The level of vitamin C (Which is a known antioxidant) may also decrease in stored rubber seeds (Okoye, 1986). In Nigeria, changes leading to deterioration are due to environmental factors such as wet conditions, and if seeds are not prepared adequately for storage, very soon become liable to deterioration.

Role of Moisture:

Rubber seeds are known to have a high moisture content, especially at time of harvest. In Nigeria, moisture content of rubber seeds about one month after seed fall has been reported to be as high as 29.7% at Iyanomo (Otoide and Begho, 1986), and 38% at Itaogholu (Okhuoya and Ige, 1986). In the tropics, 70% is regarded as the minimum relative humidity for the growth of most moulds. The moisture content of grain which is in equilibrium with surrounding air at the grain (Oyeniran, 1980; Vanek and Hoberg, 1992). This varies from grain to grain and has been determined for various Nigerian stored grains (Agboola, 1992; Oyeniran, 1980;). Igeleke (1990) determined the moisture content for deterioration – free storage of rubber seed to be 7% and below. However, poor storage facilities susceptible to moisture exposure abound in Nigeria and account greatly for post harvest deterioration of seeds generally.

Biological Factors:

Pests:

Pests of rubber seeds in Nigeria have been identify as fungi, insects and rodents.

Fungi:

Fungi species associated with pest-harvest deterioration of rubber seeds in Nigeria have been reported (Igeleke and Ekpebor, 1986; Okhuoya and Ige, 1986). They include. *Aspergillus candidus*, *A. tamaril*, *Penicillium sp* and *Rhizopus stolonifer*. Others which have been observed as early colonizers by Igeleke and Ekpebor (1986), being field fungi originating from pre-harvest infection include *Dreacheria (Heininthosporium) sp.*, *Collectotrichum sp.* And *Fusarium*, *Penicillium* and *Rhizopus* have also been reported on rubber seeds of central African origin (Berjak, 1989).

Types of deterioration caused by fungi on rubber seeds:

Discoloration, Rubber seed kernels are buff or flesh-colored when healthy. Fungi growing on kernels impact their various colours to the seeds (Igeleke and Ekpebor, 1986).

Biochemical change.

Moulds are known to increase free fatty acids (FFA) in stored products through the process or lipolysis. Two of the fungi isolated from rubber seeds by Igeleke and Ekpebor (1986) namely, *Aspergillum flavus* and *A. niger* are known to have polytypic abilities (Coursey *et al.*, 1963)

Rotting and caking seed meal (Heating, and production of mycotoxins). Heating occurs with improperly dried rubber seeds stored in heaps or in deep layers. It is an evidence of spoilage. Production of mycotoxins, Okoye (1986) reported amounts of the *Mycotoxin aflatoxin B₂* G₁ and G₂ in stored rubber seeds.

Insects:

Attack by insects is a major problem of stored rubber seeds. *Tribolium* as have been recorded on fresh rubber seeds (Otiode and Begho, 1986) and also on dried stored seeds (Okoye, 1986). Species of *Tribolium* have been associated with various stored seeds in Nigeria (Agboola, 1992) and other parts of Africa (Mlambo *et al.*, 1992). These insects cause appreciable damage to stored rubber seeds by boring holes in the kernels, after gaining entrance through the micropyle in the hard testa. In serious infestations, the kernels are reduced to powder.

Rodents:

Rodents damage storage bags thus scattering seeds in storage. They also eat up some appreciable quantity of rubber seed kernels after cracking open whole seeds. Rodent are know to transmit dangerous diseases to man. These diseases include plague, typhus, trichinosis, rabies and Lassa fever (Hoppe. 1980, Zehrer, 1980). The control of rodents is therefore imperative.

Pre-storage handling of rubber seeds

Reducing deterioration during early post-harvest operations

Much of rubber seeds deterioration occurs in storage. However, maintenance of proper handling methods in early post harvest operation have been observed at the Rubber Research Institute of Nigeria (RRIN) to greatly reduce deterioration. Some of these methods are considered.

Rubber Seed Collection

Shortly before seed fall, rubber plantations should be clean weeded, and old seeds from previous seed seasons removed from the plantations. This allows for easy collection of fresh, whole seeds (RRIN, 1989a). By collecting seeds once or twice a week, they do not remain on the ground for too long where they are exposes to moisture as well as insect and fungal attack (Aniamaka and Uraih, 1990). Onokpise (1980) observed that rubber seeds harvested in the first and second week of August had the highest weights indicating the presence of greater food reserves. Early August may therefore be the best time to collect seeds for storage.

Transportation

Care should be taken in transporting seeds from the site of collection, in order to ensure that seeds do not become crushed or cracked as this allows fungi to attack easily. Seeds should preferably be transported in bags that allow for aeration, as the seeds are still respiring. If aeration is poor, there could be a build-up of moisture and heat, encouraging growth of microorganisms.

Preparation for storage

As soon seeds have been transported from the field, they should be removed from containers, sorted, spread out and exposed to air, if artificial drying cannot be done immediately. This allows for natural drying and an appreciable drop in moisture content. Otoide and Begho (1986) observed that fresh wholes seeds with initial moisture content of

29.7%, air-dried to a moisture content of 8.4% after a months storage on the floor. If the seeds are several layers thick, they should be occasionally turned over with shovels to expose underlying layers to air.

Measures to control deterioration during storage

Drying

Apart from reducing moisture, drying or heat treatment also halts biochemical processes, which leads to degradation of fats and oils (Ihekoronye and Ngoddy, 1985). Begho, *et al.* (unpublished) observed during a 4-month storage study that there is very little increase in FFA of oil extracted from properly dried rubber seeds. Work in Malaysia (Anon, 1975) and Sri Lanka (Nadarajah *et al.*, 1973) have also shown that increase in FFA value of stored rubber seeds can be effectively checked when seeds are heat-treated before storage. Increase in FFA during storage is an important indication of the level of spoilage in oilseeds (Landers and Rathmann, 1981; Okoye, (1986). Drying also greatly reduces the cyanide content of rubber seeds to innocuous levels (Uzu *et al.*, 1986) suggested that merely toasting the kernels in the oven may be sufficient to render them suitable for use as animal feed.

Air-drying is fairly effective only for short-term storage of relatively small quantities of rubber seeds, especially if the seeds are visually free of moulds and insects at the time of collection. For effective long-term storage, it is advisable that rubber seeds are dried to a stage of moisture level of 7% or less. At RRIN, this is achieved by artificial drying, (for 9-12 hrs.) Using heated air (60-70⁰c) in a batch drier (Figure 1). Drying temperature above 70⁰c may affect some properties of rubber seeds and the resultant oil. For instance, seeds become brittle and the colour of the seed and oil extracted from it darken (Igeleke, 1990). Length storage also affects colour of extracted oil (Aigbodion, 1994). Batch drier machines are usually employed and the traditional measure of drying the seeds in storage room heated prior to spreading of the seeds on mats on the floor of the room. However, batch drier is a better measure of drying rubber seeds.

Method of storage

After seeds have been dried, they should be allowed to cool before storage. The preferred containers of storage, particularly in bulk storage, are matted polypropylene bags (Otoide and Begho, 1986). These bags are readily available in the market. The disadvantage of their use is that stored seeds are prone to insect attack. It has been observed that whole seeds (unshelled) store better than kernels (Igeleke and Ekpebor, 1986; Otoide and Begho, 1986). Consequently, only whole clean seeds should be considered for long-term storage.

Storage Hygiene

Dried rubber seeds should be stored in a dry, cool environment, to prevent growth of moulds. This can be done in a properly ventilated warehouse, where there are no leakages or floods. At RRIN, rubber seeds in 25kg bags are stored in a warehouse (35 – 40 tonnes capacity). The bags are raised from the floor, on wooden shelves spaced 65cm apart. Bagged seeds are placed on each shelf either vertically, in a single layer, or stacked horizontally, one on top of the other in two layers only (Figure 2). This arrangement allows for enough space for aeration between loaded shelves. Temperatures in the warehouse are kept relatively low (25 – 30⁰C) by a projecting roof and by ventilation points. Preventive measures against insect infestation include covering ventilation points with insect-proof netting. The warehouse should be cleaned out thoroughly and disinfested before seeds are received for storage. Rats can be excluded from the warehouse by sealing up all openings and cracks, and by ensuring that doors and windows fit well. Piles of rubbish, including old

rubber seeds, should be removed from the vicinity of the store, as these attract rodents. Regular cleaning and inspection of the warehouse reduce episodes and levels of pest attack. At the end of the storage season, the warehouse should be thoroughly cleaned out before fresh seeds are again stored. Observance of these measures of storage hygiene, as well as proper pest control, have resulted in successful bulk storage of rubber seeds for at least one year at RRIN

Use of Chemical during storage

Insects could be a serious problem in storage. At RRIN, insects have been successfully controlled in bulk storage by the use of the contact insecticide Actellic (i.e. 2% primiphos methyl) applied either as dust to the seeds (dosage: 50g/100kg seeds) or as liquid spray to storage shelves and bags containing infested seeds (application: 200ml per 18 litres of water, per 100m²). The fumigant, hydrogen phosphide, in the form of phostoxin or Gas-Ex-T tablets (i.e. aluminum phosphide) has also been effectively used to control insects (application: one tablet, in an envelope, placed in each 25kg bag of dried rubber seeds). Both pirimiphos methyl and Hydrogen phosphide are among the insecticides recommended for food storage (Agboola, 1992). Laboratory tests at RRIN indicate that treatment with these two chemicals has little or no effect on such properties as acid value, peroxide value, saponification value, iodine value and specific gravity of oil extracted from chemical treated rubber seeds (Begho *et al.*, unpublished data). If ventilation points in the storage, the need for repeated application may not be necessary. Rodents can be kept in check by use of available rodenticides in the market. However, care should be taken to handle rodenticides as most of them are extremely toxic to humans. Rodent control has been achieved by the use of blocks of klerate (i.e. *brodifacoum*). And also by Gas-Ex-T tablets. These tablet are recommended by the manufacturers for use against rats and other rodents as well as insects. The tablet are grand and mixed with food items such as dry fish, and used as bait.

Seed handling practices among farmers in Nigeria

In Nigeria, farmers collect seeds which are traditionally stored in earthenware pots containing sharp sand where they are embedded. The limitation of this practice is that the pots used are small in size to accommodate large quantities of seeds. The natural method of drying the seeds which involves drying the seeds in the sun and free air flow, may take long hours to dry and may be exposed to rain. Generally, seeds collected by farmers may not attain the desirable moisture content before storage and are often prone to insect, mite, fungal and bacterial attack and consequence of it leads to seed deterioration. Improved technology of drying seeds usually involve batch drier machines for large consignment of seeds or any mechanical device using temperature and moving air through the seeds (Okoye, 1986).

Seeds under the traditional method of drying, have short shelf life. For prolonged shelf life, seeds should be collected immediately after seed fall to prevent uptake of moisture, insect or mould activities. Seeds should be dried to safe moisture content of 7% and below capable of reducing the activities of most agents of deterioration. Seeds such as rubber seeds should be bagged in jute sacks before transportation to reduce spillage losses and mechanical damage such as splitting and bruising of seeds. These practices preserve and sustain longer shelf life of seeds for appreciable utilization of seeds.

CONCLUSION

Rubber seeds, like all other oil-rich seeds are prone to deterioration, much of which occurs in storage. The quality of oil extracted from stored rubber seeds is important in the oil-based manufacturing industries (Aigbodion, 1991). Maintaining proper handling methods

during post-harvest operations; in particular drying to a safe moisture level, as well as observance of strict storage hygiene, can greatly reduce post-harvest deterioration of rubber seeds meant for industrial use.

REFERENCES

- Achinewhu, S. C. 1986. Unconventional sources of food: Chemical composition of rubber seed (*Hevea brasiliensis*). *Food Chemical* 21: 17 – 25.
- Achinewhu, S.C. and Akpapunum, M. A. 1985. Physical and chemical characteristics of refined vegetable oils from rubber seed (*Hevea brasiliensis*) and bread – fruit (*Artocarpus untilis*). *Quality Plant Foods. Human Nutrition* 35 : 103 -107
- Agboola, S.D. 1992. Technology for small scale storage of grains in Nigeria. pp 22 – 37 In: Food storage, processing and utilization. *CODIR occasional paper No. 1. 57p.*
- Aigbodion, A. I. 1991. Preliminary studies on the preparation of rubber seed oil alkyds. *Indian Journal of Naural Rubber Research* 4 (2): 141 – 117.
- Aigbodion A. I. 1994. Effect of storage of seeds on quality of rubber seed oil. *Indian Journal of Natural Rubber Research* 7 (2) :141 – 143.
- Aniamaka E.E. and Uraih, O.B.C. 1990. Collection and preparation of rubber (*Hevea brasiliensis*) seeds for seeding production. pp 75 – 79 In: proceeding of a national workshop on Fruit / tree production NAERLS, Zaria. 160p.
- Anon 1975. Rubber Research Institute of *Malaysia Annual Report* 160 pp.
- Anon 1978. Rubber Research Institute of *Malaysia Annual Report*. 206 pp.
- Berjak, P. 1989. Storage behaviour of Rubber seeds (*Hevea brasiliensis*). *Journal of Natural Rubber Research* 4 (3): 195 – 203.
- Coursey, D. G., Simons E. A. and Sheridan, A. 1963. Studies on the quality of Nigerian palm Kernels *Journal of West Africa Science Association* 8 (1): 18 – 28.
- Eggin, H. O. W. 1961. Notes on fungal damage to palm produce in Nigerian *West Africa Stored Products Research Unit Technical Rep. 7: 53 – 59.*
- Gandhi, V. M., Cherian, K.M. and Mulky, M. J. 1990. Nutritinonal and Toxicological evaluation of rubber seed oil. *JACOBS* 67 (11) 883 – 886.
- Hoppe, A. H. 1980. The control of rodent pests in grain storage. Appendix pp 1 – 33. In: Post Harvest problems. Developmentation of a OAU/GTZ Seminar. GTZ, Germany, 258 pp.
- Igeleke, C.I. 1990. Effect of temperature on the storability of Rubber (*Hevea bresilienesis*) seeds. *Nigeria Journal of Horticultural Science* 1 (1): 27-30.
- Igeleke, C. L. and Ekpebor, S.M.C. 1986, Fungal species associated with deterioration of rubber seeds during storage pp 135 – 145. In: E. E. Enabor (ed). Industrial Utilization

of natural rubber (*Hevea brasiliensis*) seed, latex and wood. Rubber Research Institute of Nigeria. Benin City.

Thekoronye, A.I. and Ngoddy, P.C. 1985. Integrated food science and technology for the tropics. Macmillan publishers. 386 pp.

Kuku, F. O. and Adniji, M.O. 1976. The effect of moulds on the quality of Nigeria palm kernels, *International Biodeterioration Bulletin* 12(2): 37 – 41.

Landers, R. E. and Rathman, D.M 1981. Vegetable oils: Effects of processing, storage and use on nutritional values. *JAOCs* 58(3): 255 – 258.

Mlambo, S.S., Sithole, S.Z. and Tanyongana, R. (1992) Storage facilities in Zimbabwe and losses due to insects. pp 109 – 116. In: S.B. Mathur and Johs Jorgensen (eds). Seed pathology. Technical Centre for Agriculture and Rural Cooperation (CTA), Netherlands.

Moore, R. (1969). *Food preservation. Revised ordinary Series No.3*. Unilever, London. 24 pp.

Nadarajah M., Abeysinghe, A., Dayaratne, W.C. and Tharmalingam, R. 1973. The potentialities of rubber seed collection and its utilization in Sri Lanka. *Rubber Research of Sri Lanka Bulletin* 8 (1): 9 – 21.

Nair, N.M., Unnikrishan, K.C., Unnikrishan, M. and Nair, C.S.B. 1981. Utilization of rubber seed oil and karinnota oil for the preparation of air-drying oil-modified alkyd resins. *Paintindia* 31 (5): 5 – 9.

Nash, M.J. 1978. Crop conservation and storage. Pergamon Press. 393 pp.

Okoye, W.I. 1986. Strategy for the effective storage of rubber seeds, oil and cake. Pp 123 – 129. In: E.E. Enabor (ed). Industrial utilization of natural rubber (*Hevea brasiliensis*) seed, latex and wood. Rubber Research Institute of Nigeria. Benin City.

Okhuoya, J.A. and Ige, O. 1986. Fungi associated with biodeterioration of rubber seeds. pp 146 – 154. In: E.E. Enabor (ed). Industrial utilization of natural rubber (*Hevea brasiliensis*) seed, latex and wood. Rubber Research Institute of Nigeria. Benin City.

Oludemokun, A.A. and McDonald, D. 1979. A survey of fungi associated with diseased trees and stored nuts of kola, *Cola nitida* (Vent). Schott and Endlicher, in the Northern State of Nigeria. *Nigeria Agricultural Journal* 16: 77 – 81.

Olunloyo, O.A. 1979. Fungi associated with deterioration of stored kola nuts. *Nigeria Journal Agricultural Science* 1 (1) 51 – 59.

Onokpise, O.U. 1980. Germination success and seedling performance in relation to time of harvest of seeds of *Hevea brasiliensis* (Muell. Arg.) *Nigeria Journal of Science* 14 (1 and 2): 24 – 25.

Otoide, V.O. and Begho, E.R. 1986. An assessment of some methods of storage of rubber seeds and undefeated seed cake. pp 130 – 134. In: E.E. Enabor (ed) Industrial

utilization of natural rubber (*Hevea brasiliensis*) seed, latex and wood. Rubber Research Institute of Nigeria. Benin City.

Oyeniran, J.O. 1977. Fungal deterioration of maize during storage in Nigeria. *Nigeria Journal of Plant Protection* 3: 102 – 105.

Oyeniran, J. O. 1980. The role of fungi in the deterioration of tropical stored products. Nigeria Store Products Research Institute occasional paper series No. 2. 25 pp.

Oyeniran, J. O. and Adeniji M. O. 1975. Mould deterioration of cocoa during processing and storage in Nigeria *Nigeria Journal of Plant Protection* 1 (1): 48 – 51.

Rubber Research Institute of Nigeria. 1985a. Highlights of research achievements on rubber seed oil. Federal Ministry of Education, Science and Technology. 18 pp.

Rubber Research Institute of Nigeria. 1985b. Highlights of research on rubber seed meal as a constituent of livestock diets. Federal Ministry of Education, Science and Technology. 14 pp.

Rubber Research Institute of Nigeria. 1989a. Processing of rubber seeds into oil. Advisory Circular No.7. Rubber Research Institute of Nigeria. Benin City.

Rubber Research Institute of Nigeria. 1989b. Industrial utilization of natural rubber seed oil. Advisory Circular No. 8. Rubber Research Institute of Nigeria. Benin City.

Sheridan, A. 1961. Preliminary notes on the micro-organisms associated with the deterioration of groundnuts and palm kernels during storage in Nigeria *West Africa Stored Products Research Unit Technology Rep.* 6: 52 – 55.

Spannuth, H.T. 1949. Stabilization and antioxidants. *JAACS* 26 (10): 618 – 622.

Ukhun, M.E. and Uwatse, G.M. 1988. Nutritional evaluation of selected Nigerian rubber seed products – a chemical approach, *Plant Foods Human Nutrition* 38: 309 – 318.

Uzu, F.O. Ihenyen, G.A and Imoibe, S.O. 1986. Processing, analysis and utilization of rubber (*Hevea brasiliensis*) seed oil and cake. pp 19 – 26. In: E.E Enabor (ed). Industrial utilization of natural rubber (*Hevea brasiliensis*) seed, latex and wood. Rubber Research Institute of Nigeria. Benin City.

Vanek, K. and Hoberg, K. 1992. Simple means to improve seed storage. pp 141 – 147. In: S. B Mathur and Johs Jorgensen (eds.) Seed Pathology. Technical Centre for Agriculture and Rural Cooperatioin (CTA), Netherlands.

Webster, C.C. and Baulkwill, W.J. (Eds.). 1989. Rubber Longman Scientific and Technical 614 pp.

Zehrer, W. 1980. Preventive and hygienic measures in crop storage. pp 31-70. In: Post Harvest Problems Documentation of OAU/GTZ Seminar GTZ, Germany.




Fig. 1: Dried rubber seeds being removed from the batch drier into a mounted polypropylene bag for storage.

