

THE EFFECT OF *XYLOPIA AETHIOPICA* (UDA) ON INTRAOCULAR PRESSURE

BY

*UZODIKE, E. B. AND ONUOHA, I. N.

DEPARTMENT OF OPTOMETRY

ABIA STATE UNIVERSITY, UTURU, ABIA STATE

Email: ebeluzo@yahoo.com

*Corresponding author

ABSTRACT

Xylopiya aethiopica (Uda) is a popular spice used in the preparation of soups. Fifty volunteers between the ages of 18 to 30 (mean age of 22.58 ± 12.75) years were used for this research. They were given 20ml (1.16g/ml) of *Xylopiya aethiopica* (*X. aethiopica*) extract. The baseline and induced intraocular pressure (IOP) of the volunteers were measured at 30 minutes interval after the ingestion of *X. aethiopica* until the IOP approximated the baseline value. The result showed an initial 1.26% increase in IOP at 30mins post ingestion of *X. aethiopica* which started reducing from 60mins with a maximum reduction of 2.60% at 90mins. The test for significance using Z-test showed the cumulative effect to be statistically significant at 95% confidence interval ($P > 0.05$). The effects were transient as the induced mean IOP approximates the baseline IOP after 90minutes of ingestion.

KEYWORDS: *Xylopiya aethiopica*, Intraocular pressure, Ethno-medicine, Saponin, Anonecaine.

Received 11/12/09

Accepted 15/02/10

INTRODUCTION

Every food substance consumed by humans has either a therapeutic, nutritional or toxic effect on the body. These food substances when got in their crude form can be of immense help in the curing of some ailments. Plants as well have been used for therapeutic purposes and their uses are as old as the history of man. In the past decades, pharmacologists and organic chemists have synthesized a large number of interesting chemical substances from medicinal plants, which have been of great help in the practice of Optometry and Medicine; example is Belladonna plant from which atropine is derived.

Xylopiya aethiopica (*X. aethiopica*) commonly known as "African guinea pepper" or "Ethiopian pepper" is wide spread in tropical Africa, Zambia, Mozambique and Angola¹. In Nigeria, it is found all over the lowland rain forest and most fringe forest in the Savanna zones of Nigeria. Negro pepper as it is also known has been used as a pepper substitute in Europe and India².

X. aethiopica is common in ethno-medicine in West Africa. This is due to its preservative effect, the fruit extract has been shown to be active as antimicrobial agent against gram positive and gram negative bacteria, though it has not been shown to be effective against *Escherichia Coli*³. *X. aethiopica* has anti-spirochoetal properties so that it works both as a preventive measure and in

treatment of primary, secondary and tertiary stages of syphilis⁴. *X. aethiopica* has been used for treating rheumatism and arthritis as well as other inflammatory conditions. Numerous research studies have confirmed the spice's anti-inflammatory and antipyretic (fever reducing) properties^{2,3}. Indian researchers reported anti-arthritic and anti-inflammatory actions of one of the compounds of *X. aethiopica* called nimbidin⁵.

The seeds are mainly used by traditional medicine healers and can also serve as an alternative to pepper⁶. Medical uses of the plants are: as a carminative, as cough remedy and as a post partum tonic and lactation aid. Other uses include treatment for stomach ache, bronchitis, biliousness and dysentery⁶. It has also been reported to be used as a flavour in palm wine. The bark of *X. aethiopica* tree when steeped in palm wine is given for attacks of asthma and rheumatism⁷.

It is widely accepted that fruits and vegetables have many healthful properties. There are considerable amount of epidemiological evidences revealing an association between those who have a diet rich in fresh fruits and vegetables and a decreased risk of cardiovascular diseases and certain forms of cancer^{8,9}. The constituent of these fruits and vegetables that contribute to these protective effects are phytochemicals, vitamins and minerals¹⁰. Naturally occurring phytochemicals like flavonoids are potentially

anti-allergic, anti-carcinogenic, antiviral and antioxidant agents¹¹. Phytochemicals as antioxidants play vital roles in human health^{9,12}. *X. aethiopica* has been found to contain some phytochemicals which exhibit a wide range of biological effects as a consequence of their antioxidant properties¹¹.

The chemical components of *X. aethiopica* have been helpful in the avoidance and treatment of cancerous tumors. Researchers in India, Europe and Japan have found that polysaccharides and limonoids found in *X. aethiopica* reduce tumors and cancers¹³⁻¹⁵.

Flavonoids represent the most common and widely distributed of plant phenolics found in *X. aethiopica*. Flavonoids prevents oxidative cell damage, have strong anti-cancer activity and protects against all stages of carcinogenesis^{8,9}. As antioxidants, flavonoids from *X. aethiopica* provide anti-inflammatory action^{16,17}. Anonecaine an alkaloids constituent of *X. aethiopica* is known to have anti-pyretic effect¹⁸. Saponins another phytochemical constituent of *X. aethiopica* have wide range of biological properties; they are used to recover homeostasis, have anti-inflammatory and anti-cancer actions^{19,20}.

The normal IOP varies between 10mmHg - 21mmHg (Mean 16mmHg)²¹. Although there is no absolute cutoff point, 21mmHg is considered the upper limit of normal, and levels above this are viewed with suspicion. However in some patients, glaucomatous damage occurs with IOP less than 21mmHg, whereas others remain unscathed, at least in a short term with IOP up to 30mmHg²¹.

Fluctuations in IOP occur with time of the day, heartbeat, blood pressure level and respiration²¹. The pattern of diurnal curves of IOP varies in the normal and glaucomatous eye, with a tendency towards increased IOP in the morning and lower IOP in the afternoon and evening. Normal eyes have a smaller diurnal fluctuation (4mmHg) than glaucomatous eyes in which the fluctuation may be 10mmHg or more²¹.

Since *X. aethiopica* is widely used in traditional medical practice, as spice in foods and as a preservative in drugs, there may be some effects on the general body physiology, in which the ocular functions may equally be affected. The eye care practitioners should therefore have a working knowledge of the effect of *X. aethiopica* on the ocular system.

RESEARCH METHODOLOGY

This study was carried out in the Optometry clinic of Abia State University, Uturu because of convenience of the research volunteers and the availability of research instruments and materials needed. The research was designed as a prospective and clinically based one. The volunteers used were students of Abia State University within the age limit of 18-30 (mean age of 22.58±12.75) years. These volunteers passed the inclusion criteria of being emmetropic, free of any ocular or debilitating systemic disease, not under any medication and not smokers or alcoholics.

Fresh *X. aethiopica* fruits were purchased from Eke Okigwe market in Imo State, Nigeria and certified good for human consumption by a renowned botanist in the Department of Botany of the University. *X. aethiopica* extract was obtained through the batch method²², using water as the extracting solvent. The carpels of *X. aethiopica* were removed from their strands, washed and dried in the sun after which, 50g was measured out and ground with a grinding machine. The paste was soaked in 120ml of distilled water for 24 hours, after which it was filtered out using a white cotton gauge. The quantity of fluid extracted was measured. The solid filtrate was dried and weighed; the value was subtracted from the 50g original value. The balance gave the quantity of *X. aethiopica* in the solution. The value obtained after extraction was approximately 43g showing a balance of 7g in solution of 120ml. The concentration of *X. aethiopica* extract was 58.30mg/ml. The extract was sterilized using an autoclave.

Baseline IOP was measured before 20ml of *X. aethiopica* stock extract (58.30mg/ml concentration) was given to each subject to drink and the IOP was measured 30minutes after intake and subsequently at intervals of 30minutes until IOP returned to baseline. There are no chemokinetic studies available to the best of the researchers' knowledge but the rigor effects of ingesting the extract within few minutes post ingestion showed an instant absorption. This informed the choice of 30mins for monitoring of induced effects. All the IOP measurements were taken between 10.00am and 1.00pm to take into consideration the diurnal nature of IOP. Each measurement was repeated 3times and the average used. Data collected was tabulated and analyzed using Z-statistics.

RESULTS

The result in table 1 showed a sharp increase in IOP within 30minutes post ingestion of *X. aethiopica* extract (from 21.81mmHg to 22.09mmHg). After which, there was a reduction in IOP from 22.09mmHg to 21.83mmHg at 60minutes post ingestion and further reduction below baseline IOP at 90minutes (from 21.83mmHg to 21.24mmHg). The mean induced change in IOP was found to be 0.28mmHg, 0.02mmHg and -0.57mmHg with percentage change of 1.26%, 0.09% and 2.60% at 30mins, 60mins and 90mins respectively post ingestion of *X. aethiopica*. Z-statistics showed the cumulative effect of *X. aethiopica* to be statistically significant ($P > 0.05$; $Z_{cal} = \pm 0.83$). The effect of Uda extract was found to be transient and not sustained.

It is important to point out here, though not within the scope of this study, that the initial and spontaneous reaction noticed upon ingestion of *X. aethiopica* was reflex tearing.

which are antioxidants thus playing a vital role in therapies.

Also it has been shown from recent studies that anonecaine one of the constituents of *Xylopi aethiopica* has morphine like activity and that morphine acts through opion receptor in the eye, these receptors alpha (α) and kappa (K) stimulates the parasympathetic system. Opiod receptors have different actions in the autonomic nervous system that is; it has both an agonist and antagonist effect²⁶. It stimulates the oculomotor nerve center, which could result in the increase in accommodation thereby constricting the ciliary process and opening the canal of Schlemn²⁷. Moreover, the phytochemical constituents of *X. aethiopica* like limonoids, nimbidin and tannins have been found to have central nervous system depressing effect, hence a parasympathetic effect that aids in IOP reduction²⁸⁻³². According to Igwe et al³³, the aqueous extract of *X. aethiopica* was neither miotic nor mydriatic.

Table 1: Mean change in IOP after intake of 20ml of 58.30mg/ml of *Xylopi aethiopica* Extract (Baseline IOP was 21.81mmHg).

Time interval (Minutes)	Mean induced IOP (mmHg)	Mean change in IOP (mmHg)	% change in IOP (%)
30	22.09	0.28	1.26
60	21.83	0.02	0.09
90	21.24	-0.57	2.60

DISCUSSION

Drinking of 20ml of 58.30mg/ml of *X. aethiopica* stock extract was found to induce a 1.26% increase in IOP after 30mins, followed by a 0.09% reduction at 60mins and 2.60% further reduction at 90mins. The increase in IOP could be as a result of saponin which Lawal et al²³ found to be abundant in *X. aethiopica*. Saponin has been found to increase the norepinephrine and dopamine content of central nervous system (CNS) and also to have an anti-fatigue effect and excitatory effect on the CNS²⁴⁻²⁶. This effect of saponin is sympathetic in origin and production of aqueous humor is through the stimulation of the beta receptors of the sympathetic autonomic nervous system.

The reduction effect on IOP of *X. aethiopica* might be as a result of its constituents like flavonoids and anonecaine. According to Agoha⁶, *X. aethiopica* is known to contain flavonoids,

The immediate and spontaneous tearing of the volunteers upon ingestion of the stock extract showed a reflex tearing effect which is parasympathetic in origin, confirming the parasympathetic characteristics of *X. aethiopica* constituents³⁴.

Although the effect of *Xylopi aethiopica* on IOP was transient, it could affect optometric findings and analysis, it could also affect tests like tear film break up time and slit lamp biomicroscopy because *Xylopi aethiopica* intake would cause stimulation of local axon reflex resulting in tearing. There is need to probe the social history of patients with regards to what was ingested before optometric examination.

Researches are recommended for the chemokinetic analysis of *X. aethiopica* to help in the understanding of both the systemic and ocular effects.

REFERENCES

- Puri S.G. and Talata O, (1978): Paper presented in A Symposium on Recent Advances in the Development, Production and Utilization of Medicinal and Aromatic plants in India: A survey of some plants used in Native Medicine of West Africa of interest to India; p. 35.
- Sofowara, E. A. (1978): Medical Plants and traditional medicine in Africa. The pitman Press, Bath, Avon, pp 314-8.
- Iwu, M. (1993): Handbook of Africa Medical plant. CRC Press, pp 435.
- Mitra, C. R. and Misra, P. S. (1967): Amino acids of processed Seed meal proteins. J. Agric. Food Chem, 15 (4):697700.
- Pillai, N. R. and Santhakumari, G. (1981): Medicinal, Culinary and Aromatic Plants in the North East. *Planta Medica*, 43:59-63.
- Agoha, R.C. (1974): Medicinal Plant of Nigeria; Offset- Happen, Netherlands, pp. 86.
- Smith, G. M., Clegg, C. K. and Grivetti, L. (1996): Mineral values of Selected Plant Foods Common to Southern Burkina Faso and to Niamey; Niger. *Int. J. Food Sci. Nutri*, 47:41-53.
- Salah, N., Miller, N. J., Paganga, G., Tijburg, L., Bolwell, G. P. and Riceevans, C. (1995): Polyphenolic Flavanols as Scavengers of Aqueous Phase Radicals and as Chain-Breaking Antioxidants. *Arch. Biochem. Biophysics*, 322(2): 339-46.
- Del-Rio, A., Obdulio, B. G., Castillo, J. Main, F.R. and Ortuno, A. (1997): Uses and properties of Citrus flavonoids. *J. Agric food Chem*. 45: 4505-15.
- Okwu D.E. (2003): The potential of *ocimum gratissimum*, *pergturia extersa* and *tetrapleura tetraptera* as spice and flavoring agent. *Nig. Agric. J*, 34: 45 148.
- Close, D. C. and McArthur, C. (2002): Rethinking the role of many plant phenolics protection from photo damage. *J. Plant Med*, 99:166-72.
- Eleangovan, V., Sekar, N., Gorindasamy, S. (1994): Chemo preventive potential of dietary bioflavonoids against 20 methyl cholanthren induce tumor genesis. *Cancer Letters*, 87:107 13.
- Cai, Q. (1997): Dietary flavonoids, quercetin, luteolin and genistein, reduce oxidative DNA damage and lipid peroxidation and quench free radicals. *Cancer Letters*, 119(1): 99-107.
- Hartwell, J. L. (1984): Plants Used Against Cancer: A Survey (Bioactive Plants, Vol 2). Quarterman Publications Inc, pp710.
- Graham, J. G., Quinn, D. S., Fabricant, D. S. and Farnsworth, N. R. (2000): Plants used against cancer an extension of the work of Jonathan Hartwell. *J. Ethnopharmacol*. 73(3): 347-77.
- Okwu, D. E. (2001): Improving the nutritive value of cassava tapioca meal with local spices. *J. Nutraceutica Functional Med. Food* 3:43-51.
- Herthorg, M. G. L.; Feskeens, E. J. M; Hokum, C. H. and Katan, A. (1993): Dietary antioxidant flavonoids and risk of coronary heart disease: J. De Zutphen elderly study. *Lancet*, 342:1007 11.
- Clarence, T. V. (1980): Volatile oils in Remington's Pharmaceutical sciences. 17th Edn. Merck Publishing Company, Easton Pennsylvania, pp. 834-56.
- Sparg, S. G., Light, M. E. and van Staden, J. (2004): Biological activities and distribution of plant saponins. *J. Ethnopharmacol*, 94 (2-3): 219-43.
- Shoji, S. (2001): Chemistry and cancer presenting activities of Ginseng Saponins and some related triterpenoid compounds. *J. Korean Med. Sci*, 16(suppl): S28-37.
- Kanski, J. J. (1994): Clinical Ophthalmology- A Systematic Approach. 3rd Edn. Butterworth-Heinemann, England, 514pp.
- Ijeh, I. I., Omodamiro, O. D. and Nwanna, I. J. (2005): Antimicrobial effects of aqueous extract and ethanolic fractions of two spices, *Ocimum gratissimum* and *Xylopia aethiopica*. *Afri. J. Biotechnol*, 4(9):953-6.
- Lawal, I. O., Igboanugo, A. B. I., Osikarbor, B., Duyilemi, O. P., Adesoga, A. A. and Adeyanju, B. A. (2010): Evaluation of plant-based non-timber forest products (ntfps) as potential

- bioactive drug in South-western Nigeria. J. Clin. Med. Res, 3 (4): 061-6.
24. Kim, Y. C., Lee, J. H., Kim, M. S. and Lee, N. G. (1985): Effect of the saponin fraction of *Panax ginseng* on catecholamines in mouse brain. Arch. Pharmacol Res, 8(1): 45-8.
 25. Tachikawa, E., Kudo, K., Kashimoto, T. and Takahashi, E. (1997): Effects of ginseng saponins on receptor stimulation-responses. Nippon Yakurigaku Zasshi, 110 (Suppl 1): 126-31.
 26. Tachikawa, E., Kudo, K., Harada, K., Kashimoto, T., Miyate, Y., Kakizaki, A. and Takahashi, E (1999): Effects of ginseng saponins on responses induced by various receptor stimuli. Eur. J. Pharmacol, 12; 369(1):23-32.
 27. Katzung, B. G. (2006): Basic and Clinical Pharmacology. 10th Edn. McGraw-Hill Publishing, 1179pp.
 28. Kausik, B., Chattopadhyay, I., Banerjee, R. K. and Bandyopadhyay, U. (2002): Biological activities and medicinal properties of Neem (*Azadirachta indica*). Curr. Sci, 82(11): 1336-45.
 29. Danjuma, N. M., Abdu-Aguye, I., Anuka, J. A., Hussaini, I. M., Zezi, A. U., Yaro, A. H., Maiha, B. B. and Dabo, I. (2009): Central Nervous System Depressant Effect of the Hydroalcoholic Extracts of Leaves, Stem and Root Barks of *Randia Nilotica* Stapf. (Rubiaceae). Eur. J. Scientific Res, .25(3):353-61.
 30. Suárez, A., Ulate, G. and Ciccio, J. (1997): Cardiovascular effects of ethanolic and aqueous extracts of *Pimenta dioica* in Sprague-Dawley rats. J. Ethnopharma, 55(2): 107-11.
 31. Sebastián, P. F., Cristina, W., Leonardo, M. L., Renee, E. G., Graham, A. R. J., Alejandro, C. P. and Mariel, M. (2006): Central nervous system depressant action of flavonoid glycosides. Eur. J. Pharma, 539(2006): 16876.
 32. Pillai, N. R. and Santhakumari, G. (1984): Some pharmacological actions of 'nimbidin' - a bitter principle of *Azadirachta indica* A Juss (neem). Ancient Sci. Life, 4(2): 88-95.
 33. Igwe, S. A., Afonne, J. C. and Ghasi, S. I. (2003): Ocular dynamics of systemic aqueous extracts of *Xylocarpus aethiopicus* (African guinea pepper) seeds on visually active volunteers. J. Ethnopharmacol., 88(2-3): 139-42.
 34. Davson, H. (1980): Physiology of the Eye. 4th Edn. Churchill-Livingstone, Edinburgh, London, pp644.