



Microbes and Infectious Diseases

Journal homepage: <https://mid.journals.ekb.eg/>

Original article

Knowledge, perception and practices toward COVID-19 among care receivers attending a university medical center in Nigeria: A cross-sectional study

Bala Ningi Umar ^{*1}, **Nuhu Mohammed** ², **Kabiru Haliru Ahmad** ¹, **Khadijah Usman** ³, **Rahman Khaleequr** ³

1- Department of Veterinary Microbiology, Faculty of Veterinary Medicine, Ahmadu Bello University, Zaria, Nigeria.

2- Department of Clinical Pharmacology and Therapeutics, Faculty of Basic Clinical Sciences, Ahmadu Bello University, Zaria.

3- Ahmadu Bello University Medical Centre, Zaria-Nigeria.

ARTICLE INFO

Article history:

Received 8 July 2021

Received in revised form 15 August 2021

Accepted 19 August 2021

Keywords:

COVID-19

Coronavirus

Pandemic

SARS-CoV-2

ABSTRACT

Background: Knowledge is one of the contributing factors formulating the perception and attitude of people towards disease outbreaks, which in turn drives compliance to effective prevention and control measures. The current cross-sectional study assessed the knowledge, perception and practices (KPP) towards COVID-19 pandemic among care receivers attending Ahmadu Bello University medical center (ABUMC) in Zaria, Nigeria. **Methods:** Data were collected using self-administered questionnaires which covered demographic characteristics and related KPP questions. Responses were tabulated, frequencies and percentages were computed for descriptive purposes, while independent t-test and ANOVA were used to depict statistical differences between demographic groups. **Results:** Among the 150 participants who answered the questionnaires, 58.7% were female, 62.0% were undergraduate students at various level of studies, 25.5% held a bachelor degree, about half of them (52.7%) never married, and have mean age of 29.90 ± 10.66 years. Majority of the participants, especially women are knowledgeable about COVID-19 with overall mean knowledge score of 4.91 ± 1.68 (70.14%). Similarly, 16-25years age range, and higher educational status were found to possess significantly ($p < 0.05$) higher mean knowledge score. They hold positive perceptions, and have correct practices towards COVID-19 prevention. However, significant proportion of the participants (12.0%) believed that the disease does not exist in Nigeria. **Conclusion:** This study has demonstrated that, majority of the care receivers attending ABUMC have good knowledge, perception and practices toward COVID-19. Nevertheless, certain knowledge gaps and bad practices toward the pandemic were equally identified, which highlight the need to reiterate targeted health awareness campaigns about the pandemic in Nigeria.

Introduction

The pandemic novel coronavirus disease 2019 (COVID-19) is a highly contagious viral infection caused by severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). The most

commonly reportable symptoms of the disease ranged from mild (or asymptomatic) to severe illness characterized by fever, dry cough, dyspnea, sore throat, myalgia, fatigue, nausea, diarrhea, chest

DOI: 10.21608/MID.2021.86886.1173

* Corresponding author: Bala Ningi Umar

E-mail address: umarningi5@gmail.com

© 2020 The author (s). Published by Zagazig University. This is an open access article under the CC BY-NC-ND license <https://creativecommons.org/licenses/by/4.0/>.

pain, and headache [1-3]. Rapid inter human transmission through infectious respiratory droplets has been established globally [3,4].

The disease has no specific therapy; prevention through vaccination and non-pharmacological means such as compulsory wearing of face masks in public, regular and thorough hand washing using soap and water or alcohol-based sanitizers, crowd avoidance, and maintaining physical distancing among others are known to be the only ways to avoid the spread of COVID-19 [5,6]. However, despite these precautionary measures being observed in several places, the battle against COVID-19 lingers with increased confirmed cases in many parts of the world [7].

Nigerian government like most countries in the world, have adopted the above-mentioned basic infection prevention and control measures used globally to fight previous outbreaks and pandemics. The government through Nigeria center for disease control (NCDC) has been very active in disseminating prevention messages on print and social media; and there are appreciable efforts for screening, quarantine, and providing more treatment centers for COVID-19 patients. Yet, at the time of the present study, precisely, February 28, 2021 Nigeria had about 1.22% COVID-19 mortality rate [8]. Similarly, as of May 27, 2021 the country has documented over 1.24% mortality rate; while globally, approximately 2.06% COVID-19 mortality rate was reported [9]. Therefore, there is an urgent need to assess the level of knowledge, beliefs and practices towards COVID-19 among general public, so as to curtail the persistent nationwide spread of the disease in Nigeria [10].

Moreover, in a developing country like Nigeria, which have ill-equipped healthcare system, limited trained medical personnel, economic crises, as well as widespread myths and misconceptions among general populace; the significance of preventing the rapid spread of COVID-19 cannot be over emphasized. However, people's behaviors and adherence to any disease preventive or control measures is affected by their knowledge, attitudes, perceptions, and practices towards the disease [11-14]. Central to this, are the ways and manners observed in hospitals being focal point of diseases. Furthermore, several similar KPP studies about COVID-19 have been done in Nigeria, however, high majority of them were online web-based surveys [10,15-17] hence, possible selection bias for

only those having access to the internet exist, such surveys also lack information about the elderly and few have focused on health care receivers. Therefore, the present study assessed the knowledge, perceptions and practices (KPP) towards COVID-19 among care receivers attending Ahmadu Bello University Medical Centre (ABUMC) in Nigeria.

Materials and Methods

Ethical clearance

Ethical approval for the study was obtained from the Research and Ethics Committee of ABUMC. Informed consent was gained from every participant after due explanation of the aim and objectives of the study. The principles of voluntary participation, privacy, confidentiality and anonymity of subjects were strictly considered and followed

Study location, period and participants

The cross-sectional study was conducted at ABUMC, Zaria, Nigeria. Zaria is one of the two major cities in Kaduna State, occupying part of the central high plains of Northern Nigeria. It is located at about 670m above sea level on latitude 11.11°N and longitude 7.73°E. The study was done from February 16, 2021, the week immediately after the long Nigerian lockdown of schools and religious centers to March 3, 2021. All accessible care receivers attending the university medical center within the specified period were the target population for this study. The study was carried out through convenience sampling technique (i.e., only available care receivers who were willing to participate in the study were sampled).

Study instrument

A self-administered questionnaire, designed based on the information derived from previously validated similar questionnaires [14, 18-20] was used to collect the participants' data while fully observing all COVID-19 prevention protocols. The questionnaires were distributed and monitored by the authors, and any participant who cannot accurately interpret any statement was clarified in the best language he/she understood. The questionnaire covered two main parts; the demographic characteristics of the participants such as age, gender, educational level, marital status and religion; and questions related to their KPP towards COVID-19. All the questions were closed-ended with the exception of one (Pc5) designed as semi-open to allow for incorporation of "others" category

for an opinion not explicitly captured by the available options.

The knowledge section consisted of seven questions (K1-K7) regarding the etiology, clinical manifestations, disease transmission, treatment and precautions/prevention of COVID-19 (**Table 1**). Each of the knowledge questions had four related responses which were later collapsed to three as 'correct response', 'incorrect response' and 'no

knowledge' during analyses. Assessments of the participants' perceptions towards COVID-19 comprised of eight questions (Pc1-Pc8) such as government efforts in curbing the spread of the disease, vaccines and vaccination acceptance. While the section that assessed the participants' practices towards COVID-19 has four questions (P1-P4) centered around adherence to prevention and control measures (**Table 1**).

Table 1. Self-administered questionnaire on knowledge, perceptions, and practices toward COVID-19 among care receivers attending a university medical center in Nigeria.

Questions	Options
K1. What is the cause of COVID-19?	Bacteria, Virus, 5G network, I don't know
K2. The main symptoms of COVID-19 are fever, difficulty in breathing, dry cough, and catarrh.	Strongly agree, Agree, Not agree, I don't know
K3. People with COVID -19 who do not have fever or coughing cannot infect others.	Strongly agree, Agree, Not agree, I don't know
K4. COVID -19 can be acquired through	Closed contact with infected person, Contaminated surfaces, waving infected persons, I don't know
K5. The most severe form of COVID -19 is seen among elderly people (over 60 years of age) in Nigeria	Strongly agree, Agree, Not agree, I don't know
K6. There is no effective/specific treatment for COVID -19	Strongly agree, Agree, Not agree, I don't know
K7. The use of face masks, hand hygiene, social distancing, and restriction of movement through lock down does NOT reduce the spread of COVID -19	Strongly agree, Agree, Not agree, I don't know
Pc1. Government has/is doing enough to prevent the spread of the disease	Strongly agree, Agree, Not agree, I don't know
Pc2. People who have contact with COVID -19 patients should be immediately isolated, investigated and managed.	Strongly agree, Agree, Not agree, I don't know
Pc3. Following the inception of the COVID-19 pandemic, how would you rate your awareness of the disease?	Highly aware, Aware, Partially aware, Not aware
Pc4. COVID-19 will soon be eliminated in Nigeria?	Strongly agree, Agree, Not agree, I don't know
Pc5. Please, why do you choose the above answer (Pc4)?	I believe in divine intervention, Good government policies & synergistic effect of stakeholders, Corruption and greed in managing the pandemic, Others
Pc6. As the Government is trying to acquire COVID-19 vaccines, do you wish to be vaccinated?	Yes. No. Not sure. Only when giving with incentives
Pc7. Would you buy the vaccine if available to vaccinate yourself or family member?	Yes, No, Not sure, When affordable
Pc8. Globally, some people have reservations regarding COVID-19. Please, how do you view the pandemic in Nigeria?	I believe COVID-19 is real and is claiming several lives. The disease is real, but few propagandas attached, The disease is real but many conspiracies attached, I believe it does not exist in Nigeria
P1. How often do you follow COVID-19 updates from national center for disease control (NCDC)?	Very Frequent, Frequent, Sometimes, Not at all
P2. As a preventive measure to COVID-19, how often do you wash your hands especially when you interact with others?	Very Frequent, Frequent, Sometimes, Not at all
P3. Do you wear face mask when going out of your home?	Yes, No, when reminded, only when compel to do so
P4. How often do you avoid crowd because of COVID-19?	Very Frequent, Frequent, Sometimes, Not at all

Data analyses

All the data collected were entered into Microsoft Excel and cross checked for possible error to ensure its accuracy. Descriptive statistics was applied to calculate percentages and frequencies for the participants' demographics, KPP toward COVID-19 using IBM SPSS software version 26 (NY, USA). All correct responses in the knowledge section were assigned 1 point, while incorrect answers were assigned 0 point; and the total score ranged from 0-7 with higher score signifying better knowledge about COVID-19. The mean knowledge score was obtained from the summation of the correct responses, and results were expressed as mean \pm standard deviation. The percentage knowledge scores were graded as excellent, good, fair, or poor for $\geq 80\%$, ≥ 60 but $< 80\%$, ≥ 40 but < 60 , and $< 40\%$, respectively as similarly reported by some related studies [6, 21]. Independent t-tests or one-way analysis of variance (ANOVA) where appropriate, were employed to determine the statistical differences between groups for the selected demographic variables. The statistical significance level was set at $p < 0.05$

Results

Demographic characteristics of the participants

A total of 150 respondents participated in the study, out of which 58.7% were female, while 41.3% were male. The largest age range of the participants was 16-25 years old, while the overall mean age was 29.90 ± 10.66 years. In terms of occupation, majority of the participants (60.7%) were students at various levels of education, whereas significant quota of the participants was civil servants (24.7%). More than two-thirds of the participants (62.2%) were undergraduates, 25.5% held a bachelor degree or above, and about half of them (52.7%) never married. Greater portion of the participants (53.3%) were Muslims, 40.7% were Christians, while smaller proportion (6.0%) neither believe in Islam or Christianity (Table 2).

Knowledge of the participants towards COVID-19

In general, the overall knowledge of COVID-19 among the study participants was good, with a mean knowledge score of 4.91 ± 1.68 . The participants' knowledge regarding the cause of COVID-19 was good (78.7%). Majority of the participants (93.3%) knew about the clinical manifestations of the disease, 78.7% rightly identified closed contact with infected persons and contaminated surfaces as the major source of

acquiring COVID-19 (Table 2). However, large proportion of the participants erroneously understood that COVID-19 patients who do not have fever or coughing cannot infect others. Similarly, significant part of the participants (40.7%) thought that the use of face masks, hand hygiene, social distancing, and restriction of movement through lock down does not reduce the spread of COVID -19 (Table 3).

When the participant's knowledge was compared among the different demographics (Table 3), female participants (5.1 ± 1.6) were found to be more knowledgeable about COVID-19 than male (4.6 ± 1.7), however, the difference was not statistically significant ($p > 0.05$). The youths, particularly age range of 16-25 years (5.2 ± 1.5) and 26-35 years (5.1 ± 1.6) had significantly ($p < 0.05$) higher COVID-19 knowledge score as compared to the elderly who had the least COVID-19 knowledge score (2.3 ± 2.5). Undergraduate students were found to have significantly ($P < 0.05$) higher knowledge score (4.8 ± 1.6) than those at secondary (2.9 ± 1.8) and primary schools (3.3 ± 2.9). Likewise, there was a significant difference ($P < 0.05$) between Muslims (5.2 ± 1.7) and Christian respondents (4.8 ± 1.7) regarding the knowledge about COVID-19 (Table 2).

Participants perception towards COVID-19

The inclusive participant's perceptions towards COVID-19 are presented in table (4). Larger proportion of the participants (42.0%) was convinced that the Nigerian government is doing enough to prevent the spread of COVID-19; however, significant portion of the participants (34.7%) believed the opposite. Furthermore, nearly all the participants (84.0%) held positive views towards immediate quarantine of COVID-19 patients or people who had contact with the patients. Although, majority of the participants (56.0%) had confidence that COVID-19 will soon be eliminated in Nigeria, most of them cited their faith in divine intervention (36.7%), rather than good government policies and synergistic effects of stakeholders (17.3%) in curbing the disease. Similarly, about 16.0% of the participants thought otherwise citing inadequate government will, corruption and greed in managing the pandemic as their main reasons. Nearly half of the respondents (47.3%) did not wish to be vaccinated against COVID-19 even if the vaccines are freely available, 19.3% are unsure if they would accept the vaccines, whereas 6.0% are

likely to accept only when giving with incentives. However, despite the obvious negative economic impact with scoring morbidity of COVID-19, some of the participants (12.0%) believed that the disease does not exist in Nigeria.

Participants practices toward COVID-19

The participants attachment to Nigeria center for disease control concerning COVID-19 updates was low (41.4%), with approximately half of the respondents hardly following the national center for information regarding the pandemic. Nevertheless, majority of the participants (65.3%) regularly washed their hands with soap or alcohol-based sanitizer as a means to protect themselves

from getting COVID-19. However, large share of the respondents (32.0%) rarely practiced this important preventive measure; while about 2.7% never observed hand hygiene even after interacting with others. The study also showed that, though a high majority of the participants (87.3%) wore face masks when going out of their homes, 7.3% of the participants wear the masks only when compel to do so. In addition, most of the participants (61.3%) regularly avoid crowd to protect themselves from COVID-19 while some (10.7%) still socialized in defiance to government directives of physical distancing (**Table 5**).

Table 2. Demographic characteristics and knowledge about COVID-19 among care receivers attending a university medical center in Nigeria.

Characteristics	No. of participants (%)	Mean knowledge score (SD)	t / F	p – value
Gender				
Male	62 (41.3)	4.6 (1.7)	-1.855	0.066
Female	88(58.7)	5.1 (1.6)		
Age (years)				
16-25	68(45.3)	5.2 (1.5)	3.139	0.016
26-35	42(28.0)	5.1 (1.6)		
36-45	24(16.0)	4.5 (1.9)		
46-55	13(8.7)	4.4 (1.6)		
Above 55	3(2.0)	2.3 (2.5)		
Occupation				
Student	91(60.7)	5.1 (1.4)	2.012	0.115
Employed	37(24.7)	4.8 (1.9)		
Entrepreneur	18(12.0)	4.4 (1.9)		
Unemployed	4(2.7)	3.5 (2.6)		
Marital status				
Never married	79(52.7)	5.1 (1.6)	1.686	0.189
Married	60(40.0)	4.6 (1.9)		
Others	11(7.3)	5.5 (1.0)		
Educational status				
Primary	3(2.0)	3.3 (2.9)	5.983	0.000
Secondary	8(5.3)	2.9 (1.8)		
Undergraduate	93(62.0)	4.8 (1.6)		
Postgraduate	38(25.3)	5.4 (1.3)		
Others	8(5.3)	5.9 (1.5)		
Religion				
Islam	80(53.3)	5.2 (1.7)	3.164	0.045
Christianity	61(40.7)	4.8 (1.7)		
Others	9(6.0)	3.8 (1.0)		

Note: The category “others” under educational status denotes people with informal education, they have no certificates but can read and write

Table 3. Knowledge about COVID-19 among care receivers attending a university medical center in Nigeria (n = 150).

Question (hints)	Correct response (%)	Incorrect response (%)	No knowledge (%)
K1). Causes of COVID-19	118 (78.7)	22 (14.7)	10 (6.7)
K2). Symptoms of COVID-19	140 (93.3)	1 (0.7)	9 (6.0)
K3). Infectivity of patients without fever	93 (62.0)	38 (25.3)	19 (12.7)
K4). Transmission of COVID-19	118 (78.7)	8 (5.3)	24 (16.0)
K5). Most severe form of COVID-19	105 (70.0)	21 (14.0)	24 (16.0)
K6). No effective treatment for COVID-19	92 (61.3)	18 (12.0)	40 (26.7)
K7). Status of non-pharmacologic preventions	71 (47.3)	61 (40.7)	18 (12.0)

Table 4. Perceptions about COVID-19 among care receivers attending a university medical center in Nigeria (n = 150).

Question	Options	Responses	
		Frequency	Percentage (%)
Pc1). Is Nigerian government trying to control COVID-19?	Strongly agree	24	16.0
	Agree	54	36.0
	Not agree	52	34.7
	I don't know	20	13.3
Pc2). Should COVID-19 patients be quarantined?	Strongly agree	85	56.7
	Agree	41	27.3
	Not agree	18	12.0
	I don't know	6	4.0
Pc3). How do you rate your awareness about COVID-19?	Highly aware	70	46.7
	Aware	56	37.3
	Partially aware	15	10.0
	Not aware	9	6.0
Pc4). COVID-19 will be controlled?	Strongly agree	35	23.3
	Agree	49	32.7
	Not agree	24	16.0
	I don't know	42	28.0
Pc5). Reasons for choosing Pc4?	Divine intervention	55	36.7
	Good government policies	26	17.3
	Corruption and greed	53	35.3
	Others	16	10.7
Pc6). Do you wish to be vaccinated?	Yes	41	27.3
	No	71	47.3
	Not sure	29	19.3
	Only with incentives	9	6.0
Pc7) Would you buy the vaccines?	Yes	22	14.7
	No	63	42.0
	Not sure	31	20.7
	When affordable	34	22.7
Pc8). How do you view COVID-19?	COVID-19 is real & lethal	31	20.7
	Is real with few propagandas	50	33.3
	Is real but many conspiracies	51	34.0
	It does not exist in Nigeria	18	12.0

Table 5. Practices toward COVID-19 among care receivers attending a university medical center in Nigeria (n = 150).

Question	Options	Responses	
		Frequency	Percentage (%)
P1). How often do you follow NCDC for COVID-19 updates?	Very frequent	16	10.7
	Frequent	46	30.7
	Sometimes	72	48.0
	Not at all	16	10.7
P2). How often do you practice hand hygiene?	Very frequent	35	23.3
	Frequent	63	42.0
	Sometimes	48	32.0
	Not at all	4	2.7
P3). Do you wear face mask?	Yes	131	87.3
	No	4	2.7
	When reminded	4	2.7
	Only when compelled	11	7.3
P4). How often do you avoid crowd?	Very frequent	42	28.0
	Frequent	50	33.3
	Sometimes	42	28.0
	Not at all	16	10.7

Discussion

The high level of theoretical knowledge about COVID-19 recorded in the present study denotes good awareness of the disease among the study participants, and is in line with the findings of similar surveys conducted in different parts of the country [21-23]. Likewise, the significant difference in COVID-19 knowledge scores observed in the present study among different educational backgrounds is generally consistent with previous studies on COVID-19 in several parts of the world [6,24,25]. Similarly, the significantly higher knowledge score reported among youths, particularly those within the age range of 16-25years, and 26-35years old might not be far from the fact that these active age brackets are relatively more exposed to information regarding COVID-19 through both print and social media as earlier noted by **Ladiwala et al.** [6], **Gohel et al.** [20], **Bekele et al.** [26], and **Yue et al.** [27] in Pakistan, India, Ethiopia, and China, respectively. Thus, health education intercession would be most effective if it targets certain socio-demographics like elderly men and people with informal education. Conversely, the commonly reported significant difference between male and female gender was not observed in the present study. This could be chiefly due to the high share of male students in the current study as

compared to the relatively larger proportion of older females in the previous studies [6, 17, 25].

Broadly, knowledge regarding COVID-19 directly influenced people's perceptions such as perceived risk, government policies, and efficacy belief towards COVID-19 [14]. Therefore, it can be presumed that most of those who possessed poor knowledge about the disease would have negative perceptions and risky practices toward COVID-19. However, it is worth noting that in some instances, acquisition of knowledge has not vividly influenced practices towards COVID-19; possible action gap in applying the acquired knowledge into practices has been documented [26]. Consequently, we encourage Nigerian government and concerned stakeholders to set priorities in information campaigns about COVID-19, and go beyond the rhetoric of lectures and presentation. There should be a resolute, targeted and continuous public health education focusing on providing awareness about the disease while stressing on the implementation of the acquired prevention knowledge into good practices.

Non-pharmacologic methods of COVID-19 prevention have been widely employed as the major means of preventing COVID-19. Their significance as the most critical approach for decreasing the risk of COVID-19 transmission has been well elucidated [5, 20, 28, 29]. Although, the

present study reveals regular use of face mask, hand washing, and social distancing among majority of the participants, appreciable proportion of the participants did not practice these simple, yet effective measures to reduce the spread of the pandemic. Similarly, in spite of the few controversies surrounding COVID-19 vaccines, vaccination remains the only way to acquire long lasting immunity to the virus; hitherto majority of the participants were not willing to be vaccinated against the disease, and this calls for immediate and resolute enlightenment.

The strength of the current study lies in the fact that it has covered the KPP of an understudied community of public which played a critical role in the fight against COVID-19. Moreover, data presented in the study were self-reported, hence, the issue of possible demographic selection bias of those having access to internet only, as seen in web-based surveys was avoided. However, the study has few limitations which cannot be ignored. This includes the use of convenience sampling technique with limited sample representativeness, as well as failure to subject the adopted validated questionnaire to additional (dual) validation to increase the reliability of the study.

Conclusions

The current study has established that while majority of the care receivers attending ABUMC showed high level of knowledge and fairly good practices towards COVID-19 prevention and control measures; it is apparent that some people do not consider the danger posed by the pandemic serious, and are not following the government recommendations regarding the disease. The study has also highlighted the specific aspects of the KPP where incorrect responses were documented, and need to be improved. Similarly, in contrast to the high knowledge score among relatively young female care receivers; the study has equally unraveled serious misconceptions among the participants regarding the existence of COVID-19 in Nigeria. Thus, synergistic and concerted efforts from government, religious bodies and concerned stakeholders are recommended to correct the wrong perceptions, and improve the knowledge of the care receivers toward controlling the pandemic. Special preference should be giving to elderly males, unemployed individuals, people with low educational levels and unmarried because of their relatively poor knowledge scores about COVID-19.

Conflict of interest

The authors have no conflicts of interest to declare that are relevant to the content of this article.

Funding source

This research has not been funded with any specific grant from funding agencies in the public, commercial, or not-for-profit sectors.

Authors' contributions

Bala Ningi Umar conceived and designed the study, and wrote the original draft of the manuscript; Nuhu Muhammad administered the project, collected and analyzed the data; while Kabiru Haliru Ahmad, Khadijah Usman, and Rahman Khaleequr participated in administering the questionnaires, revised the article critically and approved its intellectual content. All authors have reviewed and approved the final version of the manuscript.

Acknowledgement

The authors wish to thank the management of Ahmadu Bello University, Zaria for providing a conducive environment to undertake the study. We are indebted to the study participants who spared their precious time to respond to the questionnaire. We are equally grateful to Dr. M. Babashani of Veterinary Teaching Hospital, ABU, Zaria for his guidance towards data analyses.

Other numerous professional colleagues who assisted in making the study a successful one are well appreciated.

Data availability

The datasets generated during and/or analyzed during the current study are available from the corresponding author on reasonable request

References

- 1-**Adhikari SP, Meng S, Wu YJ, Mao YP, Ye RX, Wang QZ et al.** Epidemiology, causes, clinical manifestation and diagnosis, prevention and control of coronavirus disease (COVID-19) during the early outbreak period: a scoping review. *Infect Dis Poverty* 2020; 9(1):1–12.
- 2-**Mina FB, Billah M, Karmakar S, Das S, Hasan F, Acharjee UK.** An online observational study assessing clinical characteristics and impacts of the COVID-19 pandemic on mental health: a perspective study from Bangladesh. *J Public Health: From*

- Theory to Pract 2020; <https://doi.org/10.1007/s10389-020-01445-2>
- 3-**Shereen MA, Khan S, Kazmi A, Bashir N, Siddique R.** COVID-19 infection: Origin, transmission, and characteristics of human Coronaviruses. *J Adv Res* 2020; 24(1): 91–98
- 4-**Guo YR, Cao QD, Hong ZS, Tan YY, Chen SD, Jin HJ et al.** The origin, transmission and clinical therapies on coronavirus disease 2019 (COVID-19) outbreak—an update on the status. *Mil Med Res* 2020; 7(1):1–10.
- 5-**Centre for Disease Control and Prevention.** Coronavirus Disease 2019 (COVID-19) Social Distancing, Quarantine, and Isolation. 2020; Accessed on October 20, 2020 from www.cdc.gov/coronavirus/2019-ncov/preventgetting-sick/social-distancing.html. 2020.
- 6-**Ladiwala ZFR, Dhillon RA, Zahid I, Irfan O, Khan MS, Awan S, Khan JA.** Knowledge, attitude and perception of Pakistanis towards COVID-19; a large cross-sectional Survey. *BMC Public Health* 2021; 21: 1-10
- 7-**European Centre for Disease Prevention and Control (ECDC).** COVID-19 situation updates worldwide, as of week 30, updated 5 August 2021. Available at <https://www.ecdc.europa.eu/en/geographical-distribution-2019-ncov-cases/> Accessed on August 11, 2021.
- 8-**Nigeria Centre for Disease Control (NCDC).** Nigeria-COVID-19 update. 2021; Available at <https://covid19.ncdc.gov.ng/> Accessed on June 2, 2021.
- 9-**World Health Organization (WHO).** Coronavirus disease (COVID-19) Weekly Epidemiological Update and Weekly Operational Update. 2021; Available at www.who.int/emergencies/diseases/novel-coronavirus-2019/ Accessed July 2, 2021.
- 10-**Reuben RC, Danladi MM, Saleh DA, Ejembi PE.** Knowledge, attitudes and practices towards COVID-19: an epidemiological survey in North-Central Nigeria. *J Community Health* 2020; 1-14.
- 11-**Janz NK, Becker MH.** The health belief model: A decade later. *Health Educ Q* 1984; 11(1): 1-47.
- 12-**Ramadhani HO, Muiruri C, Maro VP, Omondi M, Mushi JB, Lirhunde ES, Bartlett JA.** Association of knowledge on ART line of treatment, scarcity of treatment options and adherence. *BMC Health Serv Res* 2016; 16(1): 1-7.
- 13-**Akalu Y, Ayelign B, Molla MD.** Knowledge, attitude and practice towards COVID-19 among chronic disease patients at Addis Zemen Hospital, Northwest Ethiopia. *Infect Drug Resist* 2020; 13(1): 1949.
- 14-**Zhong B, Luo W, Li H, Zhang Q, Liu X, Li W, Li Y.** Knowledge, attitudes, and practices towards COVID-19 among Chinese residents during the rapid rise period of the COVID-19 outbreak: a quick online cross-sectional Survey. *Int J Biol Sci* 2020; 16(10): 1745-1752.
- 15-**Adesegun OA, Binuyo T, Adeyemi O, Ehioghae O, Rabor DF, Amusan O et al.** The COVID-19 crisis in Sub-Saharan Africa: Knowledge, attitudes, and practices of the Nigerian public. *Am J Trop Med Hyg* 2020; 103(5): 1997-2004.
- 16-**Ejeh FE, Saidu AS, Owoicho N, Maurice NA, Jauro S, Madukaji L, Okon OO.** Knowledge, attitude, and practice among healthcare workers towards COVID-19 outbreak in Nigeria. *Heliyon* 2020; 6(11): e05557
- 17-**Adefisoye JO, Adefisoye MA, Bisi-Johnson MA, George F.** Knowledge, Attitude and Perception of People towards the Novel Coronavirus Disease (COVID-19) Outbreak

- and its Effects on their Livelihoods: Perspectives from Diverse Populations. *Health Econ Outcome Res* 2021; 7(2): 033-042
- 18-**Bhagavathula AS, Aldhaleei WA, Rahmani J, Mahabadi MA, Bandari DK.** Novel Coronavirus (COVID-19) Knowledge and Perceptions: A Survey on Healthcare workers. *MedRxiv* 2020; (1): 1-15.
- 19-**Ogolodom MP, Mbaba AN, Alazigha N, Erundu OF, Egbe NO, Golden I et al.** Knowledge, attitudes and fears of healthCare workers towards the corona virus disease (COVID-19) pandemic in South-South, Nigeria. *Health Sci J* 2020; 1(2): 1-10.
- 20-**Gohel KH, Patel PB, Shah PM, Patel JR, Pandit N, Raut A.** Knowledge and perceptions about COVID-19 among the medical and allied health science students in India: An online cross-sectional survey. *Clin Epidemiol Glob Health* 2021; 9:104–109.
- 21-**Umezudike KA, Isiekwe IG, Fadeju AD, Akinboboye BO, Aladenika ET.** Nigerian undergraduate dental students' knowledge, perception, and attitude to COVID-19 and infection control practices. *J Dent Educ* 2021; 85:187–196.
- 22-**Egbi OG, Duru C, Kasia B.** Knowledge, attitude and practice towards COVID-19 among workers of a tertiary hospital in Bayelsa State, Nigeria. *Pan Afr Med J* 2020; 37(1): 24.
- 23-**Ogbole AJ, Bisji JS, Umar SJ, Jallo IM., Ezeh SO, James AL.** Knowledge, attitudes and perception in regard to COVID-19 pandemic in Nigerian military population. *Adv Soc Sci Res J* 2020; 7(11): 231-249.
- 24-**Azlan AA, Hamzah MR, Sern TJ, Ayub SH, Mohamad E.** Public knowledge, attitudes and practices towards COVID-19: A cross-sectional study in Malaysia. *PLoS One* 2020; 15(5): e0233668.
- 25-**Lee M, Kang B, You M.** Knowledge, attitudes, and practices (KAP) toward COVID-19: a cross-sectional study in South Korea. *BMC Public Health* 2021; 21: 1-10
- 26-**Bekele D, Tolossa T, Tsegaye R, Teshome W.** The knowledge and practice towards COVID-19 pandemic prevention among residents of Ethiopia. An online cross-sectional study. *PLoS One* 2021; 16(1): e0234585.
- 27-**Yue S, Zhang J, Cao M, Chen B.** Knowledge, attitudes and practices of COVID-19 among urban and rural residents in China: A cross-sectional study. *J Community Health* 2021; 46: 286–291
- 28-**Meng L, Hua F, Bian Z.** Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res* 2020; 99(5):481-487.
- 29-**World Health Organization (WHO).** Health Topics: Coronavirus. 2020; Available at <https://www.who.int/health-topics/> Accessed on November 7, 2020.