

Five-times sit-to-stand test following anterior cruciate ligament surgery: a cross-sectional reliability study

Hikmet KOCAMAN^{1*}, Halil ALKAN², Mehmet YETİŞ³, Mehmet CANLI⁴, Şafak KUZU⁴, Anıl ÖZÜDOĞRU⁴

1.Faculty of Health Sciences, Department of Physiotherapy and Rehabilitation, Karamanoglu Mehmetbey University, Karaman, Turkey

2.Faculty of Health Sciences, Department of Physiotherapy and Rehabilitation, Muş Alparslan University, Muş, Turkey

3.Department of Orthopaedics and Traumatology Faculty of Medicine, Kırşehir Ahi Evran University, Kırşehir, Turkey

4.School of Physical Therapy and Rehabilitation, Kırşehir Ahi Evran University, Kırşehir, Turkey

*Corresponding Author: Hikmet KOCAMAN; E-mail: m

Abstract

Aims

Patients who have had anterior cruciate ligament reconstruction (ACL-R) should periodically have their muscle strength assessed. The five-times sit-to-stand test (FTSST) can evaluate the muscle strength and balance of the lower extremities. This study's primary purpose was to assess the validity and reliability of the FTSST in patients who have undergone ACL-R.

Material and Methods

Forty-three people who had undergone ACL-R surgery were included in the study. The study's primary outcome measure, the FTSST, was assessed by two different investigators. Secondary outcome measures were body balance, quadriceps muscle strength, Tegner activity score (TAS), and Lysholm score.

Results

The FTSST's test-retest and inter-rater reliability were both high (ICC: 0.99). The FTSST also showed a strong statistically significant correlation with all secondary outcome measures, including balance, quadriceps muscle strength, TAS, and Lysholm score ($p < 0.05$).

Conclusions

According to the study results, the FTSST is a tool-free, simple method for assessing muscle strength and the body balance level, mobility level, and functional status of the knee in patients who have undergone ACL-R surgery.

Key words: Anterior cruciate ligament, Reconstruction, Five-times sit-to-stand test

Introduction

Anterior cruciate ligament (ACL) rupture is the most common knee injury and is seen primarily in adults aged 20–29. The ACL is often injured during sports activities that involve turning, jumping, and contact¹. ACL reconstruction (ACL-R) aims to restore knee stabilization, but deficiencies in knee extension strength, neuromuscular control, and proprioception persist for up to 2 years after reconstruction surgery². These deficiencies are also present in the contralateral uninjured knee³⁻⁵. Therefore, evaluation programs after ACL-R should include both knees.

Quadriceps muscle weakness and dysfunction are the most frequent symptoms in patients who have undergone ACL-R and may also occur in the years following reconstruction^{6,7}. Atherogenic muscle inhibition, which affects the periarticular musculature, is the underlying mechanism that results from damaged joint components. Theoretically, inaccurate sensory input from joint mechanoreceptors causes a reduction in efferent impulses to the quadriceps muscle in patients with knee joint injury. Consequently, a decreased number of motor units are triggered during contraction^{7,8}. Grindem et al. stated that quadriceps muscle weakness after ACL-R increases the risk of injury⁹. Therefore, continuous evaluation of quadriceps muscle strength is required after reconstruction.

Quadriceps muscle strength has traditionally been measured in clinical settings with isokinetic dynamometers or isometric dynamometers¹⁰. However, patients who require long-term follow-up, such as those who have undergone ACL-R, may find it difficult to come to clinical settings on a regular basis. Therefore, clinicians are trying to reach patients through telerehabilitation methods and frequently resort to easy, equipment-free methods.

The five-times sit-to-stand test (FTSST) is a simple, rapid, equipment-free method that evaluates balance and mobility, as well as lower extremity muscle strength^{11,12}. The FTSST's validity and reliability have been examined in different populations and diseases, with the conclusion that it is a valid and reliable method¹³⁻²⁰. After ACL-R, rapid and reliable tests such as FTSST that evaluate balance, mobility, and lower extremity muscle strength are very important in monitoring rehabilitation results. However, we could not find any study examining its validity and reliability in people with ACL-R. Thus, the aim of this study was to test the inter-rater and test-retest reliability of the FTSST in patients who had undergone ACL-R surgery. Additionally, it was to examine correlations between FTSST and other parameters.

Methods

Design

This cross-sectional study was approved by the Muş Alparslan

University Clinical Research Ethics Committee (Decision no: 1-2022/39). The Helsinki Declaration's guiding principles were followed during the study's execution. Written and verbal consent was obtained from the participants just before starting the study. Participants were randomly selected from those who had previously undergone ACL reconstruction surgery and presented applied to the Kırşehir Training and Research Hospital Orthopedics and Traumatology Clinic for a neuromuscular control exam. Then, two physiotherapists with four years of clinical expertise in ACL rehabilitation conducted each assessment one hour apart. Seven days later, the same physiotherapists conducted the evaluations once more.

Participants

Patients who underwent ACL-R surgery and were at least six months post-reconstruction and over the age of 18 were included in the study. Those who had a meniscus or posterior cruciate ligament injury accompanying an ACL injury and those who had undergone any surgery other than ACL surgery were excluded from the study. In addition, those with any orthopedic, vestibular disorders or neurological in addition to ACL-R, were excluded from the study. Current guidelines recommend the incorporation of 30 to 50 participants in reliability studies²¹. Considering the lack of precedent research concerning the validity and reliability of the FTSST among individuals with ACL-R, we diligently followed this guideline and recruited a total of 46 patients with ACL-R for the present study.

Assessments

Patients' demographic and clinical characteristics (age, height, gender, weight, injured knee, and time elapsed after surgery) were recorded. Then, clinical evaluations were started.

FTSST Test

For the FTSST, the participants were asked to sit on a 42-45 cm chair with their hands crossed on their chest, and to sit and stand up from the chair five times with the start command, respectively. The test score was assessed with a stopwatch, and the time to complete the task was recorded in seconds²².

Balance

The Biodex Stability System (BSS) [(BSS) Biodex Inc., Shirley, New York] was used to measure body balance. The BSS consists of a transportable balance platform with a 20° tiltable surface connected to computer software that provides an objective assessment of balance. Each participant in the study received information about the tests and guidelines for compliance. In both static and dynamic balance measurements, we recorded three distinct postural sway amounts in centimeters, resulting in a total of six outcome measures: anterior-posterior (AP), medial-lateral (ML), and overall score (Overall score-OS). Each session of static and dynamic balance measurement, lasting 20 seconds each, was repeated three times with 10-second intervals of rest between measurements. The device automatically computed the averages for the static and dynamic balance parameters (AP, ML, and OS)²³.

Muscle Strength

Quadriceps muscle strength of the participants was evaluated with isometric dynamometer. While the participants were

seated on a flat surface and the knees and hips were 90° flexed, the dynamometer was placed above the malleolus and the participant was asked to push the dynamometer with maximum force. Three attempts were made and the highest score was recorded and recorded in kg²⁴.

Lysholm Score

The Lysholm score was developed in the 1980s as a patient-focused subjective assessment method to assess the functional status of patients with ACL injuries [23]. Çelik et al. developed a Turkish version of the Lysholm score and found it to be a valid and reliable method. The Lysholm score consists of eight questions, and the total score is in the range of 0–100. A high score indicates good knee functional status²⁵.

Tegner Activity Scale

The Tegner activity scale is a scoring system that measures activity levels in daily life and sports from 0 to 10. It has been found to have acceptable responsiveness for use in the early return to function following ACL treatment. Activity level is graded from 0 points for those who quit the activity due to injury or dysfunction to 10 points for those who play professional sports at the national team level^{26,27}.

Statistical Analysis

SPSS software (version 24) was used to conduct the statistical analysis. Using visual and analytical techniques, the variables' conformance to normal distribution was evaluated (Shapiro–Wilk tests). Since the data were normally distributed, descriptive statistics were given as the mean and standard deviation. Nominal variables were expressed as numbers and percentages. The inter-rater and test–retest reliability of the FTSST were assessed using intraclass correlation coefficients (ICC). The ICC is rated as good (0.60–0.80) or excellent (0.80–1.0) 28. The FTSST's concordance validity and relationship with other tests were assessed with the Pearson correlation test. The correlation coefficient was considered low between 0.05 and 0.4, moderate between 0.4 and 0.7, and high between 0.7 and 1.0 29. A statistically significant level was defined as a p-value of 0.05 or lower.

Results

The demographic and descriptive characteristics of the patients are given in Table 1.

The values of the FTSST, balance, muscle strength, Lysholm, and Tegner activity scores of the ACL patients are given in Table 2.*

When the test–retest (ICC) and inter-rater (ICC) reliability of the FTSST were examined, the first assessor's test–retest result, the ICC value, was found to be 0.997. Regarding consistency (inter-rater) between the first and second assessors, the ICC value was found to be 0.997. According to these results, the test–retest and the inter-rater agreement were interpreted as excellent (Table 3).

The correlation results of the Biodex Stability System, Tegner activity score, Lysholm score, and quadriceps muscle strength (injured and uninjured knees) are given in Table 4. There was a high positive correlation between FTSST and static-dynamic parameters of the Biodex Stability System ($p < 0.05$).

Table 1. Demographic and clinical characteristics of ACL-R patients

		(n=46)			
		Mean	SD	Min	Max
Age (years)		33.26	6.22	20	44
Height (cm)		165.52	5.10	155	190
Weight (kg)		76.35	10.86	58	96
BMI (kg/m²)		27.39	6.61	20.9	34.52
Post-surgery time (month)		8.70	1.96	6	12
		n		(%)	
Gender	Male	34		73.9	
	Female	12		26.1	
Dominant side	Right	12		26.1	
	Left	34		73.9	
Injured Knee	Left	14		30.4	
	Right	32		69.6	

N: Number, %: Percent, SD: Standard deviation, BMI: Body mass index

Table 2. Measurement values of the ACL-R patients

		(n=46)		Min	Max
		Mean	SD		
Balance	Static AP (cm)	1.9	1.2	0.2	4.1
	Static ML (cm)	1.3	0.9	0.4	3.8
	Static-Overall (cm)	2.8	1.8	0.7	6.7
	Dynamic-AP (cm)	2.6	1.5	0.4	6.1
	Dynamic-ML (cm)	1.7	0.8	0.2	3.9
	Dynamic-Overall (cm)	3.6	2.0	0.9	7.6
Tegner activity score		2.5	1.7	0.0	5
Lysholm score		51.1	21.3	15.0	85
Quariceps muscle strength (kg)	Injured	9	3.7	4.0	15
	Uninjured	10.4	4.1	4.0	17.5
FTSST (1) (sec)	Test	14.6	4.5	6.8	20.6
	Retest	14.8	4.7	6.1	20.7
FTSST (2) (sec)	Test	14.8	4.4	6.9	20.4
	Retest	14.9	4.7	6.4	21.2

cm: centimeter, n: Number, sec: second, 1: First assessor, 2: Second assessor, SD: Standard deviation, AP: Anterior-posterior, ML: Medial-lateral, FTSST: Five-times sit-to-stand test

Table 3. Inter-rater (ICC) and test–retest (ICC) reliability of the FTSST

n=46	Difference (Mean±SD)	Inter-rater (ICC _{1,2}) (95% CI)	Test-retest (ICC _{1,1}) (95% CI)	SEM	MDC ₉₅
FTSST	0.20±0.48	0.997 (0.994-0.998)	0.997 (0.995-0.998)	0.02	0.05

n; Number, SD; Standard deviation, FTSST; Five-times sit-to-stand test, ICC; Intraclass correlation coefficient, CI; Confidence interval, SEM; Standard error of measurement; MDC₉₅; Minimum detectable change at the 95% confidence interval

Table 4. Correlation of the FTSSST with other tests

			FTSSST
Balance	Dynamic-AP (cm)	r	0.873
		p	<0.001
	Dynamic-ML (cm)	r	0.843
		p	<0.001
	Dynamic-Overall (cm)	r	0.881
		p	<0.001
	Static-AP (cm)	r	0.877
		p	<0.001
	Static-ML (cm)	r	<0.001
		p	<0.001
	Static-Overall (cm)	r	0.881
		p	<0.001
Tegner activity score		r	-0.912
		p	<0.001
Lysholm score		r	-0.793
		p	<0.001
Quadriceps muscle strength (kg)	Injured Knee	r	-0.896
		p	<0.001
	Uninjured Knee	r	-0.900
		p	<0.001

FTSSST: Five-times sit-to-stand test, AP: Anterior-posterior, ML: Medial-lateral, p<0.001

There was a statistically significant negative correlation between the FTSSST, Tegner activity score, Lysholm score, and quadriceps muscle strength (injured and uninjured knee) (p<0.05).

Discussion

The results of the current study demonstrate that the FTSSST has excellent test–retest and inter-rater reliability in patients who have undergone ACL-R. In addition, the FTSSST was found to be highly correlated with muscle strength, balance, activity level, and knee functionality in patients with ACL-R. These results support the validity of the FTSSST.

The validity and reliability of the FTSSST have previously been examined in intensive care patients¹³, patients with total hip arthroplasty¹⁶, patients with balance disorders¹⁴, older adults³⁰, chronic stroke patients³¹, and patients with pelvic girdle pain due to pregnancy³², and it has been concluded that the FTSSST is a valid and reliable method. In the present study, in parallel with the literature, we concluded that the FTSSST has excellent test–retest reliability in patients with ACL reconstruction. In light of these results, the FTSSST is a useful tool in patients with ACL-R.

The literature often refers to peripheral changes (morphological and cellular) in the quadriceps and weakness of the quadriceps following ACL-R³³⁻³⁵. This weakness and the resulting dysfunction have been shown to cause a decrease in knee-related quality of life, the development of

osteoarthritis, and decreased functionality^{6,36}. Noehren et al. stated that despite physical therapy after ACL-R, there was a loss of strength in the vastus lateralis muscle³⁵. In another study, Konisi et al. reported that dysfunction of the quadriceps muscles after reconstruction persisted for a long time in the uninjured knee³⁷.

In present study, a correlation was found between knee extension muscle strength and the FTSSST test. This may be because the quadriceps muscle is an antigravity muscle that plays an important role in both standing and sitting.

Proprioception is one of the most important sources of sensory information in maintaining body balance. After ACL reconstruction, proprioception decreases due to the removal of the Pacini and Ruffini bodies in the joint during surgery. Moussa et al. compared the postural stability of patients 2 years after ACL-R with that of healthy people and concluded that postural stability was worse in patients who had undergone ACL-R 38. Bonfirm et al. reported that ACL lesions have adverse effects on postural balance and proprioception³⁹. Similarly, in a study, Tookuni et al. stated that a unilateral knee ACL injury disrupts balance⁴⁰. The FTSSST assesses balance ability and muscle strength¹¹. Tiwari et al. stated that the strongest predictors of FTSSST performance in older adults were lower extremity muscle strength, and dynamic balance⁴¹. In another study, concluded that there is a strong correlation between the FTSSST and the Berg Balance Scale¹². In the present study, there was a strong

correlation between FTSSST and dynamic–static balance.

ACL injuries cause limitations in range of motion, loss of strength, muscle imbalance, atrophy, and functional instability of the knee. An attempt is made to regain stability and knee functionality through reconstruction surgery and an appropriate rehabilitation program. Patients are supposed to be followed for six months in rehabilitation clinics, but deficiencies in stabilization, mobility, and functionality can last for up to two years after surgery. Goldberg et al. stated that FTSSST is related to functional mobility in older adults⁴². Medina-Mirapeix et al. concluded that the FTSSST is related to functional mobility tests in patients with total knee arthroplasty⁴³. In the present study, a statistically significant relationship was found between the FTSSST and the Tegner activity score, which evaluates mobility, and the Lysholm score, which evaluates knee functionality. This result may be due to the fact that the sit-stand activity is also a mobility activity.

Study Limitation

The most important limitation of this study is that it was not conducted on professional athletes. Considering that ACL injury is common in professional athletes, it is recommended that future studies be conducted on different groups of athletes who have undergone ACL-R.

Conclusion

In conclusion, the FTSSST is a valid and reliable method for assessing muscle strength, balance, and mobility in patients who have undergone ACL-R. FTSSST may be preferred by clinicians and researchers working on ACL rehabilitation because it is simple, does not require equipment, and enables both face-to-face and remote evaluation.

Declarations

Conflict of interest

None

Funding

None

Authors' Contributions

HK: Conceptualization, Data curation, Formal Analysis, Methodology, Project administration Writing – review & editing. MY: Writing – original draft, Conceptualization, Formal Analysis. MC: Conceptualization, Writing – original draft. HA: Conceptualization, Writing – original draft. ŞK: Conceptualization, Formal Analysis. AÖ: Data curation, Writing – original draft.

References

- Majewski M, Susanne H, Klaus S. Epidemiology of athletic knee injuries: A 10-year study. *Knee*. Jun 2006;13(3):184-188.
- Nagelli CV, Hewett TE. Should Return to Sport be Delayed Until 2 Years After Anterior Cruciate Ligament Reconstruction? Biological and Functional Considerations. *Sports Med*. Feb 2017;47(2):221-232.
- Chung KS, Ha JK, Yeom CH, et al. Are Muscle Strength and Function of the Uninjured Lower Limb Weakened After Anterior Cruciate Ligament Injury? Two-Year Follow-up After Reconstruction. *Am J Sports Med*. Dec 2015;43(12):3013-3021.
- Lepley AS, Gribble PA, Thomas AC, Tevald MA, Sohn DH, Pietrosimone BG. Quadriceps neural alterations in anterior cruciate ligament reconstructed patients: A 6-month longitudinal investigation. *Scand J Med Sci Sports*. Dec 2015;25(6):828-839.

- Negahban H, Mazaheri M, Kingma I, van Dieën JH. A systematic review of postural control during single-leg stance in patients with untreated anterior cruciate ligament injury. *Knee Surg Sports Traumatol Arthrosc*. Jul 2014;22(7):1491-1504.
- Palmieri-Smith RM, Thomas AC. A neuromuscular mechanism of posttraumatic osteoarthritis associated with ACL injury. *Exerc Sport Sci Rev*. Jul 2009;37(3):147-153.
- Rice DA, McNair PJ. Quadriceps arthrogenic muscle inhibition: neural mechanisms and treatment perspectives. *Semin Arthritis Rheum*. Dec 2010;40(3):250-266.
- Hopkins JT, Ingersoll CD. Arthrogenic muscle inhibition: a limiting factor in joint rehabilitation. *Journal of sport rehabilitation*. 2000;9(2):135-159.
- Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after ACL reconstruction: the Delaware-Oslo ACL cohort study. *British journal of sports medicine*. 2016;50(13):804-808.
- Al Amer HS, Sabbahi MA, Alrowayeh HN, Bryan WJ, Olson SL. Electromyographic activity of quadriceps muscle during sit-to-stand in patients with unilateral knee osteoarthritis. *BMC research notes*. 2018;11(1):1-6.
- Csuka M, McCarty DJ. Simple method for measurement of lower extremity muscle strength. *Am J Med*. Jan 1985;78(1):77-81.
- Ng S. Balance ability, not muscle strength and exercise endurance, determines the performance of hemiparetic subjects on the timed-sit-to-stand test. *Am J Phys Med Rehabil*. Jun 2010;89(6):497-504.
- Melo TA, Duarte ACM, Bezerra TS, França F, Soares NS, Brito D. The Five Times Sit-to-Stand Test: safety and reliability with older intensive care unit patients at discharge. *Rev Bras Ter Intensiva*. 2019;31(1):27-33.
- Whitney SL, Wrisley DM, Marchetti GF, Gee MA, Redfern MS, Furman JM. Clinical measurement of sit-to-stand performance in people with balance disorders: validity of data for the Five-Times-Sit-to-Stand Test. *Phys Ther*. Oct 2005;85(10):1034-1045.
- Yenişehir S, Çıtak Karakaya İ, Sivaslıoğlu AA, Özen Oruk D, Karakaya MG. Reliability and validity of Five Times Sit to Stand Test in pregnancy-related pelvic girdle pain. *Musculoskelet Sci Pract*. Aug 2020;48:102157.
- Özden F, Coşkun G, Bakırhan S. The test-retest reliability and concurrent validity of the five times sit to stand test and step test in older adults with total hip arthroplasty. *Exp Gerontol*. Dec 2020;142:111143.
- Duncan RP, Leddy AL, Earhart GM. Five times sit-to-stand test performance in Parkinson's disease. *Arch Phys Med Rehabil*. Sep 2011;92(9):1431-1436.
- Mong Y, Teo TW, Ng SS. 5-repetition sit-to-stand test in subjects with chronic stroke: reliability and validity. *Arch Phys Med Rehabil*. Mar 2010;91(3):407-413.
- Özüdoğru A, Canlı M, Gürses ÖA, Alkan H, Yetiş A. Determination of five times-sit-to-stand test performance in patients with multiple sclerosis: validity and reliability. *Somatosensory & Motor Research*. 2023;40(2):72-77
- Özüdoğru A, Canlı M, Ceylan İ, Kuzu Ş, Alkan H, Karaçay BÇ. Five Times Sit-to-Stand Test in people with non-specific chronic low back pain—a cross-sectional test–retest reliability study. *Irish Journal of Medical Science (1971-)*. 2023;192(4):1903-1908.
- Lexell JE, Downham DY. How to assess the reliability of measurements in rehabilitation. *American journal of physical medicine & rehabilitation*. 2005;84(9):719-723.
- Guralnik JM, Simonsick EM, Ferrucci L, et al. A short physical performance battery assessing lower extremity function: association with self-reported disability and prediction of mortality and nursing home admission. *J Gerontol*. Mar 1994;49(2):M85-94.

23. Cachupe WJ, Shifflett B, Kahanov L, Wughalter EH. Reliability of biodex balance system measures. *Measurement in physical education and exercise science*. 2001;5(2):97-108.
24. Lee SC, Wu LC, Chiang SL, et al. Validating the Capability for Measuring Age-Related Changes in Grip-Force Strength Using a Digital Hand-Held Dynamometer in Healthy Young and Elderly Adults. *Biomed Res Int*. 2020;2020:6936879.
25. Celik D, Coşkunsu D, Kiliçoğlu O. Translation and cultural adaptation of the Turkish Lysholm knee scale: ease of use, validity, and reliability. *Clin Orthop Relat Res*. Aug 2013;471(8):2602-2610.
26. Briggs KK, Kocher MS, Rodkey WG, Steadman JR. Reliability, validity, and responsiveness of the Lysholm knee score and Tegner activity scale for patients with meniscal injury of the knee. *JBJS*. 2006;88(4):698-705.
27. Briggs KK, Lysholm J, Tegner Y, Rodkey WG, Kocher MS, Steadman JR. The reliability, validity, and responsiveness of the Lysholm score and Tegner activity scale for anterior cruciate ligament injuries of the knee: 25 years later. *Am J Sports Med*. May 2009;37(5):890-897.
28. Koo TK, Li MY. A Guideline of Selecting and Reporting Intraclass Correlation Coefficients for Reliability Research. *J Chiropr Med*. Jun 2016;15(2):155-163.
29. Schober P, Boer C, Schwart LA. Correlation Coefficients: Appropriate Use and Interpretation. *Anesth Analg*. May 2018;126(5):1763-1768.
30. Teo TW, Mong Y, Ng SS. The repetitive Five-Times-Sit-To-Stand test: its reliability in older adults. *International Journal of Therapy and Rehabilitation*. 2013;20(3):122-130.
31. Mong Y, Teo TW, Ng SS. 5-repetition sit-to-stand test in subjects with chronic stroke: reliability and validity. *Archives of physical medicine and rehabilitation*. 2010;91(3):407-413.
32. Yenişehir S, Karakaya İÇ, Sivaslıoğlu AA, Oruk DÖ, Karakaya MG. Reliability and validity of Five Times Sit to Stand Test in pregnancy-related pelvic girdle pain. *Musculoskeletal Science and Practice*. 2020;48:102157.
33. Snyder-Mackler L, Delitto A, Bailey SL, Stralka SW. Strength of the quadriceps femoris muscle and functional recovery after reconstruction of the anterior cruciate ligament. A prospective, randomized clinical trial of electrical stimulation. *J Bone Joint Surg Am*. Aug 1995;77(8):1166-1173.
34. Krishnan C, Williams GN. Factors explaining chronic knee extensor strength deficits after ACL reconstruction. *J Orthop Res*. May 2011;29(5):633-640.
35. Noehren B, Andersen A, Hardy P, et al. Cellular and Morphological Alterations in the Vastus Lateralis Muscle as the Result of ACL Injury and Reconstruction. *J Bone Joint Surg Am*. Sep 21 2016;98(18):1541-1547.
36. Ageberg E, Thomeé R, Neeter C, Silbernagel KG, Roos EM. Muscle strength and functional performance in patients with anterior cruciate ligament injury treated with training and surgical reconstruction or training only: a two to five-year followup. *Arthritis Rheum*. Dec 15 2008;59(12):1773-1779.
37. Konishi Y, Aihara Y, Sakai M, Ogawa G, Fukubayashi T. Gamma loop dysfunction in the quadriceps femoris of patients who underwent anterior cruciate ligament reconstruction remains bilaterally. *Scand J Med Sci Sports*. Aug 2007;17(4):393-399.
38. Zouita Ben Moussa A, Zouita S, Dziri C, Ben Salah FZ. Single-leg assessment of postural stability and knee functional outcome two years after anterior cruciate ligament reconstruction. *Ann Phys Rehabil Med*. Jul 2009;52(6):475-484.
39. Bonfim TR, Grossi DB, Paccola CAJ, Barela JA. Effect of additional sensory information in the proprioception and postural control of individuals with ACL lesion. *Acta Ortopédica Brasileira*. 2009;17:291-296.
40. Tookuni KS, Bolliger Neto R, Pereira CAM, Souza DRd, Greve JMDA, Ayala ADA. Análise comparativa do controle postural de indivíduos com e sem lesão do ligamento cruzado anterior do joelho. *Acta Ortopédica Brasileira*. 2005;13:115-119.
41. Tiwari D, Talley SA, Alsalaheen B, Goldberg A. Strength of association between the Five-Times-Sit-to-Stand Test and balance, knee extensor strength and lower limb power in community-dwelling older adults. *International Journal of Therapy And Rehabilitation*. 2019;26(3):1-10.
42. Goldberg A, Chavis M, Watkins J, Wilson T. The five-times-sit-to-stand test: validity, reliability and detectable change in older females. *Aging clinical and experimental research*. 2012;24(4):339-344.
43. Medina-Mirapeix F, Vivo-Fernández I, López-Cañizares J, García-Vidal JA, Benítez-Martínez JC, del Baño-Aledo ME. Five times sit-to-stand test in subjects with total knee replacement: Reliability and relationship with functional mobility tests. *Gait & posture*. 2018;59:258-260.