

**A comparative study on preservation of soy flour (*Glycine max.(L.) Merr.*) using sodium benzoate, a chemical preservative and *Aframomum melegueta* (grains of paradise), a natural preservative.**

**\*De, N. and \*\*Daniyan, S.**

\*Department of Microbiology  
Federal University of Technology, Yola.

**Department of Biological Sciences  
Federal University of Technology, Yola**

\* To whom all correspondence should be addressed

### **Abstract**

This investigation has explored the possibility of using ethanolic extract of seeds of *A. melegueta* at concentration of 1 mg/10g soy flour and sodium benzoate at a concentration of 1 mg/10g soy flour for preservation of soy flour over a period of sixty days. It has been shown that the pH of the samples treated with sodium benzoate was in the range of 6.50-6.80 whereas for samples treated with seeds of *A. melegueta* was in the range of 6.62- 6.86 over sixty days period of storage. The pH of untreated samples was in the range of 6.40-6.55 over the periods of storage. The bacterial and fungal isolates identified on 60<sup>th</sup> day of storage from the untreated samples were *Staphylococcus aureus*, *Bacillus subtilis*, *Escherichia coli*, *Aspergillus* sp., *Penicillium* sp., *Mucor* sp., *Rhizopus* sp. and *Candida* sp. whereas the organisms isolated from the samples treated with sodium benzoate were found to be *B. subtilis*, *B.cereus*, *Candida* sp. and *Aspergillus* sp. The organisms isolated from samples treated with extract of seeds of *A. melegueta* were *B. subtilis*, *B.cereus*, *S.aureus*, *Rhizopus* sp. and *Aspergillus* SP. The total bacterial counts for the untreated samples, samples treated with sodium benzoate and samples treated with seeds of *A. melegueta* were  $9.9 \times 10^{10}$ ,  $7.5 \times 10^8$  and  $6.0 \times 10^8$  respectively on 60<sup>th</sup> day of storage. The total fungal count for untreated samples, samples treated with sodium benzoate and samples treated with seeds of *A. melegueta* were  $9.9 \times 10^{10}$ ,  $9.0 \times 10^6$  and negligible. The protein content (25.1%) and the fat content (18.0%) for untreated samples were less than those of samples treated with sodium benzoate (32.0% and 19.0%) at 1.0 mg/10g soy flour and samples treated with seeds of *A. melegueta* (35.0% and 19.2%) at 1.0 mg/10 g soy flour.

**Key words:** soy flour, preservative, storage, protein and fat content.

### **Introduction**

Flour is generally regarded as a microbiologically safe product and it is a low water activity commodity. Although the growth of pathogenic bacteria may not be supported under such conditions, pathogens that contaminate flour may survive for extended periods. There are few reported incidents of food poisoning resulting from contaminated flour. Australian, European and United States of America studies indicate that *Salmonella* sp., *Escherichia coli*, *Bacillus cereus* and spoilage microorganisms are present in wheat flour at low levels (Cicognani *et al.*, 1975; Ottogali and Galli, 1979; Richter *et al.*, 1993). In 1952, an outbreak of salmonellosis caused by *Salmonella paratyphi* B phage type I occurred in New South Wales, Australia where flour was implicated, but the organism was never isolated from the suspected flour (Dack, 1961).

Soybean is a good source of lecithin and it is widely used in making pastries and sauces. Soy flour has many applications worldwide namely Com-soy milk (CSM), wheat soy blend and soy sauce (Obatolu *et al.*, 1993).

There is little information on the nutritional and microbial changes of soy flour during storage and its preservation using preservatives. Many spices have been reported to possess antimicrobial properties and have been successfully used as preservatives (Shelef, 1983). *Aframomum melegueta* Roscoc (Hausa name: Chita mai koko) is a perennial herb cultivated in tropical Africa and the seeds are used as a condiment or spice. It belongs to the family of Zingiberaceae. Ginerols, shagaols and paradols have been isolated from the grains of *A. melegueta*. The crude extracts showed considerable bactericidal activities against *E. coli*, *P.aeruginosa*, *B. subtilis*, *P. vulgaris*, *K. pneumoniae* and *S. marcescens* and fungicidal activities against *C. albicans*, *T. mentagrophytes*, and *A. niger* (Oloke and Kolawale, 1987). In this study, an attempt was made to preserve soy flour using extracts of seeds of *A. melegueta* and compare the preservative effect of seeds of *A. melegueta* of paradise with that of sodium benzoate, a chemical preservative.

## Materials and Methods

### Collection of materials

One thousand grams of soybean seeds, TGX 536-021, was collected from the Crop Production Department in the School of Agricultural Engineering, Federal University of Technology, Minna, Nigeria. The seeds of *A. melegueta* were purchased from Minna main market and were sun dried and kept in Laboratory cabinet for further use. Sodium benzoate was purchased from BDH (England).

### Preparation of materials

#### (A) Soy flour and Sodium benzoate

Preparation of soy flour from soy seeds started with cleaning and scouring of seeds to separate and remove non-seed material. Then the seeds were washed and oven dried at 55°C for 24 hours. The dried seeds were then ground into fine powder using a milling machine as described by Konan and Agbo, 1997. After sieving, the powdered soybean was kept in a sterile container for further experiments.

#### (B) Extract of seeds of *A. melegueta*

Dried seeds were ground into powder using an electric blender. Then 100 g of dried powdered samples was extracted with 400 ml of ethanol in a 2 litre conical flask for 24 hours. The extract was then recovered by filtration using Whatman no. 1 filter paper. A rotary evaporator was used, in vacuo, at 40°C to concentrate the extract. The dried extract (12.1 g/100 g of grains of paradise) was used for preservation purpose.

### Treatment of soy flour with sodium benzoate

Fifty milligrams of sodium benzoate was dissolved in 5 ml of acetone (concentration 10 mg/ml) for this purpose. One milliliter of this solution was added to 100 g of soy flour (concentration 1 mg/10g) and mixed with the flour using a sterile spoon. Soy flour so

prepared was distributed in 10 polythene bags each containing 10 g and the bags containing the samples were sealed and stored at room temperature for 60 days. The samples were withdrawn at regular intervals for microbiological and biochemical analysis.

#### *Treatment of soy flour with the seed extract*

Thirty milligrams of extract of seeds of *A. melegueta* was dissolved in 3 ml of acetone to obtain a stock solution of 10-mg/ml concentrations. To 100 g of soy flour 1 ml of this solution was added to get a concentration of 1 mg/10g soybean flour and mixed with the flour vigorously using a sterile spoon. The treated soy bean flour was dispensed in 10 polythene bags each containing 10 g and were sealed with an electric sealer and kept for 60 days at room temperature. Soy flour without any extract was used as control and also kept for 60 days for comparison. The samples were withdrawn at regular intervals for analysis.

#### **Quality assessment of treated and untreated samples**

##### (a) Determination of pH

One gram of each of the treated and untreated samples was added to 10 ml distilled water and after vigorous shaking, the pH was measured using a pH meter (Micro pH 3310 Crison).

##### (b) Isolation and enumeration of fungal and bacterial isolates

Each of the samples (0.1 g) was added to 9.9 ml of distilled water and using this as a stock solution, serial dilution up to  $10^{-6}$  were made following the procedure of Fawole and Oso (1988). An aliquot (0.1ml) of each dilution was introduced onto agar medium (for bacteria, Nutrient agar, NA and for fungi, potato dextrose agar, PDA, was used) and the plates were incubated at 28°C and 37°C for PDA and NA plates respectively. The bacterial isolates were identified using the procedure of Hudson and Sherwood (1997) and fungal isolates were identified using the procedure developed by Smith (1977).

##### (c) Determination of moisture, fat and crude protein content

The moisture content (% of flour) was determined using an electric protimeter grain master 2000. The fat content (%) of the treated and untreated soy flour was determined by soxhlet extraction using petroleum ether as solvent. Two grams of soy flour for each sample was used for this purpose (A.O.A.C, 1980). The total nitrogen content of the samples were determined by Kjeldahl method (Bermner, 1965) and then the crude protein content of soy flour was determined by multiplying the total nitrogen content by a factor of 6.25. Two hundred fifty milligrams of soy flour was used for determination of crude protein.

## **Results**

### *Quality assessment of treated and untreated samples*

#### (a) Determination of pH

For sodium benzoate treated samples and for seeds of *A. melegueta* treated samples, the pH range was 6.50-6.80 and 6.60-6.90 respectively whereas for the untreated samples the range was 6.35 to 6.65.

#### (b) Identification of isolates

The predominant bacterial isolates on 0 day of storage were *B. subtilis* (40%) followed by *S. aureus* (35%) and *B. cereus* (20%). The relative distribution of bacterial isolates on 60<sup>th</sup> day of storage for control samples were *B. cereus* (70%) followed by *B. subtilis* (20%), *S. aureus* (6%) and *E. coli* (4%). For all the treated samples, the predominant bacterial isolates were *B. cereus* (80-90%) followed by *S. aureus* (5-10%) and *B. subtilis* (3-5 %). The relative distribution of fungal isolates in soy flour on 0 day of storage and 60<sup>th</sup> day of storage was *Aspergillus* sp. 20- 30%, *Penicillium* sp. 30%-32%, *Mucor* species 20-30%, *Rhizopus* sp. 10%-15% and *Candida* sp. 5%-10%. For the samples treated with sodium benzoate and seeds of *A. melegueta*, the predominant bacterial isolates were *B. cereus* followed by *S. aureus* and *B. subtilis*. The predominant fungal isolates for sodium benzoate treated samples were *Candida* sp. followed by *Aspergillus* sp.

### (c) Enumeration

For control samples, the initial bacterial count was  $1.1 \times 10^{10}$  and on 60<sup>th</sup> day of storage the count was  $9.9 \times 10^{10}$ . For sodium benzoate treated samples the bacterial count and the fungal count were  $7.5 \times 10^8$  and  $9.0 \times 10^6$  respectively on 60th day of storage. The bacterial count for grains of paradise treated samples was  $6.0 \times 10^8$ . The results are shown in Table 1.

Table 1: Total number of bacterial and fungal isolates in untreated (UTS) and sodium benzoate (SB) and seeds of *A. melegueta* (AF) treated samples (mean± SEM; n=3).

Time (Day)	UTS		SB		AF	
	B	F	B	F	B	F
0	$1.1 \times 10^8 \pm 0.3$	$1.9 \times 10^8 \pm 0.2$	$1.1 \times 10^8 \pm 0.1$	$1.9 \times 10^8 \pm 0.2$	$1.1 \times 10^8 \pm 0.1$	$1.9 \times 10^8 \pm 0.2$
7	$1.4 \times 10^8 \pm 0.3$	$3.2 \times 10^8 \pm 0.3$	$9.0 \times 10^8 \pm 0.1$	$2.1 \times 10^7 \pm 0.2$	$9.5 \times 10^8 \pm 0.1$	$4.0 \times 10^6 \pm 0.3$
14	$9.0 \times 10^{10} \pm 0.3$	$4.3 \times 10^{10} \pm 0.2$	$8.7 \times 10^8 \pm 0.3$	$1.7 \times 10^7 \pm 0.2$	$8.3 \times 10^8 \pm 0.1$	NG
30	$9.0 \times 10^{10} \pm 0.2$	$5.1 \times 10^{10} \pm 0.2$	$7.7 \times 10^8 \pm 0.2$	$1.1 \times 10^7 \pm 0.2$	$7.5 \times 10^8 \pm 0.2$	NG
60	$9.9 \times 10^{10} \pm 0.2$	$9.9 \times 10^{10} \pm 0.1$	$7.5 \times 10^8 \pm 0.2$	$9.0 \times 10^6 \pm 0.3$	$6.0 \times 10^8 \pm 0.1$	NG

NG-negligible; B- bacterial count (cfu/g); F- fungal count (cfu/g)

(d) Determination of moisture, fat and crude protein content of treated and untreated samples.

The initial moisture count of the samples of soy flour was 17.6%. The moisture content of treated and untreated samples of soy flour on 7<sup>th</sup>, 14<sup>th</sup>, 30<sup>th</sup> and 60<sup>th</sup> day of storage are listed in Table 2.

**Table 2: Determination of moisture content (%) and fat content (%) of treated and untreated samples (mean± SEM; n=3)..**

Time (Day)	UTS		SB		AF	
	F	M	F	M	F	M
0	20.0±0.1	16.9±0.1	20.0±0.1	16.9±0.1	20.0±0.1	16.9±0.1
7	19.4±0.2	16.0±0.0	19.6±0.1	17.5±0.2	19.2±0.2	14.2±0.0
14	19.0±0.2	16.1±0.1	19.6±0.2	15.3±0.1	19.2±0.1	15.2±0.1
30	19.8±0.1	16.1±0.1	19.2±0.1	14.2±0.0	19.8±0.0	16.7±0.0
60	18.0±0.0	16.3±0.0	19.2±0.2	14.1±0.1	19.2±0.2	16.8±0.1

UTS- untreated samples; SB- sodium benzoate treated samples; AF- seeds of *A. melegueta* treated samples; F- fat content (%); M-moisture content (%)

The fat content of soy flour samples on 0 day was 20.00 (%). The fat content (Table 2) for treated and untreated samples remained almost constant.

The mean reduction in crude protein (%) /day for the untreated samples was 0.34 whereas for the sodium benzoate treated samples and seeds of *A. melegueta* samples the respective values were 0.21 and 0.15. The results are shown in Table 3.



## Discussion

Whole soy flour used in this study contains 40% protein and 20% fat (dry weight). This agrees with the crude protein (42 %) and fat content (20 %) of the whole soy flour prepared by Bressani (1981). This study revealed that there are three types of bacteria present in soy flour namely *B. cereus*, *S. aureus* and *B. subtilis*. The fungi isolated on 0 day of storage were species of *Aspergillus*, *Penicillium*, *Mucor*, *Rhizopus* and *Candida*. The initial bacterial count was  $1.1 \times 10^8$  whereas the count was  $9.9 \times 10^{10}$  on 60 days of storage for control samples. Microbiological guidelines for flour have been proposed for various countries (Potus and Suchet, 1989; Richter et al., 1993). A survey by Richter et al. (1993) found US flour contained mean counts of  $10^3$ - $10^4$  cfu/g, depending on wheat type. Spicher (1986) reported German flour contained mean counts of  $10^4$  cfu/g, coliform counts of  $10^2$  cfu/g and mould counts of  $10^3$  cfu/g. Potus and Suchet (1989) detected  $10^4$  cfu/g total aerobes and  $10^3$  cfu/g moulds in French flour. Berghofer *et al.* (2003) detected the presence of *B. cereus* in Australian wheat flour but at low levels. The initial fungal count was  $1.95 \times 10^{10}$  and on 60<sup>th</sup> day of storage the count was  $9.90 \times 10^{10}$ . Some workers claim that 15% or above moisture permits good fungal growth. Aflatoxin is not considered a problem in soybean storage. Hesseltine *et al.* (1966) were unable to detect aflatoxin on soybeans inoculated with *A. flavus* although Farag *et al.* (1986) detected aflatoxin in sterilized and non sterilized soybeans inoculated with *A. parasiticus*. *E. coli* was detected in untreated soy flour samples on 60th day of storage. Graves *et al.* (1967) found *E. coli* in only 1 of 16 US flour samples but Richter *et al.* (1993) reported 12.8% of US flour contained *E. coli*.

Sodium benzoate was selected to preserve soy flour because it is very effective at acidic pH and it has been used extensively as an antimicrobial agent in foods (the maximum level of benzoic acid permitted in food is 0.1%). It has been shown that the seed extract inhibited the growth of *B. theobromae*, *A. niger*, species of *Mucor*, *Rhizopus*, *Fusarium* and *A. flavus* (Oloke and Kolawole, 1987). The mean reduction in crude protein (%) for the control over 60 days of storage was 0.34 whereas for sodium benzoate it was 0.21 whereas the value was 0.15 for seed extract treated samples. For seed extract treated samples the fungal count is negligible but the bacterial count was in the order of  $10^8$ . This value is high compared to the standard of quality flour set by Berghofer *et al.* (2003). *A. melegueta* may be a useful



preservative for soy flour though the presence of *B. cereus* is a cause of concern because this might cause food poisoning and it also can withstand heat (Blakey and Priest, 1980).

## References

- A.O.A.C. (1980). Official Methods of Analysis 13<sup>th</sup> Edition. Association of Official Analytical Chemists Inc., Washington, D.C.
- Blakey, L.J. and Priest, F.G. (1980). The occurrence of *Bacillus cereus* in some dried foods including pulse and cereals. *Journal of Applied Bacteriology* **48**: 297-302.
- Berghofer, L.K.; Hocking, A.D; Miskelly, D. and Jansson, E. (2003). Microbiology of wheat and flour milling in Australia. *International Journal of Food Microbiology* **85**: 137-149.
- Bermner, J. M. (1965). Total nitrogen In: Methods of Soil Analysis. American Soc. Agron. Monograph No.9, 1149-1176.
- Bressani, R. (1981). The role of soybeans. *Journal of American Oil Chemists' Society* **58**: 392-400.
- Cigonani, G.; Pedretti, C. and Cerrato, A. (1975). Microbiological characteristics of wheat flour. *Industrie Alimentari* **14** (7/8): 60-64.
- Dack, G. M.(1961). Public health significance of flour bacteriology. *Cereal Science Today* **6**: 9-10.
- Farag, R. S.; El-Leithy, M. A.; Basyony, A. E. and Daw, Z. Y. (1986). Effects of varied substrates on aflatoxin production by *Aspergillus parasiticus*. *Journal of American Oil Chemists' Society* **63**: 1024-1026.
- Fawole, M. O. and Oso, B. A. (1988). Laboratory Manual of Microbiology. Eds. Spectrum Books, Ibadan, pp. 24-56.
- Graves, R. R.; Rogers, R. F.; Lyons, A. J. and Hesseltine, C. W. (1967). Bacterial and actinomycete flora of Kansa- Nebraska and Pacific North-West wheat and wheat flour. *Cereal Chemistry* **44**: 288-299.
- Hesseltine, C. W.; Shotwell, O. L.; Ellis, J. J. and Stubblefield, R. D. (1966). Aflatoxin formation by *Aspergillus flavus*. *Bacteriology Reviews* **30**: 795-805.
- Hudson, K. B. and Sherwood, L. (1997). Identifying bacteria using metabolic characteristics In: Exploitations in Microbiology. Prentice Hall, New Jersey, pp. 83-90.
- Konan, G. and Agbo, N. G. (1997). Soybean supplementation of maize and millet pap. Soybean utilization in Nigeria, Ghana and Cote D'Ivoire (West Africa). IDRIC IITA Soybean utilization project phase III P. 42049.
- Obatolu, V. A.; Osho, S. M. and Oyekan, P. M. (1993). Fortification of indigenous foods using soybeans (For the south western States of Nigeria). IITA and IARST Publication. p.41.
- Oloke, J. K. and Kolawole, D. O. (1987). The antibacterial and antifungal activities of certain components of *Aframomum melegueta* fruits. *Fitoterapia Lix*. **5**: 384-388.
- Ottogalli, G. and Gali, A. (1979). Microbiological quality of flours: sour dough for bakery products and spaghetti.** In: Jarvia, B.; Christian, J. H. B. and Michener, H. (eds.) Food Microbiology and Technology. Proceedings of the International Meeting on Food Microbiology and Technology, Tabiabo (Parma), Italy, pp. 141-153.

- Potus, J. and Suchet, P. (1989). Microbiological problems in milling. *Industries des Cereales* **58**: 27-33.
- Richter, K. S.; Dorneanu, E.; Eskridge, K. M. and Rao, C. S. (1993). Microbiological quality of flours. *Cereal Foods World* **38**: 367-369.
- Shelef, L.A. (1983). Antimicrobial effects of spices. *Journal of Food Safety* **6**: 29-44.
- Spicher, G. (1986) Judging the microbiological-hygienic quality of white flour. *Die Muhle & Mischfuttertechnik*. **33**:449.
- Smith, D. A. (1977). Enumerating fungi. *Phytopathology* **8**: 81.