

Erratum

Erratum: Complications of Arteriovenous Fistula with Polytetrafluoroethylen Grafts in Hemodialysis Patients

In the article titled “Complications of arteriovenous fistula with polytetrafluoroethylen grafts in hemodialysis patients” published in pages 120-123, issue 1, vol. 18 of Nigerian Journal of Clinical Practice^[1], the references 8, 9 and 10 in the reference list were written incorrectly.

The correct references are following:

Reference 8: Mickley V. Personal Opinion Stenosis and thrombosis in haemodialysis fistulae and grafts: The surgeon’s point of view. *Nephrol Dial Transpl.* 2004;19:309-11.

Reference 9: Odabaşı D, Arı E, Kıymaz A, Ekim H. Should we use saphenous vein graft instead of synthetic graft for creation of secondary arteriovenous fistula in hemodialysis dependent end stage renal failure patients? *Turkish J. Thorac. Cardiovasc. Surg.* 2012;16;20(2):209-16.

Reference 10: Basel H, Cegin B, Aydın U, Aydın C, Kutlu H, Keradag M, *et al.* Dialysis purpose of the operation of the arterio-venous fistulas effect of the areas in open fistulas. *Van Tıp Derg* 2010;17:118-23.

REFERENCE

1. Göksin I, Baltalarlı A, Önem G, Rendeci O, Saçar M, Kara H, *et al.* Arteriovenous fistula operations: Its early and late-term complications that need to revision. *Turk J Thorac Cardiovasc Surg* 2004;12:180-3.

DOI: 10.4103/1119-3077.212572

