

Kinetics Study of the Reduction of $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ by Thiocyanate Ion in Aqueous Hydrochloric Acid.

Umoru, P. E.

Department of Chemistry, Nigerian Defence Academy, Kaduna, Nigeria
Email: patumoru@gmail.com

ABSTRACT

The kinetics of the reduction of $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ by thiocyanate ion (SCN) has been studied in aqueous hydrochloric acid medium. The stoichiometric determination showed a mole ratio of 1:3. Kinetics study was carried out under pseudo - first order condition and showed that the rate of reaction was first order with respect to both $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ and [SCN]. The rate of the reaction was acid dependent and was found to decrease with increase in ionic strength, suggesting a negative salt effect. The addition of added cations and anions (Ni^{2+} , Mg^{2+} , NO_3^- and CH_3COO^-) increased the rate of reaction and Michaelis-Menten's plot showed a zero intercept. Based on the results of the spectroscopic studies, kinetic evidences from Michaelis-Menten's plot and interaction with added ions, an outer-sphere mechanism has been proposed for the reaction of $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ - SCN system.

Keywords: Kinetics, Reaction, $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$, Thiocyanate Ion.

INTRODUCTION

Manganese has been reported in literature to have a very important role in the active site of organisms. It is now clearly established that at least four enzymes (the photosynthetic oxygen-evolving complex, manganese catalase, manganese superoxide dismutase and ribonucleotide reductase) use bi- or multi-nuclear manganese complex as their redox catalytic center¹; Vincent *et al.*,¹; Andreasson and Vanngard²; Penner³.

Thiocyanates (SCN) has antioxidant properties that include the ability to protect cells against oxidizing agents such as hypochlorous acid (HOCl) and repair protein chloramines. SCN is an important endogenous molecule that has the potential

to interact in complex and elegant ways with its host environment and foreign organisms. SCN's diverse properties as both host defense and antioxidant agent make it a potentially useful therapeutic⁴. Chandler and Day⁴. They are found in various foods and plants; they are produced primarily from the reaction of free cyanide with sulphur⁵. Pedemonte *et al.*,⁵. It is sometimes called sulphocyanate or simply "yam vitamin" a natural substance produced in certain plants called nitrilosides among which are the foods, African yams and cassava. These edible plants were once included in the diet of native Africans before the introduction of grains such as rice and wheat. It is also known to be an important part in

biosynthesis, thus the complete absence of thiocyanate according to “Childers *et al.*,⁶ or reduced thiocyanate as reported by “Minarowski *et al.*,⁷ in the human body is damaging to the human host defence system“Moskwa *et al.*,⁸ ; “Xu *et al.*,⁹ . It is used as a precursor for the synthesis of pharmaceuticals and other specialty chemicals. The role of manganese as well as that of thiocyanate in humans has therefore prompted the investigation of the kinetics and mechanism of the reduction of $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ (referred to in the text as $[Mn^{III}O_2Mn^{IV}]^{3+}$ by (SCN^-) .

MATERIALS AND METHODS

Materials

Analytical grade reagents were used throughout this work without further purification. Hydrochloric acid (HCl) was used to maintain $[H^+]$ for the reactions while Sodium chloride (NaCl) was used to maintain constant ionic strength. Magnesium chloride ($MgCl_2$), Nickel nitrate ($NiNO_3$), Sodium acetate (CH_3COO^-) and Sodium nitrate (NO_3^-) were used to test for the effect of added species on reaction rate. Acrylamide with methanol were used for free radical tests in all the reactions.

Stoichiometric Determination

The stoichiometry of the reaction was determined by spectrophotometric titration using the mole ratio approach. The concentration of the $[Mn^{III}O_2Mn^{IV}]^{3+}$ was kept constant at $5.0 \times 10^{-3} \text{ mol dm}^{-3}$ and that of SCN^- was varied between $(3.0 - 12.0) \times 10^{-3} \text{ mol dm}^{-3}$ at $[H^+] = 1 \times 10^{-3} \text{ mol dm}^{-3}$, $I =$

0.5 mol dm^{-3} , $T = 30.0 \pm 1.0^\circ\text{C}$. The reaction mixtures were monitored at $\lambda_{\text{max}} = 683 \text{ nm}$ and the absorbance (A_∞) was plotted against the mole ratio of the reactants and the point of sharp inflection in the plot gave the stoichiometry of the reaction.

Kinetic Measurements

All kinetic measurements were performed under pseudo-first order conditions with that of SCN^- in at least 10 folds excess over that of the complex. From the logarithmic plot of $\log (A_t - A_\infty)$ against time (t) in seconds (where A_t and A_∞ are the absorbance at time, t and at infinity respectively), the slope of the plot was obtained and the pseudo-first order rate constant (k_1) was determined. The second order rate constant (k_2) was subsequently estimated from the relation:

$$k_2 = k_1/[SCN^-]^n,$$

where 'n' = reaction order with respect to SCN^- .

Effect of Hydrogen Ion Concentration

The effect of changes in $[H^+]$ on the reaction rate was investigated at constant concentration of $[(bipy)_2Mn^{III}O_2Mn^{IV}[(bipy)_2]^{3+}$ and SCN^- . $[H^+]$ was varied between $(0.4 - 2.8) \times 10^{-3} \text{ mol dm}^{-3}$ at constant ionic strength of 0.5 mol dm^{-3} (NaCl) and Temperature at $30.0 \pm 1.0^\circ\text{C}$. The order of reaction with respect to $[H^+]$ was obtained as the slope of the plot of $\log k_1$ against $\log [H^+]$. Variation of acid dependent rate constant with $[H^+]$ was obtained by plotting k_2 against $[H^+]$.

Effect of Changes in Ionic Strength and Dielectric Constant of the Reaction Medium

The ionic strength of the reaction mixture, I , was varied between $(0.2 - 1.4) \text{ mol dm}^{-3}$ (NaCl) while maintaining the concentration of the complex and SCN^- constant at $30.0 \pm 1.0^\circ\text{C}$. The relationship of the reaction rate with changes in the ionic strength was determined by plotting $\log k_2$ vs \sqrt{I} . The effect of medium dielectric constant, D , on the rate was investigated by using a binary solvent mixture of water and acetone.

Effect of Added Ions on the Reaction Medium

The effect of added ions on the reaction rate was achieved by the addition of various amounts of ions (Mg^{2+} , Ni^+ , CH_3COO^- , NO_3^-) while maintaining the concentrations of the complex, SCN^- and HCl constant. The ionic strength and temperature were also kept constant.

Test for Free Radicals

About two (2cm^3) of acrylamide solution was added to a partially reduced reaction mixture, followed by an addition of methanol in excess. The same treatment was applied to the solutions of $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ and SCN^- separately which served as control. Addition of excess methanol did not give a gel, indicating the probable absence of free radicals in the reaction mechanism.

Tests for Intermediate Complex


Electronic spectra of partially reacted reaction mixtures were recorded at various time intervals. A similar run was carried out

for reactants separately in each case. This was carried out in order to determine whether enhancement of peak or shift in λ_{max} will result as the reaction progresses. Furthermore, Michaelis-Menten's plot of $1/k_1$ versus $1/[\text{SCN}^-]$ was plotted which provided information on the presence or absence of intermediate complex formation through the identification or non-identification of intercept.

RESULTS AND DISCUSSION

Stoichiometry Studies

A mole ratio of 1:3 was observed for the stoichiometry result between $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- . This is consistent with the report of "Lohdip and Iyun¹⁰ for the reaction of $[(\text{phen})_2\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}(\text{phen})_2]$ and SCN^- . Based on the stoichiometry observed, the overall equation for the redox reaction can be represented by the following equation:



Order of Reaction

The kinetics investigation indicated an order of one with respect to $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ from the plot of $\log (A_t - A_\infty)$ versus time which was linear over 80 % extent of the reaction (Fig. 1.0). The logarithmic plot of k_1 versus $\log [\text{SCN}^-]$ was linear with a gradient approximately equal to unity (Fig. 2.0), suggesting a first order with respect to SCN^- , thus a second order overall. A second order

rate constant, k_2 , was obtained from $k_2 = k_1/[\text{SCN}^-]$ and were found to be fairly constant for the system (Table 1.0), further confirming first order with respect to the $[\text{SCN}^-]$. The rate law for the reaction is presented as:

$$\frac{-d[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}}{dt} = k_2[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}[\text{SCN}^-] \quad \mathbf{2.0}$$

where $k_2 = 12.59 \pm 0.25 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$




Figure 1: Typical Pseudo-first Order Plot of $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- System.




Figure 2: Plot of $\log k_1$ vs $\log [\text{SCN}^-]$ for the Reaction of $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- System.

Table 1.0: Pseudo-first Order and Second Order Rate Constant for the Reaction of $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+} = 5 \times 10^{-3} \text{ mol/dm}^3$ by SCN^- , $[\text{H}^+] = 1.0 \times 10^{-3} \text{ mol dm}^{-3}$, $I = 0.5 \text{ mol dm}^{-3}$ at $30.0 \pm 1.0 \text{ }^\circ\text{C}$ and $\lambda_{\text{max}} = 683 \text{ nm}$.

$10^2 [\text{SCN}^-]$ mol dm^{-3}	$[\text{H}^+], 1.0 \times 10^{-3}$ mol dm^{-3}	$I, 0.5 \text{ mol dm}^{-3}$	$10^4 k_1, \text{ s}^{-1}$	$k_2, \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
5.0	1.0	0.5	6.17	12.34
6.0	1.0	0.5	7.66	12.77
7.0	1.0	0.5	8.90	12.71
8.0	1.0	0.5	10.34	12.93
9.0	1.0	0.5	11.29	12.54
10.0	1.0	0.5	12.68	12.68
11.0	1.0	0.5	13.83	12.57
12.0	1.0	0.5	14.59	12.16
13.0	1.0	0.5	16.04	12.34
8.0	0.4	0.5	5.75	7.19
8.0	0.8	0.5	6.95	8.69
8.0	1.2	0.5	8.08	10.10
8.0	1.6	0.5	9.21	11.51
8.0	2.0	0.5	10.34	12.92
8.0	2.4	0.5	11.44	14.30
8.0	2.8	0.5	12.64	15.80
8.0	1.0	0.2	11.37	14.21
8.0	1.0	0.4	10.32	12.90
8.0	1.0	0.6	9.61	12.01
8.0	1.0	0.8	8.65	10.82
8.0	1.0	1.0	8.47	10.59
8.0	1.0	1.2	7.43	9.92
8.0	1.0	1.4	7.02	8.78

Effect of Hydrogen Ion Concentration

The results for the effect of $[H^+]$ indicated that the rate of reaction increased with increase in the concentration of HCl (Fig. 3.0). This result is in good agreement with the reports of “Banerjee *et al.*,¹¹ for the reduction of $[Mn^{III}O_2Mn^{IV}]^{3+}$ and hydrogen peroxide, “Lohdip and Iyun¹⁰, for the reaction of $[(phen)_2Mn^{III}O_2Mn^{IV}(phen)_2]^{3+}$ with SCN^- and “Chaudhuri and Banerjee¹² for the reaction of $[Mn^{III}O_2Mn^{IV}]^{3+}$ with NO_2^- respectively. Hence

the overall equation for the reaction is presented as:

$$\frac{-1/3d[Mn^{III}O_2Mn^{IV}]^{3+}}{b[H^+][Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-]} = \quad (a + \quad) \quad \mathbf{3.0}$$

where $a = 5.805 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$, $b = 3.56 \times 10^{-3} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$: a = intercept and b = slope.




Figure 3.0: Plot of $\log k_2$ vs $[H^+]$ for $[Mn^{III}O_2Mn^{IV}]^{3+}$ - SCN^- System.

Effect of Changes in Ionic Strength and Dielectric Constant of the Reaction Medium

The rate constant was found to decrease with increase in ionic strength from a plot of $\log k_2$ vs \sqrt{I} (Fig. 4.0), suggesting a negative salt effect. This observation suggests that the activated complex is formed from two ions of unlike charges. This result is in good

agreement with the reports of “Arabel *et al.*,¹³ for the reaction of $[(bipy)_2Mn^{III}O_2Mn^{IV}(bipy)_2]^{3+}$ and hydroxyacids; “Leal *et al.*,¹⁴ for the reactions of hexacyanoferrate(III) and L-ascorbic acid and “Iyun *et al.*,¹⁵ for $[\text{diaquotetrakis}(2, 2' - \text{bipyridine}) - \mu - \text{oxodiruthenium(III)}]^{3+}$ and ascorbic acid respectively.




Figure 4.0: Plots of $\log k_2$ vs \sqrt{I} for $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- System.

The effect of change in dielectric constant showed that the rate constant increased as the concentration of acetone increased.

Effect of Added Ions in the Reaction Medium

The addition of added cations and anions (Mg^{2+} , Ni^{2+} , CH_3COO^- and NO_3^-) increased the rate of reaction (Table 2.0). The catalysis of the added cations and anions is consistent with the reports of "Lohdip and Iyun¹⁰ for the reaction of $[(\text{phen})_2\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}(\text{phen})_2]^{3+}$ and SCN^- . The cations and anions catalysis for this reaction indicates that the reaction did proceed via outer-sphere mechanism. This result is also consistent with the result reported by "Mohammed *et al.*¹⁶ for the reaction of $[(\text{bipy})_2(\text{H}_2\text{O})\text{RuORu}(\text{H}_2\text{O})(\text{bipy})_2]^{4+}$ and thiourea.

Test for Free Radicals

Addition of a solution of acrylamide to partially reacted mixture did not give a gel

even in the presence of excess methanol, indicating the probable absence of free radicals in the reaction mechanism.

Tests for Intermediate Complex

The results of spectroscopic studies indicated no shift from λ_{max} of 683 nm this suggests the absence of the formation of an intermediate complex in the reaction. Furthermore, the plot of $1/k_1$ versus $1/[\text{SCN}^-]$ (Fig. 5.0) gave a straight line which passed through the origin also suggesting absence of intermediate complex formation prior to redox reaction process. This result is also consistent with the results of "Lohdip and Iyun¹⁷," "Iyun *et al.*,¹⁸ for the reaction of (diaquotetrakis (2, 2' - bipyridine) - μ - oxodiruthenium(III))³⁺ and 2-mercaptoethanol and diAquotetrakis(2, 2' - bipyridine) - μ - OxoDiRuthenium(III)³⁺ and L- ascorbic acid respectively.

Table 2.0: Effect of Added Cations and Anions for the $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- System. $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+} = 5 \times 10^{-3} \text{ mol/dm}^3$, $[\text{H}^+] = 1.0 \times 10^{-3} \text{ mol dm}^{-3}$, $I = 0.5 \text{ mol dm}^{-3}$ at 30.0 ± 1.0 °C and $\lambda_{\text{max}} = 683 \text{ nm}$.

Ion	$10^3 [\text{Ion}], \text{ mol dm}^{-3}$	k_1, s^{-1}	$k_2, \text{dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$
Mg²⁺	0.2	0.97	12.13
	0.4	0.99	12.41
	0.6	1.01	12.71
	0.8	1.03	12.90
	1.0	1.05	13.11
	1.2	1.07	13.32
Ni²⁺	0.2	16.42	12.32
	0.4	16.56	12.42
	0.6	16.65	12.49
	0.8	16.88	12.66
	1.0	17.25	12.94
	1.2	17.36	13.02
CH₃COO⁻	0.2	0.94	11.73
	0.4	0.94	11.80
	0.6	0.99	12.38
	0.8	1.01	12.65
	1.0	1.04	12.99
	1.2	1.07	13.40
NO₃⁻	0.2	1.01	12.68
	0.4	1.03	12.83
	0.6	1.05	13.06
	0.8	1.06	13.25
	1.0	1.09	13.68
	1.2	1.12	14.04








Figure 5.0: Plot of $1/k_1$ versus $1/[\text{SCN}^-]$ for $[\text{Mn}^{\text{III}}\text{O}_2\text{Mn}^{\text{IV}}]^{3+}$ - SCN^- System.

Reaction Mechanism

The observed acid catalysis was due to successive protonation and hydration of the complex "Chaudhuri and Banerjee"¹² Under the

acid condition used for the system, the following reaction mechanism was proposed:





Equations 5 and 6 are the rate determine steps:

$$\text{Rate} = k_2[Mn^{III}O_2Mn^{IV}]^{4+}[SCN^-] + k_3[Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-] \quad 8.0$$

From equation 4

$$Mn^{III}O_2Mn^{IV}]^{4+} = K_1[Mn^{III}O_2Mn^{IV}]^{3+}[H^+] \quad 9.0$$

Substituting equation 9.0 into 8.0

$$\text{Rate} = k_2K_1[Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-][H^+] + k_3[Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-] \quad 10.0$$

Rearranging equation 10.0

$$\text{Rate} = k_3 + k_2K_1[H^+][Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-] \quad 11.0$$

where $k_3 = a$, $k_2K_1 = b$

Equation 11.0 becomes:

$$\text{Rate} = (a + b[H^+])[Mn^{III}O_2Mn^{IV}]^{3+}[SCN^-] \quad 12.0$$

where $a = 5.805 \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$, $b = 3.56 \times 10^{-3} \text{ dm}^3 \text{ mol}^{-1} \text{ s}^{-1}$

This equation is similar to the rate equation in 3.0 above.

CONCLUSION

The kinetics of the reduction of $[Mn^{III}O_2Mn^{IV}]^{3+}$ by thiocyanate ion (SCN^-) in aqueous hydrochloric acid medium showed a mole ratio of 1:3, order of reaction is first order with respect to the complex and thiocyanate ions respectively - second order overall. The rate of reaction increased with increase in hydrogen ion concentration, decreased with increase in ionic strength of the reaction medium. The rate of reaction was affected by added ions and an intermediate complex was not detected.

Based on the above results, the reaction system has been rationed to proceed via outer-sphere mechanism.

ACKNOWLEDGEMENT

I am grateful to Nigerian Defence Academy, Kaduna for providing all the facilities used for this work.

REFERENCES

1. Vincent, L. P., Michael, J. B., and Andrew, G., Interaction of Manganeses with Dioxygen and its Reduced Derivatives, Chem. Rev., 94 (1994) 807

2. Andreasson, L. E. and Vanngard, T., Manganese: Oxygen-evolving Complex & Models, in Encyclopedia of Inorganic Chemistry, Vol. 4 (1992) (ed. Bruce, K. R.), New York: John Wiley & Sons, 2101
3. Penner-Hahn, J. E. (1992). Manganese Proteins with Mono- & Dinuclear Sites, in Encyclopedia of Inorganic Chemistry, Vol. 4 (ed. Bruce, K. R.), New York: John Wiley & Sons, 1992, 2111
4. Chandler, J. D. and Day, B. J., Thiocyanate: A Potentially useful Therapeutic Agent with Host Defense and Antioxidant Properties, Biochem Pharmacol. 84 (2012) 1381
5. Pedemonte, N., Caci, E., Sondo, E., Caputo, A., Rhoden, K., Pfeffer, U., di Candia, M. Bandettini, R., Ravazzolo, R., Zegarra-Moran, O. and Galiotta, L. J., "Thiocyanate Transport in Resting and IL-4-Stimulated Human Bronchial Epithelial Cells: Role of Pendrin and Anion Channels"(pdf). Journal of Immunology 178 (2007) 5144
6. Childers, M., Eckel, G., Himmel, A. and Caldwell, J. "A new Model of Cystic Fibrosis Pathology: Lack of Transport of Glutathione and its Thiocyanate Conjugates". Medical Hypotheses, 68 (2007) 101.
7. Minarowski, Ł., Sands, D., Minarowska, A., Karwowska, A., Sulewska, A., Gacko, M. and Chyczewska, E., "Thiocyanate concentration in saliva of cystic fibrosis patients" (pdf). Folia Histochemica et Cytobiologica 46 (2008) 245
8. Moskwa, P., Lorentzen, D., Excoffon, K. J., Zabner, J., McCray, P. B. Jr., Nauseef, W. M., Dupuy, C. and Bánfi, B., "A Novel Host Defense System of Airways is Defective in Cystic Fibrosis" (pdf). American Journal of Respiratory and Critical Care Medicine 175 (2007) 174
9. Xu, Y., Szép, S., Lu, Z. and Szep, L., "The Antioxidant Role of Thiocyanate in The Pathogenesis of Cystic Fibrosis And Other Inflammation-Related Diseases"(pdf). Proceedings of the National Academy of Sciences of the United States of America 106 (2009) 20515
10. Lohdip, Y.N. and Iyun, J.F., Kinetics and Mechanism of the Oxidation of Thiocyanate Ion by Di-u-oxo-tetrakis (1, 10-Phenanthroline) – Dimanganese (III, IV) ion in Acid Medium, Global Journal of Pure and Applied Sciences, 9 (2003) 105
11. Banerjee, R., Mondal, B. and Kundu, S., Kinetics of Oxidation of Hydrogen Peroxide by Di-oxo-Bridged Manganese (III, IV) Complexes of (2,2' bipyridine and 1,10-Phenanthroline) . J. Chem. Soc., Dalton Trans. 15 (1997)4341
12. Chaudhuri, S., Mukhopadhyay, and Banerjee, R., Solution Equilibria and Redox Reactivities of di-oxo-bridged Manganese Complex. Journal of Chemical Society, Dalton Trans (1995) 621.
13. Arabel, J.M., Ayoko, G.A. and Singh, K. Kinetics of Oxidation α -hydroxyacids by High Valent Oxo-manganese Complexes, Unpublished Material, 8 (1997) 123
14. Leal JM, Domingo PL, Garcla B, Ibeas S., Alkali metal ion catalysis of the oxidation of L-ascorbic acid by hexacyanoferrate(III) in strongly acidic media. J. Chem. Soc. Faraday Trans. 89 (1993) 3571
15. Iyun, J.F., A.O. Peters and Babatunde O.A., Kinetics of the reduction of dioxotetrakis(1,10 phenanthroline)di manganese (III,IV) perchlorate by L-Ascorbic

acid in aqueous acid solution. J. Chem. Soc. Nig., 30 (2005)114.

16. Mohammed, Y, Idris, S. O. and Iyun, J. F., Redox Kinetics and Mechanism of the Oxidation of Thiourea by Diaquatetrakis(2,2'-Bipyridine) - μ - Oxodiruthenium (III) Ion, in Aqueous Perchloric Acid, International Research Journal of Pure and Applied Chemistry, 4 (2014)819

17. Lohdip, Y.N. and Iyun, J.F., Kinetics of the Reduction of Di-u-oxo-tetrakis (1, 10-Phenanthroline) – Dimanganese (III, IV) by DL- Methionine in Acid Solution. Bull Chem Soc.Ethip. 12 (1998) 113

18. Iyun J.F.; Ayoko, G.A, and Lawal H.M., Kinetics and Mechanism of the Reduction of diaquatetrakis(2,2'-bipyridine)- μ -oxo diruthenium (III) by Ascorbic Acid. Transition Metals Chemistry. 20 (1995) 30